

REFERENCES

- Abo - Khatwa, N. (1978). Cellulase of fungus - growing termites : a new hypothesis on its origin. *Experimentia* **34** : 559 - 560.
- Aregger, E. (1992). Conidial production of the Fungus *Beauveria brongniartii* on Barley and Quality Evaluation during storage at 2⁰C. *Journal of Invertebrate Pathology* **59** : 2 - 10.
- Barson, G.(1977). Laboratory evaluation of *Beauveria bassiana* as a pathogen of larval stage of large elm bark beetle, *Scolytus scolytus*. *Journal of Invertebrate Pathology* **29**: 261-266.
- Bentley, B.L. (1984). Nitrogen fixation in termites fate of newly fixed nitrogen. *Journal of Insect Physiology* **30** : 653 - 655.
- Bermudes, D., Chase, D., and Margulis, L. (1988). Morphology as a basis for taxonomy of large spirochetes symbiotic in wood - eating cockroaches and termites : *Pillotina* gen. nov., nom. rev.; *Pillotina calotermitidis* sp. nov., nom. rev.; *Diplocalyx* gen. nov., rev.; *Diplocalyx calotermitidis* sp. nov. nom. rev.; *Hollandina pterotermitides* sp. nov. rev.; and *Clevelandina reticulitermitidis* gen. nov., sp. nov. *International Journal of Systematic Bacteriology* **38** : 291 - 302.
- Bidochka M.J. and Khachatourians,G.G (1987). Purification and properties of an extra cellular protease produced by the entomopathogenic fungus *Beauveria bassiana*. *Applied Environmental Microbiology* **53**: 1679-1684.
- Bidochka M.J. and Khachatourians,G.G (1990). Identification of *Beauveria bassiana* extracellular protease as a virulence factor in pathogenicity toward the migratory grasshopper, *Melanoplus sanguinipes*. *Journal of Invertebrate Pathology* **56**: 362-370.
- Bignell, D.E. and Anderson, J.M. (1980). Determination of pH and oxygen status in the guts of lower and higher termites. *Journal of Insect Physiology* **26** : 183 - 188.

- Bignell, D.E., Oskarsson, H. and Anderson, J.M. (1980). Colonization of the epithelial face of the peritrophic membrane and the ectoperitrophic space by actinomycetes in a soil-feeding termite. *Journal of Invertebrate Pathology* **36** : 426 - 428.
- Bignell, D.E., Oskarsson, H. and Anderson, J.M. (1980a). Specialization of the hindgut wall for the attachment of symbiotic micro- organisms in a termite *Procupitermes aburiensis* (Isoptera, Termitidae, Termitinae). *Zoomorphology* **96** : 103 - 112.
- Bignell, D.E., Oskarsson, H., Anderson, J.M., Ineson, P. and Wood ,T.G. (1983). Structure, microbial associations and function of the so-called "mixed segment" of the gut in two soil-feeding termites, *Procupitermes aburiensis* and *Cubitermes severus* (Termitidae, Termitinae). *Journal of Zoology London* **201** :445 - 480.
- Bignell, D.E.(1984). Direct potentiometric determination of redox potentials of the gut contents in the termites *Zootermopsis nevadensis* and *Cubitermes severus* and in three other arthropods. *Journal of Insect Physiology* **30** :169 - 174.
- Blomquist, G.J., Howard, R.W. and Mc Daniel, C.A. (1979). Biosynthesis of cuticular hydrocarbons of the termite *Zootermopsis angusticollis* (Hagen).Incorporation of propionate into dimethylalkanes. *Insect Biochemistry* **9** : 371 - 374.
- Blomquist, G.J., Dwyer, L.A., Chu, A.J., Ryan, R.O. and de Renobales, M. (1982). Biosynthesis of linoleic acid in a termite, cockroach and cricket. *Insect Biochemistry* **12** : 349 - 353.
- Booth, S.R., Tanigoshi, L and Dewes, I (2000). Potential of a dried mycelium formulation of an indigenous strain of *Metarhizium anisopliae* against subterranean pests of cranberry. *Biocontrol Science and Technology* **10(5)**: 659 - 668.

- Breznak, J.A. and Pankratz, H.S. (1977). In situ morphology of the gut microbiota of wood-eating termites [*Reticulitermes flavipes* (Kollar) and *Coptotermes formosum* Shiraki]. ***Applied Environmental Microbiology* 33** : 406 - 426.
- Breznak, J.A. (1982). Intestinal microbiota of termites and other xylophagous insects. ***Annual Review of Microbiology* 36** : 323 - 343.
- Breznak, J.A. (1984). Hindgut spirochetes of termites and *Cryptocercus punctulatus*, In: ***Production Ecology of Ants and Termites***, (Ed. M.V.Brian) Cambridge Univ. Press, New York, pp. 67 - 70.
- Breznak, J.A. (1990). Metabolic activities of the microbial flora of termites. In: ***Microbiology in Poeciloterms***, (Ed. R. Lesel), Elsevier ,Amsterdam, pp. 63 - 68.
- Breznak, J.A. and Brune, A. (1994). Role of Microorganisms in the digestion of lignocellulose by termites. ***Annual Review of Entomology* 39** : 453 - 487.
- Burdeos, A.T. and Villacarlos, L.T. (1989). Comparative pathogenicity of *Metarhizium anisopliae*, *Beauveria bassiana* and *Paecilomyces lilacinus* to adult sweet potato weevil, *Cylas formicarius* (F.) (Coleoptera: Curculionidae). ***Phillippines Entomology* 7(6)** : 561-571
- Butt, T.M. and Brownbridge, M. (1997). Fungal pathogens of thrips. In: ***Thrips as crop pests***, (Ed.T. Lewis}, CAB International, Wallingford. pp. 399-433.
- Canale- Parola, E. (1984). Order I. Spirochaetales Buchanan.163AL (Eds. N.R.,Krieg,J.G. Holt) ***Bergey's Manual of Systematic Bacteriology***. Baltimore : Williams & Wilkins **1** : 38 - 39.
- Carruthers, R.I. and R.S. Soper (1987). Fungal disease. In : ***Epizootiology of Fungal Diseases***, (Ed. J.R. Fuxa and Y. Tanada) John Wiley, New York. pp. 357-416.

- Chaîneau, C.H. Morel, J., Dupont, J., Bury, E. and Oudot, J. (1999). Comparison of the fuel oil biodegradation potential of hydrocarbon-assimilating microorganisms isolated from a temperate agricultural soil. *Science of the Total Environment* **227** (2-3) : 237 - 247.
- Chakraborty, B.N. and Saha, A. (1994). Detection and cellular location of cross reactive antigens shared by *Camellia sinensis* and *Bipolaris carbonum*. *Physiology and Molecular Plant Pathology* **44**: 403-416.
- Chakraborty, B.N., Basu, P., Das, R., Saha, A. and Chakraborty, U. (1995). Detection of cross reactive antigens between *Pestalotiopsis theae* and tea leaves and their cellular location. *Annals of Applied Biology* **127** : 11 - 21.
- Chakraborty, B.N. and Chakraborty, U. (2002). Immunodetection of plant pathogenic fungi. In : *Frontiers of Fungal Diversity in India* (Eds. G.P.Rao, C. Manoharachary, D.J.Bhat , R.C. Rajak and T.N.Lakhanpal) International Book Distributing Co., Lucknow, pp. 23-42
- Cherry, A.J., Jenkins, N.E., Heviefo, G., Bateman, R. and Lomer, C.J. (1999). Operational and economic analysis of a West African pilot-scale production plant for aerial conidia of *Metarhizium* spp. for use as a mycoinsecticide against locusts and grasshoppers. *Biocontrol Science and Technology* **9**(1) : 35 - 51.
- Clausen, J. (1988). Laboratory technique in Biochemical and Molecular Biology. Vol.I. Part -III, (Eds. By R.H. Burdon and P.H. Van knippenberg), pp. 64-65.
- Cleveland, L.R. and Grimstone, A.V., (1964). The fine structure of the flagellate *Mixocricha paradoxa* and its associated micro-organisms. *Proceedings of Royal Society of London Ser. B* **159** : 668 - 686.
- Collins, N.M. and Wood, T.G. (1984). Termites and atmospheric gas production. *Science* **224** : 84 - 86.

- Cookson, L.J. (1992). Studies of lignin degradation in mound material of the termite *Nasutitermes exitiosus*. *Australian Journal of Scientific Research* 30 : 189 - 193..
- Cornelius, M.L and Grace, J.K. (1994). Semiochemicals extracted from a Dolichoderine ant affects the feeding and tunneling behaviour of the Formosan subterranean termite (Isoptera : Rhinotermitidae). *Journal of Economic Entomology* 87 (3) : 705-708.
- Czolij, R., Slaytor, M. and O'Brien, R.W. (1985). Bacterial flora of the mixed segment and the hindgut of the higher termite *Nasutitermes exitiosus* Hill. (Termitidae, Nasutitermitinae). *Applied Environmental Microbiology* 49 : 1226 - 1236.
- De. Croos, J.N.A. and Bidochka, M.J. (1999). Effects of low temperature on growth parameters in the entomopathogenic fungus *Metarhizium anisopliae*. *Canadian Journal of Microbiology* 45(12) : 1055 - 1061.
- Driver, F., Milner, R.J. and Trueman, J.W.H. (2000). A taxonomic revision of *Metarhizium* and sequence analysis of ribosomal DNA. *Mycological Research* 104:134-150.
- Edgington, S., Segura, H., William, D.L.R. and Williams, T. (2000). Photoprotection of *Beauveria bassiana* : Testing simple formulation for control of the coffee berry borer. *International Journal of Pest Management* 46(3) : 169 - 176.
- Ekese, S., Maniania N.K. and Onu, I. (1998). Antibiosis and antixenosis of two cowpea varieties to the legume flower thrips. *Africa Crop Science Journal* 6:49-59.
- Ekese, S., Maniania, N.K., and Lwanda, W. (2000). Susceptibility of the legume flower thrips to *Metarhizium anisopliae* on different varieties of cowpea. *Biocontrol Dordrecht* 45(1) : 79 - 95.

- Esenther, G.R. and Beal, R.H. (1978).** Insecticidal baits on field plot perimeters suppress *Reticulitermes*. ***Journal of Economic Entomology* 71** : 604-607.
- Eutick, M.L., O'Brien, R.W. and Slaytor, M. (1978).** Bacteria from the gut of Australian termites. ***Applied Environmental Microbiology* 35** : 823 - 828.
- Eutick, M.L., Veivers, P., O'Brien, R.W. and slaytor, M. (1978).** Dependence of the higher termite, *Nasutitermes exitiosus* and the lower termite, *Coptotermes lacteus* on their gut flora. ***Journal of Insect Physiology* 24** : 363 - 368.
- Fargues, J. (1972).** Etude des conditions D' infection des larves des doryphose, *Leptinotarsa decemlineata* say, par *Beauveria bassiana* (Bals.) Vuill. (Fungi Imperfecti). ***Entomophaga* 17**: 319-377.
- Ferron, P. (1978).** Biological control of insect pests by entomogenous fungi. ***Annual Review of Entomology* 23**:409-442.
- Gillespie, J.P., Burnett, C. and Charnley, A.K. (2000).** The immune response of the desert locust *Schistocerca gregaria* during mycosis of the entomopathogenic fungus, *Metarhizium anisopliae* varacridum. ***Journal of Insect Physiology* 46(4)** : 429 - 437.
- Grace, J.K. (1994).** Protocol for Testing Effects of Microbial Pest control Agents on Nontarget Subterranean Termites (Isoptera : Rhinotermitidae). ***Journal of Economic Entomology* 87(2)** : 269 - 274.
- Guo, L., Quilici, D.R., Chase, J. and Blomquist, G.L. (1991).** Gut tract microorganisms supply the precursors for methylbranched hydrocarbon biosynthesis in the termite, *Zootermopsis nevadensis*. ***Insect Biochemistry* 21** : 327 - 333.

- Gurusubramanian, G., Tamuli, A.K. and Ghosh, R.K. (1999). Susceptibility of *Odontotermes obesus*(Rambur) to *Beauveria bassiana* (Bals.) Vuill. *Insect Science and Application* 19 (2/3) : 157 - 162.
- Hajek, A.E., and Leger, R.J.S. (1994). Interactions between Fungal Pathogen and Insect Hosts. *Annual Review of Entomology* 39 : 293 - 322.
- Hallsworth, J.E. and Magan, N. (1999). Water and temperature relations of growth of the entomogenous fungi *Beauveria bassiana*, *Metarhizium anisopliae*, and *Paecilomyces farinosus*. *Journal of Invertebrate Pathology* 74(3) : 261 - 266.
- Harris, R.J., Harcourt, S.J., Glare, T.R., Rose, E.A.F., Nelson, T.J. (2000). Susceptibility of *Vespula vulgaris* (Hymenoptera : Vespidae) to generalist entomopathogenic fungi and their potential for wasp control. *Journal of Invertebrate Pathology* 75(4) : 251 - 258.
- Hazarika, L.K., Puzari, K.C. and Wahab, S. (2002). Entomopathogenic fungi as a myco-insecticides for management of plantation crops. Proc. Asian Congress of Mycology and Plant Pathology, Mysore, pp. 152 -153.
- Hethener, P., Brauman, A. and Garcia, J.L. (1992). *Clostridium termitidis* sp. nov., a cellulolytic bacterium from the gut of the wood-feeding termite, *Nasutitermes lujae*. *Systematic Applied Microbiology* 15 : 52 - 58.
- Hobbie, J.E. and Melillo, J.M. (1984). Role of microbes in global carbon cycling In: *Current Perspectives in Microbial Ecology*, (Eds. MJ klug. CA Reddy) American Society of Microbiology, Washington DC. pp. 389-393.
- Hogan, M.E., Slaytor, M. and O'Brien, R.W. (1985). Transport of vol. de fatty acids across the hindgut of the cockroach *Panesthia cribrata* Sarssure and the termite, *Mastotermes darwiensis* Froggatt. *Journal of Insect Physiology* 31 : 587 - 591.

- Hogan, M.E., Schulz, M.W., Slayton, M., Czolij, R.T. and O'Brien, R.W. (1988). Components of termite and protozoal cellulases from the lower termite, *Coptotermes lacteus* Froggatt. *Insect Biochemistry* **18** : 45 - 51.
- Hollande, A.C., Gharagozlou, I.D. (1967). Morphologie infrastructurale de *Pillotina calotermitidis* nov. gen. nov. sp. spirochaetale de l'intestin de *Calotermes praecox*. *Sci. Ser. D.* **265** : 1309 - 1312.
- Holland, H.L., Morris, T.A., Nova, P.J. and Zobic, M. (1999). A new paradigm of biohydroxylation of *Beauveria bassiana* ATCC 7159. *Tetrahedron* **55(24)** : 7441 - 7460.
- Hong, T.D., Jenkins, N.E. and Ellis, R.H. (2000). The effects of duration of development and drying regime on the longevity of conidia of *Metarhizium flavoviride*. *Mycological Research* **104(6)** : 662 - 665.
- Honigberg, B.M. (1970). Protozoa associated with termites and their role in digestion In: *Biology of Termites* (Eds.. K. Krishna and F.M. Weesner) Academic Press, New York , pp. 231 - 236.
- Inyang, E.N., Mc Cartney , H.A., Oyejola, B., Ibrahim, L., Pye, B.J., Archer, S.A. and Butt, T.M. (2000). Effect of formulation, application and rain on the persistence of the entomogenous fungus *Metarhizium anisopliae* on oilseed rape. *Mycological Research* **104(6)** : 653 - 661.
- Jacquelyn C., Pendland and Boucias, D.G. (1999). Comparative analysis of the binding of antibodies prepared against the insect *Spodoptera exigue land* against the Mycopathogen *Nomuraea rileyi*. *Journal of Invertebrate Pathology* **75** : 107-116.
- Jaros, S.J., Groden, E. and Zhang, J. (1999). Effects of selected fungicides and the timing of fungicide application on *Beauveria bassiana*-induced mortality of the Colorado potato beetle (Coleoptera : Chrysomelidae). *Biological Control* **15(3)** : 259 - 269.

- Jeffs, L.B., Xavier, I.J., Matai, R.E., and Khachatourians, G.G. (1999). Relationships between fungal spore morphologies and surface properties for entomopathogenic members of the genera *Beauveria*, *Metarhizium*, *Paecilomyces*, *Tolypocladium* and *Verticillium*. **Canadian Journal of Microbiology** 45(ii) : 936 - 948.
- Kang, S.C., Park, S. and Lee, D.G. (1999). Purification and characterization of a novel chitinase from the entomopathogenic fungus, *Metarhizium anisopliae*. **Journal of Invertebrate Pathology** 73(3) : 276 - 281.
- Keerthi, T.R., Suresh, P.V., Sabu, A., Rajeev Kumar, S. and Chandrasekaran, M. (1999). Extracellular production of L-glutaminase by alkalophilic *Beauveria bassiana* BTMF S10 isolated from marine sediment. **World Journal of Microbiology and Biotechnology** 15(6) : 751 - 752.
- Kenne, M., Schatz, B., Durand, J.L. and Dejean, A. (2000). Hunting strategy of a generalist ant species proposed as a biological control agent against termites. **Entomologia Experimentalis et Applicata** 94 (1) : 31-40.
- Khalil, M.A.K. and Rasmussen, R.A. (1990). The influence of termites on atmospheric trace gases : CH₄, CO₂, CHCl₃, N₂O, CO, H₂, and light hydrocarbons. **Journal of Geophysical Research** 95 : 3619 - 3634.
- Kmitowa, K., Augustyniuk, A. and Bajan, C. (2000). A response of entomopathogenic fungi to plant extracts. **Polish Ecological Studies** 23 (3-4) : 171 - 185.
- Kovoor, J. (1968). L'intestin d'un termite superieur (*Microcerotermes edentatus*, Wasman, Amitermitinae). Histophysiologic et flore bacterienne symbiotique. **Biological Bulletin of France** 102 : 45 - 84.
- Krishna, K. (1969). Introduction, In: **Biology of Termites** (Ed., K. Krishna and F.M. Weesner) Academic Press, New York, pp. 1 - 17.

- Krishna, K. (1970). Taxonomy, phylogeny and distribution of termites. In : *Biology of Termites* (Ed.K. Krishna and F.M. Weesner) Academic Press, New York, pp. 127-152.
- Kulye, M., Nahar, P., Chavan, S., Kulkarni, M. and Deshpande, M.V. (2002). Fungi and fungal products- as a Potential Biocontrol agents. Prroc. Asian Congress of Mycology and Plant Pathology, Mysore, pp. 153.
- Laemmli, U.K. (1970). Cleavage of structural proteins during assembly of the head of the bacteriophage T₄. *Nature* **227** : 680 - 685.
- Lai, Y.Z. and Sarkanen, K.V.(1971). Isolation and structural studies. In : *Lignans : Occurrence, Formulation, Structure and Function* (Eds. K.V.Sarkanen, and C.H. Ludwin) Wiley-Interscience , New York, pp.165 - 240.
- Lange, L.; Heide, M. and Olson, L.W. (1989). Serological detection of *Plasmodiophora brassicae* by dot immunobinding and visualization of the serological reaction by scanning electron microscopy. *Phytopathology* **79** : 1066 - 1071.
- Lee, K.E. and Wood, T.G. (1971). Termites and Soils. Academic press, New York, pp. 251.
- Ljungdahl, L. G. and Erikson, K.E. (1985). Ecology of microbial cellulose degradation. *Advance Microbial Ecology* **8**:237-299.
- Lopez Lorica, L.V., Carbone II, T. and Salinas, J. (1999). Colonization of plant waste substrates by entomopathogenic and mycoparasitic fungi - A SEM Study. *Micron* **30**(4) : 325 -333.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with folin phenol reagent. *Journal of Biology and Chemistry* **193**: 265 - 275.

- Luz, C. and Fargues, J. (1999). Dependence of the entomopathogenic fungus, *Beauveria bassiana*, on high humidity for infection of *Rhodnius prolixus*. *Mycopathologia* 146(1) : 33 - 41.
- Magalhaes, B.P., Lecoq, M., De, F.M.R., Schmidt, F.G.V. and Guerra, W.D. (2000). Field trial with the entomopathogenic fungus *Metarhizium anisopliae* var. *acridum* against bands of the grasshopper *Rhammatocerus schistocercoides* in Brazil. *Biocontrol Science and Technology* 10(4) : 427 - 441.
- Margulis, L. and Hinkle, G. (1992). Large symbiotic spirochetes : *Clevelandina*, *Cristispira*, *Diplocalyx*, *Hollandina*, and *Pillotina*. In : *The Prokaryotes* (Eds. A., Balows, H.G Truper, M. Dworkin, W . Harder and K.H Schleffer) Springer Verlag, New York . 4 : 3965 - 3978.
- Martin, M.M. and Martin, J.S. (1978). Cellulose digestion in the midgut of the fungus -growing termite *Macrotermes natalensis* : the role of acquired digestive enzymes. *Science* 199 : 1453 - 1455.
- Martin, M.M. and Martin, J.S. (1979). The distribution and origins of the cellulolytic enzymes of the higher termite, *Macrotermes natalensis*. *Physiological Zoology* 52 : 11 - 21.
- Martin, M.M. (1983). Cellulose digestion in insects. *Compendium of Biochemical Physiology* 75 A : 313 -324.
- Martin, M.M. (1987). Invertebrate - Microbial Interactions : Ingested Fungal Enzymes in Arthropod Biology. Comstock Pub. Assoc, Ithaca, N.Y. pp.148
- Martin, P.A.W., Schroder, R.F.W., Poprawski, T.J., Lipa, J.J., Hausvater, E. and Rasocha, V. (2000). Temperature effects on the susceptibility of the Colorado potato beetle (Coleoptera : Chrysomelidae) .to *Beauveria bassiana* (Balsamo) Vuillemin in Poland, the Czech Republic and the United States. *Journal of Entomological Science*. 35(3) : 251-258.

- Mauldin, J.K. (1982).** Lipid Synthesis from [^{14}C]- acetate by two subterranean termites, *Reticulitermes flavipes* and *Coptotermes formosanus*. ***Insect Biochemistry* 12** : 193 - 199.
- Mc Ewen, S.E., Slaytor, M. and O'Brien, R.W. (1980).** Cellobiase activity in three species of Australian termites. ***Insect Biochemistry* 10** : 563 - 567.
- Mishra, S.C. (1980).** Carbohydrases in *Neotermes bosei* Snyder (Isoptera : Kalotermitidae). ***Mater.Org.* 15** : 253 – 26.
- Mishra S.C., Sen-Sarma P.K. (1987).** Origin and distribution of digestive enzymes in soldiers of *Neotermes bosei* Snyder with a note on gut morphology (Isoptera : Kalotermitidae). ***Mater.Org.* 22**: 127 – 138.
- Mohan, C.M., Lakshmi, K.A., and Devi, K.U. (1999).** Laboratory evaluation the Pathogenicity of three isolates of the entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuillemin on the American cockroach (*Periplanata americana*). ***Biocontrol Science and Technology* 9(1)** : 29 - 33.
- Mohan, S. B. (1988).** Evaluation of antisera raised against *Phytophthora fragariae* for detecting red core disease of strawberries by enzyme-linked immunosorbent assay (ELISA). ***Plant Pathology* 37**: 206-216.
- Moussou, P., Archelas, A., Furstoss, R. and Baratti, J.C. (2000).** Clues for the existence of two different epoxide hydrolase activities in the fungus *Beauveria bassiana* . ***Enzymes and Microbial Technology* 26(5-6)** : 414 - 420.
- Mulock, B. and Chandler, L. (2000).** Field cage studies of *Beauveria bassiana* (Hyhomyces : Moniliaceae) for the suppression of adult western corn rootworm, *Diabrotica virgifera virgifera*(Coleoptera : Chrysomelidae). ***Biocontrol Science and Technology* 10(1)** : 51 - 60.
- Nankinga, C.M. and Moore, D (2000).** Reduction of banana weevil population using different formulations of the entomopathogenic fungus *Beauveria bassiana*. ***Biocontrol Science and Technology* 10(5)**; 645 - 657.

- Niranjana, S.R. (2002). Biological control of Coffee Berry Borer using an Entomopathogenic fungus *Beauveria bassiana*. Asian Congress of Mycology and Plant Pathology, Mysore, pp. 151.
- Noirot, C. and Noirot, T.C. (1969). The digestive system, In: ***Biology of Termites***. (Eds. K. Krishna and F.M. Weesner), Academic Press, New York, 49- 88 pp.
- Noma, T. and Strickler, K. (1999). Factors affecting *Beauveria bassiana* for control of Lygus bug (Hemiptera : Miridae) in alfalfa seed fields. ***Journal of Agricultural and Urban Entomology* 16(4) : 215 - 233.**
- O'Brien, G.W., Veivers, P.C., Mc Ewen, S.E., Slaytor, M. and O' Brien, R.W. (1979). The origin and distribution of cellulase in the termites, *Nasutitermes exitiosus* and *Coptotermes lacteus*. ***Insect Biochemistry* 9: 619 -625.**
- O' Brien, R.W. and Slaytor, M. (1982). Role of microorganisms in the metabolism of termites. ***Australian Journal of Biological Science* 35 : 239 - 262.**
- O' Brien, R.W. and Breznak, J.A. (1984). Enzymes of acetate and glucose metabolism in termites. ***Insect Biochemistry* 14: 639 - 643.**
- Odelson, D.A. and Breznak, J.A. (1983). Volatile fatty acid production by the hindgut microbiota of xylophagous termites. ***Applied Environmental Microbiology* 45 : 1602 - 1613.**
- Odelson, D.A. and Breznak, J.A. (1985). Nutrition and growth characteristics of *Trichomitopsis termopsidis*, a cellulolytic protozoan from termites. ***Applied Environmental Microbiology* 49 : 614 - 621.**
- Odelson, D.A. and Breznak, J.A. (1985)a. Cellulase and other polymer hydrolyzing activities of *Trichomitopsis termopsidis*, a symbiotic protozoan from termites. ***Applied Environmental Microbiology* 49: 622 - 626.**

- Ouchterlony, O.(1967). *Handbook of experimental immunology*,(ed. Weir) Blackwell Ins. ,London, pp.655.
- Potrikus, C. J. , Breznak, J.A. (1981).Gut bacteria recycle uric acid nitrogen in termites : a strategy for nutrient conservation. *Proceedings of National Academy of Science, USA 78* : 4601 - 4605.
- Potts, R.C., Hewitt, P. H. (1973). The distribution of intestinal bacteria and cellulase activity in the harvester termite *Trinervitermes trinervoides* (Nasutitermitinae). *Insect Sociology 20* : 215 - 220.
- Potts, R.C., and Hewitt, P. H. (1974). The partial purification and some properties of the cellulase from the termite *Trinervitermes trinervoides* (Nasutitermitinae). *Compendium of Biochemical Physiology 47 B* : 317 - 326.
- Potts, R. C. and Hewitt, P. H. (1974)a. Some properties and reaction characteristics of the partially purified cellulase from the termite *Trinervitermes trinervoides* (Nasutitermitinae). *Compendium of Biochemical Physiology 47B* : 327 - 337.
- Prasad, V.D., Jayaraj, S. and Rabindra, R.J. (1989). Susceptibility of tobacco caterpillar, *Spodoptera litura* Fab. (Noctuidae :Lepidoptera) to certain entomogenous fungi. *Journal of Biological Control 3 (1)* : 53-55.
- Presturich, G.D., Jones, R.W. and Collins, M.S. (1981). Terpene biosynthesis by nasute termite soldiers (Isoptera Nasutitermitinae).*Insect Biochemistry 11* : 331 - 336.
- Pusti, M. B., Belli, M. L. (1985). Cellulolytic activity of actinomycetes isolated from termite (Termitidae)gut. FEMS. *Microbiological Letters 26* : 107 - 112.
- Rath, C.A. (2000). The use of entomopathogenic fungi for control of termites. *Biocontrol Science and Technology 10(5)* : 563 – 581.

- Renn, N., Bywater, A.F. and Barson, G. (1999). A bait formulated with *Metarhizium anisopliae* for the control of *Musca domestica* L. (Dipt., Muscidae) assessed in large- scale laboratory enclosures. *Journal of Applied Entomology* 123(5) : 309 - 314.
- Retief, L.W. and Hewitt, P. H. (1973). Digestive β - glycosidases of the harvester termite, *Hodotermes mossambicus* : properties and distribution . *Journal of Insect Physiology* 19 : 1837 - 1847.
- Rica, W.C. and Cogburn, R.R. (1999). Activity of the entomopathogenic fungus *Beauveria bassiana* (Deuteromycota : Hyphomycetes) against three Coleopteran pests of stored grain. *Journal of Economic Entomology* 92(3) : 691 – 694.
- Rombach, M. C. , Rombach, G. M. , Aguda, R. M. and Roberts, D.W. (1987)
Counting of conidia of *Metarhizium anisopliae*, *Metarhizium flavoviride*, and *Beauveria bassiana* by light Transmission measurements on conidial suspensions. *Philippines Entomology* 7(1): 43 - 50.
- Rosa, D.L., Segura, H.R., Barrera, J.F. and Williams, T. (2000). Laboratory evaluation of the impact of entomopathogenic fungi on *Prorops nasuta* (Hymenoptera : Bethyridae), a parasitoid of Coffee berry borer. *Environmental Entomology* 29(1) : 126-131.
- Rouland, C., Mora, P. Matoub, M. and Renoux. J. (1986). Etude comparative entre la β -glucosidase presente dans le digestif du termite *Macrotermes Mulleri* et la β -glucosidase du champignon symbiotique *Termitomyces* sp. *Acta Insectes Soc.* 3 : 109 - 118.
- Rouland, C., Civas, A., Renoux, J. and Petek, F. (1988). Purification and properties of cellulases from the termite *Macrotermes mulleri* (Termitidae, Macrotermitinae)and its symbiotic fungus *Termitomyces* sp. *Compendium of Biochemical Physiology* 91 B : 449 - 458.

- Rouland, C., Civas, A., Renoux, J. and Petek, F. (1988)a. Synergistic activities of the enzymes involved in cellulose degradation, purified from *Macrotermes mulleri* and from its symbiotic fungus *Termitomyces* sp. ***Compendium of Biochemical Physiology* 91B** : 459 - 465.
- Rouland, C., Mora, P. and Renoux, J. (1988)b. Essai D'interpretation de la symbiose digestive chez *Macrotermes mulleri* (Termitidae : Macrotermitinae). ***Actes Coil Insectes Soc.* 5**: 69 - 77.
- Rouland, C., Renoux, J. and Petek, F. (1988)d. Purification and properties of two xylanases from *Macrotermes mulleri* (Termitidae. Macrotermitinae) and its symbiotic fungus *Termitomyces* sp. ***Insect Biochemistry* 18** : 709 - 715.
- Rouland, C., Chararas, C. and Renoux, J. (1989). Les osidases digestives presentes dans l'intestin moyen. l' intestin posterieur it les glandes salivaires du termite humivore *Crenetermes albotarsalis*. ***Academic Science Paris* 308** : 281 - 285.
- Samsinakova, A. and Kalalova, S. (1983). The Influence of single - spore isolate and Repeated Subculturing on the pathogenecity of conidia of the Entomophagous Fungus *Beauveria bassiana*. ***Journal of Invertebrate Pathology* 42** : 156 -161.
- Schutz, J.E. and Breznak, J.A. (1978). Heterotrophic bacteria present in guts of wood-eating termites [*Reticulitermes flavipes* (kollar)]. ***Applied and Environmental Microbiology* 35** : 930 - 936.
- Schutz, M.W., Slaytor, M., Hogan, M. and O'Brien, R.W. (1986). Components of cellulase from the higher termite, *Nasutitermes walkeri*. ***Insect Biochemistry* 16** : 929 - 932.
- Sen-Sarma, P. K. (1995). Recent trends and strategies in subterranean termite control in buildings. In: ***Termite Management in Buildings*** (Ed. Y.Singh) Tata Mc Graw Hill Publishing Company Limited, New Delhi, pp. –21.

- Sharma, S., Gupta, R.B.L. and Yadava, C.P.S. (1999).** Mass multiplication and formulation of entomopathogenic fungi and their efficacy against whitegrubs. *Journal of Mycology and Plant Pathology* **29 (3)** : 299 - 305.
- Slaytor, M. (1993).** Cellulose digestion in termites and cockroaches : what role do symbionts play? *Compendium of Biochemical Physiology* **103B** : 775 – 784.
- Smits, J.E., Johnson, D. L. and Lomer, C. (1999).** Pathological and Physiological responses of ring-necked pheasant chicks following dietary exposure to the fungus *Metarhizium flavoviride*, a biocontrol agent for locusts in Africa. *Journal of Wildlife Diseases* **35 (2)** : 194-203
- St. Leger, R.J., May, B., Allee, L.L., Frank, D.C., and Roberts, D.W. (1992).** Genetic differences in allozymes and in formation of infection structures among isolates of the entomopathogenic fungus, *Metarhizium anisopliae*. *Journal of Invertebrate Pathology* **60**: 89 - 101.
- St. Leger, R.J., Nelson, J.O. and Screen, S.E. (1999).** The entomopathogenic fungus *Metarhizium anisopliae* alters ambient pH, allowing extracellular protease production and activity. *Microbiology Reading* **145(10)** : 2691 - 2699.
- Su, N.Y., Ban, P. M. and Scheffrahn, R.H. (1991).** Suppression of Foraging Population of the Formosan Subterranean Termite (Isoptera: Rhinotermitidae) by Field Application of a Slow- Acting Toxicant Bait. *Journal of Economic Entomology* **84(5)** : 1525 - 1531.
- Su, N.Y. (1994).** Field evaluation of Hexaflumuron Bait for population suppression of subterranean Termites (Isoptera : Rhinotermitidae). *Journal of Economic Entomology* **87(2)**: 389 - 397.

- To, L.P., Margulis, L., Chase, D. and Nutting, W.L. (1980). The symbiotic microbial community of the Sonoran Desert termite: *Pterotermes occidentis*. ***Bio Systems* 13** : 109 - 137.
- Vanninen, I., Tyni, J.J. and Hakkanen, H. (2000). Persistence of augmented *Metarhizium anisopliae* and *Beauveria bassiana* in Finnish agricultural soils. ***Biocontrol Dordrecht* 45(2)** : 201 - 222.
- Veivers, P.C., O'Brien, R.W. and Slaytor, M. (1980). The redox state of the gut of termites. ***Journal of Insect Physiology* 26** : 75 - 77.
- Veivers, P.C., O'Brien, R.W. and Slaytor, M. (1982). Role of bacteria in maintaining the redox potential in the hind gut of termites and preventing entry of foreign bacteria. ***Journal of Insect Physiology* 28** : 947 - 951.
- Veivers, P.C., O'Brien, R.W., Slaytor, M. (1983). Selective defaunation of *Mastotermes darwiniensis* and its effect on cellulose and starch metabolism. ***Insect Biochemistry* 13** : 95 - 101.
- Veivers, P.C., Mutilemann, R., Slaytor, M., Leuthold, R.H. and Bignell, D.E. (1991). Digestion, diet and polyethism in two fungus - growing termites : *Macrotermes subhyalinus* Rambur and *M. michaelsoni* Sjostedt. ***Journal of Insect Physiology* 37** : 675 - 682.
- Verkerk, R.H.J.; Bravery, A.F.; and Wright, D.J. (1999). Considering the feasibility of using novel techniques for termite control in tropical and subtropical agriculture. Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit ***Gent* 64(3a)**: 177-184.
- Wagner, B.L. and Lewis, L.C. (2000). Colonization of corn, *Zea mays* by the entomopathogenic fungus *Beauveria bassiana*. ***Applied and Environmental Microbiology* 66(8)** : 3468 - 3473.

- White, J.G., Lyons, N.F. Wakeham A.J. Mead, A and Green J.R. (1994). Serological profiling of the fungal genus *Pythium*, *Physiology and Molecular Plant Pathology* **44**:349-361.
- Wood, T.G. and Sands, W.A. (1978). The role of termites in ecosystems. In : *Production Ecology of Ants and Termites* (ed. M.V. Brian) Cambridge Univ. Press, New York. pp. 245 - 249..
- Wood, T.G. and Johnson, R.A. (1986). The biology , physiology and ecology of termites. In: *Economic Impact and Control of Social Insects*, (Ed. SB Vinson), Praeger ,New York ,pp. 1 - 68.
- Wood, T.G. and Thoma, R.J. (1989). The mutualistic association between Macrotermitinae and Termitomyces. In :*Insect- Fungus Interactions*. (ed. N. Wilding, N.M. Collins, P.M. Hammond, J.F. Weber), Academic press, New York, pp. 69 -92
- Wraight, S.P. , Carruthers, R.I., Jaronski, S.T., Bradley, C.A., Garza, C.J. and Galaini, W.S.(2000). Evaluation of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* for microbial control of the silver leaf whitefly, *Bemisia argentifolii*. *Biological Control* **17 (3)** : 203 - 217.
- Xavier, S.S., Magalhaes, B. and Luna, A.L.E.A. (1999). Differentiation of the entomopathogenic fungus *Metarhizium flavoviride* (Hyphomycetes). *Revista de Microbiologia* **30(1)** : 47 - 51.
- Yamaoka, I. and Nagatani, Y. (1975). Cellulose digestion system in the termite, *Reticulitermessperatus* (kolbe). I. producing sites and physiological significance of two kinds of cellulase in the worker. *Zoological Magazine* **84** : 23 - 29.
- Yamin, M.A. (1978). Axenic cultivation of the cellulolytic flagellate *Trichomitopsis termopsidis*(Cleveland) from the termite *Zootermopsis*. *Journal of Protozoology* **25** : 535 - 538.

- Yamin, M.A. and Trager, W. (1979). Cellulolytic activity of an axenically-cultivated termite flagellate, *Trichomitopsis termopsidis*. **Journal of General Microbiology** 113 : 417 - 420.
- Yamin, M.A. (1980). Cellulose metabolism by the termite flagellate *Trichomitopsis termops*. **Applied and Environmental Microbiology** 39 : 859 - 863.
- Yamin, M.A. (1981). Cellulose metabolism by the flagellate *Trichonympha* from a termite is independent of endosymbiotic bacteria. **Science** 211 : 58 - 59.
- Yokoe, Y. (1964). Cellulase activity in the termite, *Leucotermes speratus*, with new evidence in support of a cellulase produced by the termite itself. **Sci. Papers Coll. Gen. Educ. University of Tokyo** 14 : 115 - 120.
- Yoshimura, T. Azuma, J., Tsunoda, K. and Takahashi, M. (1993). Cellulose metabolism of the symbiotic protozoa in termite, *Coptotermes formosanus* Shiraki (Isoptera : Rhinotermitidae)l. Effect of degree of polymerization of cellulose. **Mokuzai Gakkaishi** 39 : 221 - 226.
- Zurek, L. and Keddie, B.A. (2000). *Beauveria bassiana* (Balsamo) Vuillemin: a promising microbial control agent of the satin moth (Lepidoptera : Lymantriidae). **Biocontrol Science and Technology** 10(5) : 641 - 644.

