

7. REFERENCES

- Abraham, K.J. and Daniel, M. (1988) Phytoalexins of Cassia fistula L. and Morinda tomentosa Heyne. Nat. Acad. Sci. Lett. 11 : 101-102.
- Agrios, G.N. (1978) Plant Pathology, p. 703. Academic Press, New York.
- Ahl, P. and Gianinazzi, S. (1982) b-protein as a constitutive components in highly (TMV) resistant interspecific hybrids of N. glutinosa x N. debbeyi. Plant Sci. Lett. 26 : 173.
- Akazawa, T. and Wada, K. (1961) Analytical study of ipomeamarone and chlorogenic acid alterations in sweet potato roots infected by Ceratocystis fimbriata. Plant Physiology 36 : 139-141.
- Alam, M., Sattar, A. and Janardhanan, K.K. (1991) Changes in phenol and peroxidase in the leaves of Java citronella infected with Curvularia andropogonis. Biol. Plant 33 : 211-215.
- Alba, A.P.C. and Devay, J.E. (1985) Detection of cross reactive antigens between Phytophthora infestans (Mont.) de Bary and Solanum species by indirect enzyme-linked immunosorbent assay. Phytopath. Z. 112 : 97-104.
- Albersheim, P. and Anderson-Prouty, A.J. (1975) Carbohydrates, proteins, cell surfaces and biochemistry of pathogenesis. Ann. Rev. Plant Physiol. 26 : 31-52.
- Albersheim, P., Valent, B.S., Hahn, M., Wade, M. and Cline, K. (1978) Elicitation and sites of formation of phytoalexins and induced resistance. 3rd Int. Congr. Plant Pathol., Munchen, Germany (Abstr.).
- Anderson, A.J. (1980) Studies on the structure and elicitor activity of extra-cellular components produced by three races of fungal plant pathogen Colletotrichum lindemuthianum. Can. J. Microbiol. 26 : 1473-1479.

- Anderson-Prouty, A.J. and Albersheim, P. (1975) Host pathogen interactions. VIII. Isolation of a pathogen synthesized fraction rich in glucan that elicits a defense response in the pathogen's host. Plant Physiol. 56 : 286-291.
- Andrews, J.H. (1990) Biological control in the phyllosphere : Realistic goal or false hope ? Can. J. Plant Pathol. 12 : 300-307.
- Andrews, J.H. (1992) Biological control in the phyllosphere. Ann. Rev. Phytopath. 30 : 603-635.
- Andrews, J.H., Kenerley, C.M. and Nordheim, E.V. (1980) Positional variation in phylloplane microbial populations within an apple tree canopy. Microbial Ecology 6 : 71-84.
- Ando, Y. and Hamaya, E. (1986) Defense reaction of tea plant against infection of the tea anthracnose fungus. Study of Tea 69 : 35-43.
- Anonymous (1980) The United States Pharmacopeia. XX; United States Pharmacopoeial Convention Inc. 12601 Twin brook, Parkway, Rockville, MD 20852, p. 980.
- Arnows, E.L. (1933) Colorimetric determination of the component of 2,4-dihydroxyphenyl-alanine-tyrosine mixture. J. Biol. Chem 118 : 513-537.
- Austin, B., Dickinson, C.H. and Goodfellow, M. (1977) Antagonistic interactions of phylloplane bacteria with Drechslera dictyoides (Drechsler) Shoemaker. Can. J. Microbiol. 23 : 710-715.
- Azevedo, E.C., Rios, E.M., Fukushima, K. and Campos-Takaki, G.M. (1993) Bacitracin production by a new strain a Bacillus subtilis : Extraction, purification and characterization. Appl. Biochem. Biotechnol. 42 : 2-7.

- Bailey, J.A. and Deverall, B.J. (1983) The Dynamics of Host Defence. Academic Press, Sydney, p. 233.
- Ballou, C. (1976) Structure and biosynthesis of the mannan component of the yeast cell envelope. Adv. Microb. Physiol. 14 : 93-158.
- Baker, C.J., O'Neill, N.R. and Tomerlin, J.R. (1989) Accumulation of phenolic compounds incompatible clone/race interactions of Medicago sativa and Colletotrichum trifolii. Physiol. Mol. Plant Pathol. 35 : 231-241.
- Baker, C.J., Stavley, J.R. and Mock, N. (1985) Biocontrol of bean rust by Bacillus subtilis under field conditions. Plant Dis. 69 : 770-772.
- Barbora, B.C. (1988) Latest concept on management of young tea. In : Field management in tea (Ed. S.K. Sarkar, and H.P. Bezbaruah), pp. 56-62. Tocklai Experimental Station, Jorhat, India.
- Barnett, H.L. (1963) The nature of mycoparasitism by fungi. Ann. Rev. Microbiol. 17 : 1-4.
- Barry, A.L. and Feeney, K.L. (1967) Two quick methods for Voges-Proskauer test. Appl. Microbiol. 15 : 1138.
- Bartholomew, J.W. (1962) Variables influencing results, and the precise definition of steps in Gram staining as a means of standardizing the results obtained. Stain Technol. 37 : 139-155.
- Bayer, O. (1989) Phenolic metabolism in bark of sweet cherry after infection by 2 bark pathogen. Eur. J. Biochem. 125 : 213-219.
- Bazzalo, M.E., Heber, E.M., Martinez, M.A.P.P. and Caso, O.H. (1985) Phenolic compounds in stems of sunflower plants inoculated with

Sclerotinia sclerotiorum and their effects on fungus. Phytopath.
Z. 112 : 322-332.

Beissman, B., Engels, W., Kogel, K., Marticke, K.H. and Reisener, H.J.
(1992) Elicitor active glycoproteins in apoplastic fluids of
stem rust infected wheat leaves. Physiol. Mol. Plant Pathol. 40
: 79-89.

Bergey's Manual of Systemic Bacteriology (Eds. P.H.A. Sneath, N.S.
Mair, M.E. Sharpe and J.G. Holt) Vol. 2. Williams & Wilkins,
Baltimore.

Bezbaruah, H.P. and Singh, I.D. (1988) Advances in the use of planting
material. In : Field management in tea. (Ed. S.K. Sarkar and
H.P. Bezbaruah). Tocklai Experimental Station, Jorhat, India.

Bhattacharyya, B. and Purkayastha, R.P. (1983) Microbial activities and
leaf surface environment. Indian Rev. Life Sc. 3 : 185-206.

Bhattacharyya, B. and Purkayastha, R.P. (1988) Alteration of chemical
environment of jute phylloplane and its impact on foliar
microbes and anthracnose disease. Int. J. Tropical Plant
Diseases 6 : 77-84.

Blakeman, J.P. (1973) The chemical environment of leaf surfaces with
special reference to spore germination of pathogenic fungi.
Pestic. Sci. 4 : 575-588.

Blakeman, J.P. (1981) Microbial Ecology of the phylloplane. Academic,
London.

Blakeman, J.P. and Atkinson, P. (1981) Antimicrobial substances
associated with the aerial surfaces of plants. In: Microbial
Ecology of the Phylloplane, pp. 254-263. Academic Press, London.

- Blakeman, J.P. and Brodie, I.D.S. (1976) Inhibition of pathogens by epiphytic bacteria on aerial plant surfaces. In : Microbiology of Aerial Plant Surfaces (Eds. C.H. Dickinson and T.F. Preece), pp. 529-557. Academic Press, London.
- Blakeman, J.P. and Fokkema, N.J. (1982) Potential for biological control of plant disease of the phylloplane. Ann. Rev. Phytopathol. 20 : 167-192.
- Boland, G.J. and Inglis, G.D. (1989) Antagonism of white mold (Sclerotinia sclerotiorum) of bean by fungi from bean and rapeseed flowers. Can. J. Bot. 67 : 1775-1781.
- Bolwell, G.P., Robbins, M.P. and Dixon, R.A. (1985) Metabolic changes in elicitor-treated bean cells. Enzymic responses associated with rapid changes in cell wall components. Eur. J. Biochem. 148 : 571-578.
- Borkar, S.G. and Verma, J.P. (1991) Dynamics of phenolics and diphenoloxidase contents of cotton cvs. during hypersensitivity and susceptible reaction induced by Xanthomonas campestris pv. malvacearum. Indian Phytopathol. 44 : 281-290.
- Bornside, G.H. and Kallio, R.E. (1956) Urea hydrolyzing bacilli - a physiological approach to identification. J. Bact. 71 : 627.
- Bostock, R.M. and Stermer, B.A. (1989) Perspectives on wound healing in resistance to pathogens. Ann. Rev. Phytopathol 27 : 343-371.
- Bourdreau, M.A. and Andrews, J.H. (1987) Factors influencing antagonism of Chaetomium globosum to Venturia inaequalis : A case study in failed biocontrol. Phytopathology 77 : 1470-1475.

- Brodie, I.D.S. and Blakeman, J.P. (1975) Competition for carbon compounds by a leaf surface bacterium and conidia of Botrytis cinerea. Physiol. Plant Pathol. 6 : 125-135.
- Brown, A.E., Finlay, R. and Ward, J.S. (1987) Antifungal compounds produced by Epicoccum purpurascens against soil borne plant pathogenic fungi. Soil Biol. Biochem. 19 : 657-664.
- Brown, R.G. and Kimmins, W.C. (1977) Glycoproteins. International Review of Biochemistry 13 : 183-207.
- Bruegger, B.B. and Keen, N.T. (1979) Specific elicitors of glyceollin accumulation in the Pseudomonas glycinea-soybean host parasite. Physiol. Plant Pathol. 15 : 43-51.
- Bruzzese, E. and Hasan, S. (1983) A whole leaf clearing and staining technique for host specificity studies of rust fungi. Plant Pathol. 32 : 335-338.
- Bryngelsson, T. and Green, B. (1989) Characterization of a pathogenesis-related, thaumin-like protein isolated from barley challenged with an incompatible race of mildew. Physiol.Mol. Plant Pathol. 35 : 45-52.
- Bryngelsson, T., Gustaffsson, M., Leal, M.R. and Bartonek, E. (1988) Induction of pathogenesis-related proteins in barley during the resistance reaction to mildew. J. Phytopathol. 123 : 193-198.
- Callow, J.A. (1977) Recognition, resistance and the role of plant lectins in host-parasite interaction. Adv. Bot. Res. 4 : 1-49.
- Chakraborty, B.N. (1988) Antigenic Disparity. In : Experimental and Conceptual Plant Pathology (Eds. R.S. Singh, U.S.Singh, W.M. Hess and D.J. Webber), pp. 477-484. Oxford & I.B.H. Publishing, New Delhi, India.

- Chakraborty, B.N., Chakraborty, U. and Basu, K. (1994a) Antagonism of Erwinia herbicola towards Leptosphaeria maculans causing blackleg disease of Brassica napus. Letters in Applied Microbiology 18 : 74-76.
- Chakraborty, B.N., Chakraborty, U., Das, S.K., Das, G. and Saha, A. (1994b) Changes in phenolics of susceptible and resistant tea leaves after challenge with foliar fungal pathogens. In: 16th International Congress of Biochemistry and Molecular Biology, New Delhi, p. 161.
- Chakraborty, B.N., Das, G., Das, S.K. and Chakraborty, U. (1994c) Phyllosphere microflora of tea (Camellia sinensis (L.) O. Kuntze) and their interaction with Glomerella cingulata, the causal agent of brown blight disease. Tea 15 : 27-34.
- Chakraborty, B.N., Chakraborty, U. and Saha, A. (1995a) Defense strategies of tea (Camellia sinensis) against fungal pathogens. In : Handbook of Phytoalexin Metabolism and Action. (Eds. M. Daniel and R.P. Purkayastha), pp. 485-502. Marcel Dekker Inc., N.Y.
- Chakraborty, U., Das, G., Das, S.K. and Chakraborty, B.N. (1995b) Screening of phyllosphere microorganisms of Camellia sinensis for antagonism to Glomerella cingulata. In : Recent Advances in Phytopathological Researches. (Eds. A.K.Roy and K.K.Sinha) pp. 205-211. M.D. Publications, New Delhi, India.
- Chakraborty, U. and Chakraborty, B.N. (1989) Interaction of Rhizobium leguminosarum and Fusarium solani f. sp. lisi on pea affecting disease development and phytoalexin production. Can. J. Bot. 67 : 1698-1701.

- Chakraborty, B.N. and Purkayastha, R.P. (1983) Serological relationship between Macrophomina phaseolina and soybean cultivars. Physiological Plant Pathology 23 : 197-205.
- Chakraborty, B.N. and Purkayastha, R.P. (1987). Alteration of glyceollin synthesis and antigenic patterns after chemical induction of resistance in soybean to Macrophomina phaseolina. Can. J. Microbiol 33 : 835-840.
- Chakraborty, B.N. and Saha, A. (1989) Biological activity of leaf diffusates of tea in relation to resistance to Bipolaris carbonum. Environ. Ecol. 7 : 131-133.
- Chakraborty, B.N. and Saha, A. (1994a) Accumulation of antifungal compounds in tea leaf tissue infected with Bipolaris carbonum. Folia Microbiol. 39 : 409-414.
- Chakraborty, B.N. and Saha, A. (1994b) Detection and cellular location of cross-reactive antigens shared by Camellia sinensis and Bipolaris carbonum. Physiol. Mol. Plant Pathol. 44 : 403-416.
- Chakraborty, B.N. and Saha, A. (1995) Detection of antifungal compounds in tea leaves after infection with Bipolaris carbonum. In : Recent Advances in Phytopathological Researches (Eds. A.K.Roy and K.K. Sinha), pp. 205-211. M.D. Publications, New Delhi.
- Chambers, C.W. (1950) Relationship of coliform bacteria to gas production in media containing lactose. Public Health Rept. 65 : 619.
- Christ, U. and Mosinger, E. (1989) Pathogenesis-related proteins of tomato : I. Induction by Phytophthora infestans and other biotic

and abiotic inducers and correlations with resistance. Physiol. Mol. Plant Pathol. 35 : 53-65.

- Cimino, G. and Bracci, R. (1960) Analisi quantitative della produzione di indole in E. coli. Riv. Biol. 52 : 395.
- Clark , C.A. and Lorbeer, J.W. (1977) Comparative nutrient dependency of Botrytis squamosa and B. cinerea for germination of conidia and pathogenicity on onion leaves. Phytopathology 67 : 212-218.
- Clarke, A.E. and Knox, R.B. (1978) Cell recognition in flowering plants. Q. Rev. Biol. 53 : 3-28.
- Clarkson, J.P. and Lucas, J.A. (1993a) Screening for potential antagonists of Pseudocercospora herpotrichoides, the causal agent of eyespot disease of cereals : 1. Bacteria. Plant Pathol. 42 : 543-551.
- Clarkson, J.P. and Lucas, J.A. (1993b) Screening for potential antagonists of Pseudocercospora herpotrichoides, the causal agent of eyespot disease of cereals : 2-fungi. Plant Pathol. 42 : 552-559.
- Coleman, M.J., Mainzer, J. and Dickerson, A.G. (1992) Characterization of a fungal glycoprotein that elicits a defense response in French bean. Physiol. Mol. Plant Pathol. 40 : 333-351.
- Cook. R.J. (1993) Making greater use of introduced microorganisms for biological control of plant pathogens. Ann. Rev. Phytopathol. 31 : 58-80.
- Cook, R.J. and Baker, K.R. (1983) The nature and practice of biological control of plant pathogens. Am. Phytopathol. Soc. pp. 539.

- Cook, R.T.A. (1989) Control of Glomerella cingulata f. sp. camelliae with fungicides. Plant Pathol. 38 : 514-519.
- Cruickshank, I.A.M. (1963) Phytoalexins. Ann. Rev. Phytopathol. 1 : 351-374.
- Cruickshank, I.A.M. (1978) A review of the role of phytoalexins in disease resistance mechanisms. Prontificae Academiae Scientiarum Scripta Varia (21) IV : 1-61.
- Cruickshank, I.A.M. (1980) Defenses triggered by the invader : chemical defenses. Plant Disease : An Advanced Treatise, Vol. 5. (Eds. J.G.Horsfall and E.B. Cowling) pp. 247-267. Academic Press, London.
- Cruickshank, I.A.M. and Perrin, D.R. (1968) The isolation and partial characterization of monilicolin A, a polypeptide with phaseollin-inducing activity from Monilina fructicola. Life Sci. 7 : 449-458.
- Cubeta, M.A., Hartman, G.L. and Sinclair, J.B. (1985) Interaction between Bacillus subtilis and fungi associated with soybean seeds. Plant Dis. 69 : 506-509.
- Cullen, D. and Andrews, J.H. (1984) Evidence for the role of antibiosis in the antagonism of Chaetomium globosum to the apple scab pathogen, Venturia inaequalis. Can. J. Bot. 62 : 1819-1823.
- Cuypers, B., Schmelzer, E. and Hahlbrock, K. (1988) In situ localization of rapidly accumulated phenylalanine ammonia lyase mRNA around penetration sites of Phytophthora infestans in potato leaves. Mol. Plant Microb. Interaction 1 : 157-160.

- Daly, J.M. (1972) The use of near-isogenic lines in biochemical studies of the resistance of wheat to stem rust. Phytopathology 62 : 392-400.
- Daniel, M. (1995) Phytoalexins and other post-infectional compounds of some economically important plants of India. In : Handbook of Phytoalexin Metabolism and Action (Eds. M. Daniel and R.P. Purkayastha), pp. 533-553. Marcel Dekker Inc., New York.
- Daniels, D.L. and Hadwiger, L.A. (1976) Pisatin inducing components in filtrates of a virulent and avirulent Fusarium solani cultures. Physiol. Plant Pathol. 8 : 9-19.
- Darshika, P. and Daniel, M. (1992) Changes in the chemical content of Adhatoda and Trianthema due to fungal diseases. Indian J. Pharm. 54 : 73-75.
- Darvill, A.G. and Albersheim, P. (1984) Phytoalexins and their elicitors : a defense against microbial infection in plants. Ann. Rev. Plant Physiol. 35 : 243-275.
- Da Silva, S.R. and Pascholati, S.F. (1992) Saccharomyces cerevisiae protects maize plants, under green house conditions, against Colletotrichum graminicola. Journal of Plant Disease and Protection 99 : 159-167.
- Deverall, B.J. (1972) Phytoalexins and disease resistance. Proc. R. Soc. Lond. B. 181 : 233-246.
- Deverall, B.J. (1976) Current perspectives in research on phytoalexins. In : Biochemical Aspects of Plant-Parasite Relationship (Eds. J. Friend and D.R. Threlfall) pp. 207-223. Academic Press, New York.

- Dickens, J.S.W. and Cook, R.T.A. (1982) Glomerella cingulata on Camellia. Plant Pathol. 38 : 75-85.
- Dickinson, C.H. (1971) Cultural studies of leaf saprophyte. In : Ecology of Leaf Surface Microorganisms (Eds. T.F. Preece and C.H. Dickison) pp.129-137. Academic Press, London.
- Dixon, R.A. and Christopher, J.L. (1979) Stimulation of de novo synthesis of L-phenyl alanine ammonia lyase in relation to phytoalexin accumulation in Colletotrichum lindemuthianum elicitor-treated cell suspension cultures of French bean (Phaseolus vulgaris). Biochem. Biophys. Acta 586 : 453-463.
- Doherty, M.A. and Preece, T.F. (1978) Bacillus cereus prevents germination of uredospores of Puccinia allii and the development of rust disease of leek, Allium porrum in controlled environments. Physiol. Plant Pathol. 12 : 123-132.
- Douville, Y. and Boland, G.J. (1992) A note on the antibiotic properties of Bacillus subtilis against Colletotrichum trifolii. Phytoprotection 73 : 31-36.
- Dubos, B. (1987) Fungal antagonism in aerial agrobiocenoses. In : Innovative Approaches to Plant Disease Control. (Ed. I. Chet) pp. 107-135. New York.
- Ebel, J., Ayers, A.R. and Albersheim, P. (1976) Host pathogen interactions. XII. Response of suspension cultured soybean cells to the elicitor isolated from Phytophthora megasperma var. sojae, a fungal pathogen of soybean. Plant Physiol. 57 : 775-779.
- Ebel, J. and Grisebach, H. (1988) Defense strategies of soybean against the fungus Phytophthora megasperma f.sp. glycinea : molecular analysis. Trends in Biochemical Sciences 13 : 23-27.

- El-Abyad, M.S., El-Sayed, M.A., El-Shanshoury, A.R. and El-Sabbagh, S.M. (1993) Towards the biological control of fungal and bacterial disease of tomato using Streptomyces spp. Plant and Soil 149 : 189-195.
- Eswaran, K.S.S. (1971) Physiology of resistance to a bacterial disease in sorghum. Phytopath. Z. 71 : 141.
- Etenbarian, H.R. (1989) Studies on quantitative changes in phenolic compound on barley varieties during the development of Puccinia hordei and the relationship between these substance and brown rust resistance in barley. Physiol. Mol. Plant Pathol. 33 : 88-89.
- Facklam, R. and Wilkinson, H.W. (1981) The family Streptococcaceae (medical aspects). In : The Prokaryotes : A Handbook on Habitats, Isolation and Identification of Bacteria (Eds M.P. Starr, H.Q. Stolop, H.G. Trupper, A. Barlows and H.G. Schleyel) Vol. 2 : 1527-1597. Springer - Verlag, New York.
- Farkas, G.L. and Kiraly, Z. (1962) Role of phenolic compounds in the physiology of plant disease and disease resistance. Phytopathol. Z. 44 : 105-150.
- Farmer, E.E. (1985) Effects of fungal elicitor on lignin biosynthesis in cell suspension cultures of soybean. Plant Physiol. 78 : 338-342.
- Fernandez, M.R. and Heath, M.C. (1989) Interaction of the non-host French bean plant (Phaseolus vulgaris) with parasitic and saprophytic fungi. III. Cytologically detectable responses. Can. J. Bot. 67 : 676-686.

- Ferreira, J.H.S., Matthee, F.N. and Thomas, A.C. (1991) Biological control of Eutypa lata on grapevine by an antagonistic strain of Bacillus subtilis. Phytopathology 81 : 283-287.
- Fiala, V., Glad, C., Martin, M., Jolivet, E. and Derridj, S. (1990) Occurrence of soluble carbohydrates on the phylloplane of maize (Zea mays L.) : variations on relation to leaf heterogeneity and position on the plant. New Phytol. 115 : 609-615.
- Fiddaman, P.J. and Rossall, S. (1993) The production of antifungal volatiles by Bacillus subtilis. J. Appl. Bacteriol (Oxf.) 74 : 119-126.
- Fiddaman, P.J. and Rossall, S. (1994) Effect of substrate on the production of antifungal volatiles from Bacillus subtilis. Journal of Applied Bacteriology 76 : 395-405.
- Fischer, W., Christ, U., Baumgartner, M., Erismann, K.H. and Mosinger, E. (1989) Pathogenesis-related proteins of tomato II. Biochemical and immunological characterization. Physiol. Mol. Plant Pathol. 35 : 67-87.
- Fokkema, N.J. (1971) Influence of pollen on saprophytic and pathogenic fungi on rye leaves. In : Ecology of Leaf Surface Microorganisms (Eds. T.F. Preece and C.H. Dickinson) pp. 277-282. Academic, London.
- Fokkema, N.J. (1973) The role of saprophytic fungi in antagonism against Drechslera sorokiniana (Helminthosporium sativum) on agar plates and on rye leaves with pollen. Physiol. Plant Pathol. 3 : 195-205.
- Fokkema, N.J. (1981) Fungal leaf saprophytes, beneficial or detrimental ? In : Microbial Ecology of the Phylloplane. (Ed. J.P. Blakeman) pp. 433-454. Academic, London.

- Fokkema, N.J. (1993) Opportunities and problems of control of foliar pathogens with microorganisms. Pestic. Sci. 37 : 411-416.
- Fokkema, N.J., Dan Houter, J.S., Keslerman, T.J.C. and Nelis, A. (1979) Manipulation of yeasts on field grown wheat leaves and their antagonistic effect on Cochliobolus sativus and Septoria nodorum. Trans. Br. Mycol. Soc. 72 : 19-29.
- Fokkema, N.J., Van de Laar, J.A.J., Nelis-Blomberg, A.L. and Schippers, B. (1975) The buffering capacity of the natural mycoflora of rye leaves to infection by Cochliobolus sativus and its susceptibility to benomyl. Neth. J. Plant Pathol. 81 : 176-186.
- Fravel, D.R. (1988) Role of antibiosis in the biocontrol of plant diseases. Ann. Rev. Phytopathol. 26 : 75-91.
- Fridlender, M., Inbar, J. and Chet, I. (1993) Biological control of soil borne plant pathogens by a beta-1, 3-glucanase producing Pseudomonas cepacia. Soil Biol. Biochem. 25 : 1211-1221.
- Friend, J. (1977) Biochemistry of plant pathogens. Int. Rev. Biochem. 13 : 141-182.
- Friend, J. (1981) Plant phenolics, lignification and plant disease. Prog. Phytochem. 7 : 197-261.
- Fry, S.C. (1986) Cross-linking of matrix polymers in the growing cell walls of angiosperms. Ann. Rev. Plant Physiol. 37 : 165-186.
- Fry, S.C. (1987) Intracellular feruloylation of pectic polysaccharides. Planta 171 : 205-211.

- Gaber, B.A. and Hutchinson, T.C. (1988) Chemical changes in stimulated rain drops following contact with leaves of four boreal forest species. Can. J. Bot. 66 : 2445-2451.
- Gans, P.T. (1978) Physiological response of potato tubers to damage and to infection by Phoma exigua f.sp. foveata. Ann. Appl. Biol. 89 : 307-309.
- Gessler, C. and Kuc,¹ J. (1982) Appearance of a host protein in cucumber plants infected with viruses, bacteria and fungi. J. Exp. Bot. 33 : 58.
- Gianinazzi, S., Ahl, P., Cormu, A., Scalla, R. and Cassini, R. (1980) First report of host b-protein appearance in response to fungal infection in tobacco. Physiol. Plant Pathol. 16 : 337.
- Gordon, R.E., Haynes, W.C. and Pang, C.H.N. (1973). The genus Bacillus. Handbook No. 427. U.S. Dept. Agric., Washington D.C.
- Goyal, B.K., Verma, P.R. and Reddy, M.S. (1995) Suppression of white rust on mustard by foliar and seed application of microbial inoculants. In : Global conference on advances in research on plant diseases and their management, Udaipur, p. 152.
- Gustine, D.L., Sherwood, R.T., Moyer, B.G. and Lukezic, F.L. (1990) Metabolites from Pseudomonas corrugata elicit phytoalexin biosynthesis in white clover. Phytopathology 80 : 1427-1432.
- Gutterson, N., Howie, W. and Suslow, T. (1990) Enhancing efficiencies of biocontrol agents by use of biotechnology. UCLA Symp. Mol. Cell. Biol. pp. 749-756. Wiley-Liss, Inc., N.Y.
- Hachler, H. and Hohl, H.R. (1984) Temporal and spatial distribution patterns of collar and papillae wall appositions in resistant

and susceptible tuber tissue of Solanum tuberosum infected by Phytophthora infestans. Physiol. Plant Pathol. 24 : 107-118.

- Hahlbrock, K. and Scheel, D. (1989) Physiology and molecular biology of phenyl propanoid metabolism. Ann. Rev. Plant Physiol. Plant Mol. Biol. 40 : 347-369.
- Hait, G.N. and Sinha, A.K. (1986) Protection of wheat seedlings from Helminthosporium infection by seed treatment with chemicals. J. Phytopathol. 115 : 97-107.
- Hammerschmidt, R. (1984) Rapid deposition of lignin in potato tissue as a response to fungi non-pathogenic on potato. Physiol. Plant Pathol. 24 : 33-42.
- Hammerschmidt, R. and Nicholson, R.L. (1977) Resistance of maize to anthracnose : change in host phenols and pigments. Phytopathology 67 : 251-258.
- Handelsman, J., Raffel, S., Mester, E.H., Wunderlich, L. and Grau, C.R. (1990) Biological control of damping-off of alfalfa seedlings with Bacillus cereus UW85. Applied and Environmental Microbiology 56 : 713-718.
- Harborne, J.B. (1973) Phytochemical Methods. Chapman and Hall, London.
- Harborne, J.B. and Ingham, J.L. (1978) Biochemical Aspects of Plant and Animal Coevolution pp. 343-405. Academic Press, London.
- Harding, V.K. and Heale, J.B. (1981) The accumulation of inhibitor compounds in the induced resistance response of carrot root slices to Botrytis cinerea. Physiol. Plant Pathol. 18 : 7-15.
- Hassanein, F.M. and Elgoorani, M.A. (1991) The effect of Bacillus subtilis on in vitro growth and pathogenicity of Agrobacterium tumefaciens. Journal of Phytopathology 133 : 239-246.

- Heath, M.C. (1980) Reactions of non-suspects to fungal pathogens. Ann. Rev. Phytopathol. 18 : 211-236.
- Henderson, S.J. and Friend, J. (1979) Increase in PAL and lignin-like compounds as race-specific resistance responses of potato tubers to Phytophthora infestans. Phytopathol. Z. 94 : 323-334.
- Hodges, C.F., Campbell, D.A. and Christians, N. (1993) Evaluation of Streptomyces for biocontrol of Bipolaris sorokiniana and Sclerotinia homoeocarpa on the Phylloplane of Poa pratensis. J. Phytopathology 139 : 103-109.
- Hofmans, A.I. and Fuchs, A. (1970) Direct bioautography on thin-layer chromatograms as a method for detecting fungitoxic substances. Journal of Chromatography 51 : 327.
- Holliday, M.J., Keen, N.T. and Long, M. (1981) Cell death patterns and accumulation of fluorescent material in the hypersensitive response of soybean leaves to Pseudomonas syringae pv. glycinea. Physiol. Plant Pathol. 18 : 279.
- Ingham, J.L. (1972) Phytoalexins and other natural products as factors in plant disease resistance. Bot. Rev. 38 : 343-424.
- Ingham, J.L. (1973) Disease resistance in higher plants: the concept of pre-infectional and post-infectional resistance. Phytopathol. Z. 78 : 314-335.
- Ingham, J.L. (1982) Phytoalexins from the Leguminosae. In : Phytoalexins (Eds. J.A. Bailey and J.W. Mansfield) pp. 21-80. Blackie, London.
- Inglis, G.D. and Boland, G.R. (1990) The microflora of beans and rapeseed petals and the influence of the microflora of bean petals on white mold. Can. J. Plant Pathol. 12 : 129-134.

- Inglis, G.D. and Boland, G.R. (1992) Evaluation of filamentous fungi isolated from petals of bean and rapeseed for suppression of white mold. Can. J. Microbiol.
- James , D.W. Jr. and Gutterson, N.I. (1986) Multiple antibiotics produced by Pseudomonas fluorescens HV37 and their differential regulation by glucose. Appl. Environ. Microbiol. 52 : 1183-1189.
- Janisiewicz, W., Yourman, L., Roitman, J. and Mahoney, N. (1991) Post-harvest control of blue mold and gray mold of apples and pears by dip treatment with pyrrolnitrin, a metabolite of Pseudomonas cepacia. Plant Disease 75 : 490-494.
- Jayaswal, R.K., Fernandez, M., Upadhyay, R.S., Kurz, M., Webb, J. and Rinehart, K. (1993) Antagonism of Pseudomonas cepacia against phytopathogenic fungi. Curr. Microbiol. 26 : 17-22.
- Johnson, K.B., Stockwell, V.O., McLaughlin, R.J, Sugar, D., Loper, J.E. and Roberts, R.G. (1993) Effect of antagonistic bacteria on establishment of honey bee dispersed Erwinia amylovora in pear blossoms and on fire blight control. Phytopathology 83 : 995-1002.
- Johnson, L.B., Brammaman, B.L. and Zscheile, F.P. Jr. (1966) Protein and enzyme changes in barley leaves infected with Erysiphe graminis f. sp. hordei. Phytopathol. 56 : 1405.
- Joosten, M.H.A.J. and Dewit, P.J.G.M. (1988) Isolation, purification and preliminary characterization of a protein specific for compatible Cladosporium fulvum (Syn. Fulgia fulva) - tomato interaction. Physiol. Mol. Plant Pathol. 33 : 241-253.
- Kabir, S.E., Ghosh Hajra, N. and Chaudhuri, T.C. (1991) Performance of certain clones under agro-ecological conditions of Darjeeling. Technical Bulletin No. 5. Tea Board, Darjeeling Tea Research Centre, Kurseong, India.

- Kalaichelvan, P.T. and Mahadevan, A. (1988) Distribution of prohibitins in groundnut I. solvent effect and phenol estimation. Ind. Phytopathol. 41 : 581.
- Kauss, H., Franke, R., Krause, K., Conrath, U., Jeblick, W., Grimming, B. and Matern, U. (1993) Conditioning of parsley (Petroselinum crispum L.) suspension cells increase elicitor induced incorporation of cell wall phenolics. Plant Physiol 102 : 459-466.
- Kawamura, J. and Takeo, T. (1989) Antibacterial activity of tea catechin to Streptomyces mutans. Nippon Shokuhin Kogyo Gakkaishi 36 : 463.
- Keen, N.T. (1978) Phytoalexins : efficient extraction from leaves by a facilitated diffusion technique. Phytopathology 68 : 1237-1239.
- Keen, N.T. (1981) Evaluation of the role of phytoalexins. In : Plant Disease Control (Eds. R.C. Staples and G.H. Joerniesson) pp. 155-157. Wiley, N.Y.
- Keen, N.T. (1982) Specific recognition in gene-for-gene host parasite systems. Advances in Plant Pathology Vol. 1 (Eds. P.H. Williams and D. Ingram) pp. 35-82. Academic Press, N.Y.
- Keen, N.T. (1990) Gene-for-gene complimentary in plant-pathogen interactions. Ann. Rev. Genetics 24 : 447.
- Keen, N.T. and Bruegger (1977) Phytoalexins and chemicals that elicit their production in plants. In : Host Plant Resistance to Pests (Ed. P. Hedin) Am. Chem. Soc. Symp. Ser. 62 : 1-26.
- Keen, N.T. and Legrand, M. (1980) Surface glycoproteins : evidence that they may function as the race specific phytoalexin elicitors of Phytophthora megasperma f.sp. glycinea. Physiol. Plant Pathol. 17 : 175-192.

- Kempf, H.J. and Wolf, G. (1989) Erwinia herbicola as a biocontrol agent of Fusarium culmorum and Puccinia recondita f. sp. tritici on wheat. Phytopathology 79 : 990-994.
- Koomen, I. and Jeffries, P. (1993) Effects of antagonistic microorganisms on the post-harvest development of Colletotrichum gloeosporioides on mango. Plant Pathol. 42 : 230-237.
- Kranz, J. (1981) Hyperparasitism of biotrophic fungi. In : Microbial Ecology of the Phylloplane (Ed. J.P. Blakeman) pp. 327-352. Academic Press, London.
- Kuc, J. (1966) Resistance of plants to infectious agents. Ann. Rev. Microbiol. 20 : 337.
- Kuc, J. (1972) Phytoalexins. Ann. Rev. Phytopathol. 10 : 207-232.
- Kuc, J. (1976) Phytoalexins. Encyclopedia of Plant Physiology, Vol. 4. Physiological Plant Pathology (Eds. R. Heitefuss and P.H. Williams), pp. 637-652. Springer-Verlag, Berlin.
- Kumar, G.R., Ganesh, R. and Pathak, V.N. (1990) Some biochemical changes in the shoots of pearl millet infected Downey Mildew. Adv. Plant Sci. 5 : 103-108.
- Kumar, S., Shukla, H.S., Singh, H.N. and Singh, K.P. (1993) Isolation and characterization of mycelial wall polysaccharide of Drechslera longistrata eliciting glyceollins. Ind. Phytopathol. 46 : 267.
- Kumar, S., Singh, H.N. and Shukla, R.S. (1995) Host-pathogen interactions influencing the susceptibility of Drechslera rostrata causing leaf blight disease of Costus speciosus. In : Global Conference on Advances in Research on Plant Diseases and Their Management p. 43. Udaipur, India.

- Laemmli, U.K. (1970) Cleavage of structural proteins during assembly of the head of bacteriophage T4. Nature 227 : 680-685.
- Langcake, P. and MacCarthy, W.V. (1979) The relationship of resveratrol production to infection of grapevine leaves by Botrytis cinerea. Vitis 18 : 244-253.
- Langcake, P. and Pryce, R.J. (1976) The production of resveratrol by Vitis vinifera and other members of the Vitaceae as a response to infection or injury. Physiol. Plant Pathol. 9 : 77-86.
- Laville, J., Voisard, C., Keel, C., Maurhofer, M., Defago, G. and Haas, D. (1992) Global control in Pseudomonas fluorescens mediating antibiotic synthesis and suppression of black root rot of tobacco. Proc. Natl. Acad. Sci. (USA) 89 : 1562-1566.
- Legrand, M. (1983) Phenylpropanoid metabolism and its regulation in disease. In : Biochemical Plant Pathology (Ed. J.A. Callow) pp. 367-384. Wiley : Chichester.
- Lim, M., Uri, Z., Michael, N. and Hzhack, P. (1986) An improved procedure for the isolation of medicagenic acid 3-O- β -D-glucopyranoside from alfalfa roots and its antifungal activity on plant pathogens. J. Agric. Food Chem. 34 : 960.
- Lindow, S.E. (1979) Frost damage to potato reduced by bacteria antagonistic to ice nucleation-active bacteria. Phytopathology 69 : 1036 (Abstr.).
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951) Protein measurement with the Folin phenol reagent. J. Biol. Chem. 193 : 265.
- Luthra, Y.P. (1989) Total phenols and their oxidative enzymes in Sorghum leaves resistant and susceptible to Ramulispora sorghicola. Phytopath. 20 : 105-150.

- Lyons, P.C., Wood, K.V. and Nicholson, R.L. (1990) Caffeoyl ester accumulation in corn leaves inoculated with fungal pathogens. Phytochemistry 29 : 97-101.
- Macri, F., DiLerma, P. and Vianello. A. (1974) Preliminary research on peroxidase, polyphenol oxidase activity and phenol content in healthy and infected corn leaves, susceptible and resistant to Helminthosporium maydis race T. Rev. Patol. Vegetable 4 : 109.
- Mahadevan, A. (1966) Biochemistry of infection and resistance. Phytopath. Z. 57 : 96-99.
- Mahadevan, A. (1991) Post-infectional defense mechanisms. Biochemical aspects of plant disease resistance Vol. 2 : Today and Tomorrow's Printers and Publishers, New Delhi.
- Mahadevan, A. and Sridhar, R. (1982) Methods in Physiological Plant Pathology. 2nd Edn., Sivakami Publications, Madras.
- Maloney, S.E., Maule, A. and Smith, A.R.W. (1993) Purification and preliminary characterization of permethrinase from a pyrethroid transforming strain of Bacillus cereus. Appl. Environ. Microbiol. 59 : 2007-2013.
- Mansfield, J.W. (1983) Antimicrobial compounds. In : Biochemical Plant Pathology (Ed. J.A. Callow), pp. 237-265. Wiley : Chichester.
- Mansfield, J.W. (1990) Recognition and response in plant/fungus interactions. In : Recognition and Response in Plant Virus Interactions (Ed. R.S.S. Fraser), pp. 31-52. Springer-Verlag : Berlin.
- Mansfield, J.W., Hargreaves, J.A. and Boyle, F.C. (1974) Phytoalexin production by live cells in broad bean leaves infected with Botrytis cinerea. Nature 252

- Martins, E.M.F., de Maria, A.C., Grunewaldt-Stocker, G. and Moraes, W.B.C. (1986) Changes in the resistance of detached coffee leaves by yeast extract filtrate and heat treatment. Fitopatol. Bras. 11 : 899-909.
- Matern, U. and Kneusel, R.E. (1988) Phenolic compounds in plant disease resistance. Phytoparasitica 16 : 153-170.
- Matern, U., Strasser, H., Wendorff, H. and Hamerski, D. (1988) Coumarins and furanocoumarins. In : Cell Culture and Somatic Cell Genetics of Plants (Ed. I.K. Vasil), Vol. 5 : 3-21. Academic : New York.
- Maurhofer, M., Keel, C., Schinder, U., Voisard, C., Haas, D. and Defago, G. (1992) Influence of enhanced antibiotic production in Pseudomonas fluorescens strain CHAO on its disease suppressive capacity. Phytopathology 82 : 190-195.
- Mayama, S. and Tani, T. (1982) Microspectrophotometric analysis of the location of avenalumin accumulation in oat leaves in response to fungal infection. Physiol. Plant Pathol. 21 : 141-149.
- Mayama, S., Tani, T., Matsuura, Y., Ueno, T. and Fukami, H. (1981) The production of phytoalexins by oat in response to crown rust, Puccinia coronata f.sp. avenae. Physiol. Plant Pathol. 19 : 217-226.
- McDade, J.J. and Weaver, R.H. (1959) Rapid methods for the detection of gelatin hydrolysis. J. Bact. 77 : 60.
- McKeen, C.D., Reily, C.C. and Pusey, P.L. (1986) Production and partial characterization of antifungal substances antagonistic to Monilinia fructicola from Bacillus subtilis. Phytopathology 76 : 136-139.

- McKenzie, E.H.C. and Hudson, H.J. (1976) Mycoflora of rust-infected and non-infected plant material during decay. Trans. Br. Mycol. Soc. 66 : 223-238.
- Millar, R.J. and Higgins, V.J. (1970) Association of cyanide with infection of birds foot trefoil by Stemphylium loti. Phytopathology 50 : 104-110.
- Mishra, A. and Pundhir, V.S. (1995) Studies on potato phylloplane fungi. In : Global conference on advances in research on plant disease and their management, Udaipur, p. 159.
- Mishra, R.R. and Tewari, R.P. (1976) Studies on biological control of Puccinia graminis tritici. In : Microbiology of Aerial Plant Surfaces (Eds. C.H. Dickinson and T.F. Preece), pp. 559-567. Academic Press : London.
- Morris, C.E. and Rouse, D.I. (1985) Role of nutrients in regulating epiphytic bacterial populations. In : Biological Control on the Phylloplane (Eds. C.E. Windels and S.E. Lindow), pp. 63-82. St. Paul; MN . Am. Phytopathol. Soc.
- Mukerji, K.G. and Subba Rao, N.S. (1982) In : Advances in Agricultural Microbiology (Ed. N.S. Subba Rao), pp. 111-138. Oxford & I.B.H., New Delhi.
- Mukhopadhyay, R. . and Purkayastha, R.P. (1981) Effects of age and senescence retarding compounds on biological activities of leaf diffusates and disease susceptibility of rice. Ind. J. Exp. Biol. 21 : 216.
- Myatt, P.M., Dart, P.J. and Hayward, A.C. (1993) Potential for biological control of Phytophthora root rot of chickpea by antagonistic root-associated bacteria. Aust. J. Agric. Res. 44 : 773-784.

- Nakajima, T., Tamari, K. and Matsuda, K. (1977) A cell wall proteoheteroglycan from Pyricularia oryzae isolation and partial structure. J. Biochem. 82 : 1647-1655.
- Narain, A. and Mohanty, A.P. (1983) Bacterial antagonists of some phytopathogenic fungi. Indian J. Mycol. Plant Pathol. 13 : 28-31.
- Nicholson, R.L. and Hammerschmidt, R. (1992) Phenolic compounds and their role in disease resistance. Ann. Rev. Phytopathol. 30 : 369-389.
- Nicholson, R.L., Hipskind, J. and Hanau, R.M. (1989) Protection against phenol toxicity by the spore mucilage of Colletotrichum graminicola, an aid to secondary spread. Physiol. Mol. Plant Pathol. 35 : 243-252.
- Nieman, G.J., Van der Kerk, A., Niessen, M.A. and Versluis, K. (1991) Free and cell wall-bound phenolics and other constituents from healthy and fungus-infected carnation (Dianthus caryophyllus L.) stems. Physiol. Mol. Plant Pathol. 38 : 417-432.
- Norris, J.R., Berkeley, R.C.W., Logan, N.A. and O'Donnell, A.G. (1981) The genus Bacillus and Sporolactobacillus. In : The Prokaryotes : A Handbook on Habitats, Isolation and Identification of Bacteria (Eds. M.P. Starr, H. Stolp, H.G. Truper, A. Balows, H.G. Schlegel), 2 : 1711-1742. Springer-Verlag, N.Y.
- O'Brien, R.W. and Stern, J.R. (1969) Requirement for sodium in the anaerobic growth of Aerobacter aerogenes on citrate. J. Bact. 98 : 388.
- Oedjijono, Line, M.A. and Drager, C. (1993) Isolation of bacteria antagonistic to a range of plant pathogenic fungi. Soil Biol. Biochem. 25 : 247-250.

- Oke, O.A. (1988). Changes in chemical constituents of tobacco leaves infected with Caspicola corynespora and Colletotrichum nicotianae. J. Photopathol. 122 : 181-185.
- Omar, M. and Heather, W.A. (1979) Effect of saprophytic phylloplane fungi on germination and development of Melampsora larici - populina. Trans. Br. Mycol. Soc. 72 : 225-231.
- Omar, S.A.M., Salem, D.E. and El-Gantiry, S.M.M. (1989) Methods for estimating microorganism populations in faba bean phylloplane. FABIS Newsletter 24 : 16-20.
- Ouchi, S., Oku, H., Hibino, C. and Akiyama, I. (1974) Induction of accessibility and resistance in leaves of barley by some races of Erysiphe graminis. Phytopathol. Z. 79 : 24-34.
- Pandey, R.R. and Dwivedi, R.S. (1984) Seasonal incidence of phylloplane mycoflora of guava (Psidium guajava) with reference to fungal pathogens. Acta Bot. Indica 12 : 1-8.
- Parashar, R.D. and Sindhan, G.S. (1987) Biochemical changes in resistant and susceptible varieties of pea in relation to Powdery Mildew disease. Prog. Hortic. 18 : 137-138.
- Parent, J.G. and Asselin, A. (1984) Detection of pathogenesis related proteins (PR or b) and/or other proteins in the inter-cellular fluid of hyper-sensitive plants infected with tobacco mosaic virus. Can. J. Bot. 62 : 564-569.
- Partridge, J.E. and Keen, N.T. (1976). Association of the phytoalexin kievitone with single resistance of cowpeas to Phytophthora vignae. Phytopathology 66 : 426-429.
- Paschenko, I.N. (1978) Role of phenols in resistance of tobacco to blue mold. Mikol. Phytopathol. 12 : 343-346.

- Paxton, J.D. (1988) Phytoalexins in plant-parasite interactions. In : Experimental and Conceptual Plant Pathology (Eds. R.S. Singh, U.S. Singh, W.M. Hess and D.J. Weber) pp. 537. Oxford & I.B.H. Publishing Co. Pvt. Ltd., New Delhi.
- Paxton, J. (1995) Soybean phytoalexins : nature, elicitation, mode of action and role. In : Handbook of Phytoalexin Metabolism and Action (Eds.M. Daniel and R.P. Purkayastha) pp. 69-84. Marcel Dekker Inc., N.Y.
- Peng, G. and Sutton, J.C. (1991). Evaluation of microorganisms for biocontrol of Botrytis cinerea. Can. J. Plant Pathol. 13 : 247-257.
- Perumalla, C.J. and Heath, M.C. (1991) The effect of inhibitors of various cellular processes on the wall modifications induced in bean leaves by the cowpea rust fungus. Physiol. Mol. Plant Pathol. 38 : 293-300.
- Phae, C.G., Shoda, M., Kita, N., Nakano, M. and Ushiyama, K. (1992) Biological control of crown and root rot and bacterial wilt of tomato by Bacillus subtilis. Ann. Phytopathol. Soc. Jpn. 58 : 329-339.
- Pierce, M. and Ersenberg, M. (1987) Localization of phytoalexins in fluorescent mesophyll cells isolated from cotyledons and separated from other cells by fluorescence activated cell sorting. Physiol. Mol. Plant Pathol. 31 : 273.
- Powell, K.A. and Jutsum, A.R. (1993) Technical and commercial aspects of biocontrol products. Pestic Sci. 37 : 315-321.
- Prasada, B.K., Sinha, T.S.P. and Shankar, U. (1988) Biochemical changes in tomato fruits caused by Sclerotium rolfsii. Indian J. Bot. 14 : 122-129.

- Purkayastha, R.P. (1973) Phytoalexins : plant antigens and disease resistance. Sci. Culture 39 : 528-535.
- Purkayastha, R.P. (1976) Induction of phytoalexins in relation to plant disease control. Sci. & Cult. 42 : 586.
- Purkayastha, R.P. (1985) Phytoalexins. In: Frontiers in Applied Microbiology (Eds. K.G. Mukherjee, N.C. Pathak and V.P. Singh) Vol. 1, pp. 363-407. Print House, Lucknow.
- Purkayastha, R.P. (1986) Elicitors and elicitation of phytoalexins. In : Vistas in Plant Pathology (Eds. A. Varma and J.P. Verma) pp. 25-44. Malhotra Publishing House, New Delhi, India.
- Purkayastha, R.P. (1995) Progress in phytoalexin research during the past 50 years. In : Handbook of Phytoalexin Metabolism and Action (Eds. M. Daniel and R.P. Purkayastha) pp. 1-39. Marcel Dekker Inc., N.Y.
- Purkayastha, R.P. and Bhattacharyya, B. (1982) Antagonism of microorganisms from jute phyllosphere towards Colletotrichum corchori. Trans. Br. Mycol. Soc. 78 : 509-513.
- Purkayastha, R.P. and Ghosh, S. (1983) Elicitation and inhibition of phytoalexin biosynthesis in Myrothecium-infected soybean leaves. Indian J. Exp. Biol. 21 : 216-218.
- Purkayastha, R.P., Ghoshal, A. and Biswas, S. (1983) Production of momilactone associated with resistance of rice cultivars to sheath rot disease. Curr.Sci. 52 : 131-132.
- Purkayastha, R.P. and Ray, C. (1975) The detection of phytoalexins in jute leaves after infection by Colletotrichum corchorum and their possible role in lesion development. Physiol. Plant Pathol. 6 : 265-273.

- Purushothaman, D. (1974) Phenylalanine ammonialyase and aromatic amino acid in rice varieties infected with Xanthomonas oryzae. Phytopath. Z. 80 : 171-175.
- Pusey, P.L. (1991) Biological control of fruit rot. U.S. Dep. Agric. Pat. (5,047,239) p. 1.
- Pusey, P.L. and Wilson, C.L. (1984) Post-harvest biological control of stone fruit brown rot by Bacillus subtilis. Plant. Dis. 68 : 753-756.
- Ransom, R.F., Hipskind, J., Leite, B., Nicholson, R.L. and Dunkle, L.D. (1992) Effects of elicitor from Colletotrichum graminicola on the response of sorghum to Periconia circinata and its pathotoxin. Physiol. Mol. Plant Pathol. 41 : 75-84.
- Rao, K.V. and Manoharachary, C. (1985) Studies on pollen fungal association. Indian Bot. Rep. 4 : 20-22.
- Ricci, P., Trentin, F., Bonnet, P., Venard, P., Mouton-Perronnet and Bruneteau, M. (1992) Differential production of parasiticein, an elicitor of necrosis and resistance in tobacco, by isolates of Phytophthora parasitica. Plant Pathol. 41 : 298-307.
- Ride, J.P. (1983) Cell walls and other structural barriers in defence. In : Biochemical Plant Pathology (Ed. J.A. Callow) pp. 215-236. Wiley : Chichester.
- Roitman, J.N., Mahoney, N.E., Janisiewicz, W.J. and Benson, M. (1990) A new chlorinated phenylpyrrole antibiotic produced by the antifungal bacterium Pseudomonas cepacia. J. Agric. Food Chem. 38 : 538-541.

- Rosales, A.M., Vantomme, R., Swings, J., DeLey, J. and Mew, T.W. (1993) Identification of some bacteria from paddy antagonistic to several rice fungal pathogens. J. Phytopathol (Berl.) 138 : 189-208.
- Rothrock, C.S. and Gottlieb, D. (1981) Importance of antibiotic production in antagonism of selected Streptomyces species of two soil-borne plant pathogens. J. Antibiot. 34 : 830-835.
- Rouxel, T., Sarniguet, A., Kollmann, A. and Bousquet, J.F. (1989) Accumulation of a phytoalexin in Brassica spp. in relation to a hypersensitive reaction to Leptosphaeria maculans. Physiol. Mol. Plant Pathol. 34 : 507-517.
- Rytter, J.L., Lukezic, F.L., Craig, R. and Moorman, G.W. (1989) Biological control of geranium rust by Bacillus subtilis. Phytopathology 79 : 367-370.
- Saikia, P. and Chowdhury, H.D. (1993) Phylloplane microflora for the control of bacterial leaf blight of rice caused by Xanthomonas oryzae pv. oryzae. Ind. Phytopathol. 46 : 218-223.
- Sako, N. and Stahmann, M.A. (1972) Multiple molecular forms of enzymes in barley leaves infected with Erysiphe graminis f. sp. hordei. Physiol. Plant Pathol. 2 : 217.
- Sambandam, T., Sivswamy, N. and Mahadevan, A. (1982) Microbial degradation of phenolic substances. Ind. Rev. Life Sci. 2 : 1.
- Saxena, A.K., Sharma, A.K., Nityagopal, N. and Bakshi, A.K. (1986) Changes in total phenolics and sugars in relation to Puccinia recondita (brown rust) of some wheat varieties. J. Res. Punjab Agric. Univ. 24 : 359-362.

- Schonbeck, F. and Schlosser, E. (1976) Preformed substances as potential protectants. In : Physiological Plant Pathology (Eds. R. Heitefuss and P.H. Williams), Vol. 4 : 653-678. Springer-Verlag, New York.
- Schroth, M.N. and Hancock, J.G. (1981) Selected topics in Biological control. Ann. Rev. Microbiol. 35 : 453-476.
- Segrest, J.P. and Jackson, R.L. (1972) Molecular weight determination of glycoproteins by polyacrylamide gel electrophoresis in sodium dodecyl sulfate. In : Methods in Enzymology (Ed. V. Ginsburg) Vol. 28B : 54-63. Academic Press, New York.
- Sekhawat, P.S. and Chakravarti, B.P. (1977) Epiphytic bacterial and actinomycetes population and control of bacterial leaf spot of chillies by an antagonistic bacterium associated with bacterial lesions. Ind. Phytopathol. 30 : 358-360.
- Sen, S., Pal, S.C. and Sen, S.K. (1993) Characterization and medium suitability of extra-cellular amino acid producing strains of Micrococcus. Acta Microbiol. Pol. 41 : 75-82.
- Shahabuddin, M. and Anwar, M.N. (1992) Metabolic changes in the jute plant due to fungal infection. Bangladesh J. Bot. 21 : 205-212.
- Sharma, P.D. (1985) Potential of phylloplane microorganisms for biological control of rust diseases. In: Frontiers in Applied Microbiology 1 : 201-236.
- Shirashi, T., Oku, H., Yamashita, M. and Ouchi, S. (1978) Elicitor and suppressor of pisatin induction in spore germination fluid of pea pathogen Mycosphaeralla pinoides. Ann. Phytopathol. Soc. Jpn. 44 : 659-665.
- Simeoni, L.A., Lindsay, W.L. and Baker, R. (1987) Critical iron levels associated with biological control of Fusarium wilt. Phytopathology 77 : 1057-1061.

- Sinha, A.K. and Hait, G.N. (1982) Host sensitization as factor in induction of resistance in rice against Drechslera by seed treatment with phytoalexin inducers. Trans. Br. Mycol. Soc. 79 : 213-219.
- Skidmore, A.M. and Dickinson, C.J. (1976) Colony interactions and hyphal interference between Septoria nodorum and phylloplane fungi. Trans. Br. Mycol. Soc. 66 : 57-64.
- Smith, C.J., Milton, J.M. and Williams, J.M. (1995) Induction of Phytoalexin synthesis in Medicago sativa (Lucerne) - Verticillium albo-atrum interaction. In : Handbook of Phytoalexin Metabolism and Action (Eds. M. Daniel and R.P. Purkayastha) pp. 405-444. Marcel Dekker Inc., N.Y.
- Snary, D. and Hudson, L. (1979) Trypanosoma cruzi cell surface proteins : Identification of one major glycoprotein. FEBS Lett. 100 : 166-170.
- Spurr, H.W. Jr. (1981) Experiments on foliar disease control using bacterial antagonists. In : Microbial Ecology of the Phylloplane, pp. 369-381. Academic Press, London.
- Sridhar, R. and Ou, S.H. (1974) Phenolic compounds detected in rice blast disease. Biol. Plant. 16 : 67-70.
- Stahl, E. (1967) Dunnschicht-chromatographie. Springer-Verlag, Berlin-Heidelberg, New York.
- Stangarlin, J.R. and Pascholati, S.F. (1994) Protection of maize seedlings against Exserohilum turcicum by Saccharomyces cerevisiae. Summa Phytopathologica 20 : 16-21.
- Staples, R.C. and Stahmann, M.A. (1964) Changes in proteins and several enzymes in susceptible bean leaves after infection by the bean rust fungus. Phytopathol. 54 : 760.

- Stoessl, A. (1983) Secondary plant metabolites in preinfectious resistance. In : The Dynamics of Host Defence (Eds. J.A. Bailey and B.J. Deverall) pp. 71-122. Academic : New York.
- Sutton, J.C. and Peng, G. (1993) Biocontrol of Botrytis cinerea in strawberry leaves. Phytopathology 83 : 615-621.
- Tanaka, F., Takeuchi, S. and Tanaka, N. (1987) Bacimethrin, a new antibiotic produced by B. megatherium. The Journal of Antibiot. Ser. A. 14: 161.
- Tepper, C.S. and Anderson, A.J. (1986) Two cultivars of bean display a differential response to extra-cellular components from Colletotrichum lindemuthianum. Physiol. Mol. Plant Pathol. 29 : 411-420.
- Terce-Laforgue, T., Huet, J.C. and Pernollet, J.C. (1992) Biosynthesis and secretion of cryptogein, a protein elicitor secreted by Phytophthora cryptogea. Plant Physiol. 98 : 936-941.
- Thomashow, L.S., Weller, D.M., Bonsall, R.F. and Pierson, L.S. III (1990) Production of the antibiotic phenazine-1-carboxylic acid by fluorescent Pseudomonas species in the rhizosphere of wheat. Appl. Environ. Microbiol. 54 : 908-912.
- Thomson, S.V., Schroth, M.N., Moller, W.J. and Reil, W.O. (1976) Efficacy of bactericides and saprophytic bacteria in reducing colonization and infection of pear flowers by Erwinia amylovora. Phytopathology 66 : 1457-1459.
- Thornley, M.J. (1960) The differentiation of Pseudomonas from other Gram-negative bacteria on the basis of arginine metabolism. J. Appl. Bacteriol. 23 : 37-52.
- Tomiyama, K. (1966) Double infection by an incompatible race of Phytophthora infestans of a potato cell which has been

- previously infected by a compatible race. Ann. Phytopathol. Soc. Jpn. 32 : 181.
- Torre, G.D. and Tosi, L. (1989) Alteration in the composition of phenolic compounds and amino acids in tobacco seedlings infected with Thielaviopsis basicola. Ann. Fac. Agrar. Univ. Stud. Perugia 43 : 279-288.
- Toyoda, K., Kobayashi, I. and Kunoh, H. (1992) Elicitor activity of a fungal product assessed at the single-cell level by a novel gel-bead method. Plant and Cell Physiol. 34: 775-780.
- Tronsmo, A. (1986) Use of Trichoderma spp. in biological control of necrotrophic pathogens. Microbiology of the phyllosphere (Eds. N.J. Fokkema and J. Vanden Heuvel), pp. 348-362. Cambridge University Press, Cambridge.
- Tschen, J.S.M. and Kuo, W.L. (1985) Antibiotic inhibition and control of Rhizoctonia solani by Bacillus subtilis. Plant Prot. Bull. Taiwan 27 : 95-103.
- Tsuji, H., Tsuji, M. and Deba, H. (1990a) Antifungal activity of the culture supernatant of Bacillus spp. and examination of the culture medium compositions. J. Pestic. Sci. 15 : 1-6.
- Tsuji, H., Tsuji, M., Deguchi, T. and Deba, H. (1990b) Properties, isolation and purification of the antifungal substances produced by Bacillus spp. J. Pestic. Sci. 15 : 7-12.
- Uritani, I. (1971) Protein changes in diseases plants. Ann. Rev. Phytopathol. 9 : 211.
- Uritani, I. (1976) Protein metabolism. In : Encyclopedia of Plant Physiology Vol.4 Physiological Plant Pathology (Eds. R. Heitefuss and P.H. Williams) pp. 509. Springer-Verlag, Berlin, Heidelberg, New York.

- Uritani, I. and Stahman, M.A. (1961a) Changes in nitrogen metabolism in sweet potato with black rot. Plant Physiol. 36 : 770.
- Uritani, I. and Stahman, M.A. (1961b) The relationship between antigenic compounds produced by sweet potato in response to black rot infection and the magnitude of disease resistance. Agr. Biol. Chem. 25 : 479.
- Ushiyama, K. (1980) Studies on citrus melanose disease of Satsuma mandarin-5 influence of microorganism isolated from citrus orchard on the occurrence of citrus melanose by Diaporthe citri. Bull. Kanagawa Hortic. Exp. Stn. 0 : 9-16.
- Utkhede, R.S. and Rahe, J.E. (1983) Interactions of antagonist and pathogen in biological control of onion white rot. Phytopathology 72 : 890-893.
- Valdes S.N., Gotz, H. and Busse, M. (1991) Antagonistic effect of coryneform bacteria from red smear cheese against Listeria species. Int. J. Food Microbiol. 13 : 119-130.
- VanEtten, H.D. (1973) Differential sensitivity of fungi to pisatin and to phaseollin. Phytopathology 63 : 1477-1482.
- VanEtten, H.D., Matthews, D.E. and Matthews, P.S. (1989) Phytoalexin detoxification : importance for pathogenicity and practical implications. Ann. Rev. Phytopathol. 27 : 143-164.
- VanEtten, H.D., Matthews, D.E. and Smith, D.A. (1982) Metabolism of phytoalexins. In : Phytoalexins. (Eds. J.A. Bailey and J.W. Mansfield) pp. 181-217. Blackie, Glasgow.
- VanEtten, H.D. and Pueppke, S.G. (1976) Isoflavonoid phytoalexins. In : Biochemical aspects of Plant-Parasite Relationship (Eds. J. Friend and D.R. Threlfall) pp. 230-289. Academic Press, New York.

- Varneste, J.L., Yu, J. and Beer, S.V. (1992) Role of antibiotic production by Erwinia herbicola Eh 252 in biological control of Erwinia amylovora. J. Bacteriol. 174 : 2785-2796.
- Van Sumere, C.F., Wolf, G., Teuchy, H. and Kint, J. (1965) A new thin-layer method for phenolic substances and coumarins. J. Chromat. 20 : 48-65.
- Vaughn, S.F. and Lulai, F.C. (1991) The involvement of mechanical barriers in the resistance response of a field-resistant and a field-susceptible potato cultivar to Verticillium dahliae. Physiol. Mol. Plant Pathol. 38 : 455-465.
- Vidyasekharan, P. (1988) Phenolics and disease resistance. In : Physiology of Disease Resistance in Plants I. (Ed. P. Vidyasekharan) pp. 49. CRC Press, Boca Raton.
- Viswanathan, R. and Narayansamy, P. (1991) Nature of rice phylloplane micro-organisms and their sensitivity to fungicides. Acta Botanica Indica 19 : 236-238.
- Walker, R., Powell, A.A. and Seddon, B. (1994) Tests for biological control of seed and seedling damping-off disease of peas and beans using Bacillus species. BCPC MONOGRAPH : Seed treatment : Progress and Prospects 57 : 333-338.
- Wang, Z.N. (1991) Chinese famous teas and their characteristic constituents bioformation World Tea. International Symposium on Tea Science. Shizuoka, Japan, pp. 23.
- Ward, E.W.B. (1986) Biochemical mechanisms involved in resistance of plants to fungi. In : Biology and Molecular Biology of Plant Pathogen Interactions (Ed. J. Bailey) pp. 107-131. Springer-Verlag, Berlin, Heidelberg.

- Weller, D. and Thomashow, L.S. (1990) Antibiotics : evidence for their production and sites where they are produced. In : New Directions in Biological Control : Alternatives for Suppressing Agricultural Pests and Diseases (Eds. R.R. Baker and P.E. Durr) pp.703-711. Liss, N.Y.
- Werder, J. and Kern, H. (1985) Resistance of maize to Helminthosporium carbonum : changes in host phenolics and their antifungal activity. Journal of Plant Diseases and Protection 92 : 477-484.
- Whipps, J.M., McQuilken, M.P. and Budge, S.P. (1993) Use of fungal antagonists for biocontrol of damping-off and Sclerotinia diseases. Pestic Sci. 37 : 309-313.
- Williams, D.H. and Fleming, I. (1988) Spectroscopic methods in organic chemistry. Tata McGraw Hill Publishing Co., New Delhi.
- Williams, S.T. (1982) Are antibiotics produced in soil ? Pedobiologia 23 : 427-435.
- Wilson, M., Epton, H.A.S. and Sigeo, D.C. (1990) Biological control of fireblight of hawthorn (Crataegus monogyna) with Erwinia herbicola under protected conditions. Plant Pathology 39 : 301-308.
- Wilson, M. and Lindow, S.E. (1993) Interactions between the biological control agent Pseudomonas fluorescens A506 and Erwinia amylovora in pear blossoms. Phytopathology 83 : 117-123.
- Wölk, M. and Sarkar, S. (1993) Antagonism in vitro of fluorescent pseudomonads against Rhizoctonia solani and Pythium aphanidermatum. Zentralbl Microbiol. 148 : 237-245.
- Wood, R.K.S. (1982) Active defense mechanisms in plants. Plenum, New York.

- Yamoto, Y., Oku, H., Shirashi, T., Ouchi, S. and Koshizawa, K. (1986) Non-specific induction of pisatin and local resistance in Pea leaves by elicitors from Mycosphaerella pinoides, M. melonis and M. lignilicola and the effect of suppressor from M. pinoids Jr. Phytopathology 117 : 136.
- Yanase, Y. and Takeda, Y. (1987) Method for testing the resistance to tea grey blight caused by Pestalotia longiseta spegazzini in tea breeding. Bulletin of the National Research Institute of Vegetables, Ornamental Plants and Tea B. (Kanaya) 1 : 1.
- Ye, X.S., Pan, S.Q. and Kuc', J. (1990) Association of pathogenesis related proteins and activities of peroxidase, β -1, 3-glucanase and chitinase with systemic induced resistance to blue mould of tobacco but not to systemic tobacco mosaic virus. Physiol. Mol. Plant Pathol. 36 : 523-531.
- Yoshikawa, M. (1983) Macromolecules, recognition and the triggering of resistance. In : Biochemical Plant Pathology (Ed. J.A. Callow) pp. 267-298. John Wiley & Sons, N.Y.
- Yoshikawa, M. (1995) Elicitor and the molecular bases of phytoalexin elicitation. In : Handbook of Phytoalexin Metabolism and Action (Eds. M. Daniel and R.P. Purkayastha) pp. 61-68. Marcel Dekker Inc., N.Y.
- Yoshikawa, M. and Sugimoto, K. (1993) A specific binding site on soybean membranes for a phytoalexin elicitor released from fungal cell walls by β -1,3-endoglucanase. Plant and Cell Physiol. 34 : 1229-1247.
- Yoshikawa, M., Yamaoka, N. and Takeuchi, Y. (1993) Elicitors : their significance and primary modes of action in the induction of plant defense reactions. Plant and Cell Physiol. 34:1163 - 1173.

- Zaher, E., Barakat, F.M., Osman, A.R. and El-Khaleely, M.I. (1985) Antagonism between phyllosphere bacteria and actinomycetes and Ulocladium botrytis Preuss. causing tomato leaf spot. Egypt J. Phytopathol. 17 : 15-22.
- Zhou, T. and Reeleder, R.D. (1989) Application of Epicoccum purpurascens spores to control white mold of snap bean. Plant Dis. 76 : 639-642.
- Zhou, T. and Reeleder, R.D. (1991) Colonization of bean flowers by Epicoccum purpurascens. Phytopathology 81 : 774-778.