

CONTENTS

| CHAPTER - I | Page No. |
|--|-----------------|
| 1. INTRODUCTION | 1 - 5 |
| CHAPTER - II | |
| 2. REVIEW OF LITERATURE | 6 -17 |
| 2.1 Impact of mulberry varieties on the bivoltine silkworm rearing. | 6 -7 |
| 2.1.1. Quality of leaves. | 7 -10 |
| 2.2. Silkworm breed and rearing seasons. | 10 -13 |
| 2.3. Effect of fertilizer on mulberry plant and bearing upon silkworm rearing performance. | 13-15 |
| 2.4. Nutritional efficiencies. | 15-17 |
| CHAPTER - III | |
| 3. MATERIALS AND METHODS | 18-26 |
| 3.1. Geographical location , soil and climate. | 18 |
| 3.2. Experimental site | 19 |
| 3.3. Insect and food plant. | 19 |
| 3.4. Rearing of silkworm. | 19-20 |
| 3.4.1. Meteorological data or rearing room. | 20 |
| 3.4.2. Diet. | 21 |
| 3.5. Rearing performance. | 22 |
| 3.6. Consumption and utilization of food by the fifth instar larva. | 22 |
| 3.6.1. Consumption and growth. | 23 |
| 3.6.2. Digestibility and efficiency of conversion. | 23 |
| 3.7. Reproduction performance. | 24 |

| | | |
|--------|---|-------|
| 3.8. | Reeling characters of cocoon and filament characters of silk. | 24 |
| 3.9. | Biochemical assay of mulberry leaves. | 24 |
| 3.9.1. | Estimation of total protein. | 24-25 |
| 3.9.2. | Estimation of total carbohydrate | 25-26 |
| 3.9.3. | Determination of leaf moisture | 26 |
| 3.10. | Statistical analysis. | 26 |

CHAPTER - IV

| | | |
|----------|---|--------------|
| 4. | RESULTS | 27-58 |
| 4.1. | Screening of mulberry varieties . | 27 |
| 4.1.1. | Total leaf yield . | 27 |
| 4.1.2. | Chemical contents of leaf. | 27-28 |
| 4.1.3. | Rearing performance. | 28 |
| 4.2. | Selection of superior variety from the initially screened varieties. | 28-29 |
| 4.2.1. | Seasonal variations of chemical contents in mulberry leaves of different varieties. | 29 |
| 4.2.1.1. | Total protein content. | 29 |
| 4.2.1.2. | Total carbohydrate content. | 29-30 |
| 4.2.1.3. | Moisture content. | 30 |
| 4.2.2. | Rearing performance of worm as influenced by variety over seasons. | 30 |
| 4.2.2.1. | Weight of 10 mature larvae. | 30-31 |
| 4.2.2.2. | Single cocoon weight. | 31-32 |
| 4.2.2.3. | Single shell weight. | 32-33 |
| 4.2.2.4. | Shell ratio. | 33-34 |
| 4.2.2.5. | Absolute silk content. | 34-35 |
| 4.2.2.6. | Effective rate of rearing in weight . | 35-36 |

| | | |
|----------|--|-------|
| 4.2.2.7. | Effective rate of rearing in number. | 36-37 |
| 4.2.2.8. | Fecundity. | 37 |
| 4.2.3. | Quality of cocoon as influenced by variety. | 38 |
| 4.2.3.1. | Filament length. | 38 |
| 4.2.3.2. | Reelability. | 38 |
| 4.2.3.3. | Denier. | 38-39 |
| 4.2.4. | Consumption and utilization efficiencies of food as influenced by variety over seasons. | 39 |
| 4.2.4.1. | Consumption indices. | 39-40 |
| 4.2.4.2. | Nutritional efficiencies. | 40-42 |
| 4.3. | Enrichment of nutritional quality of leaves through manipulation of fertilizer supplementation and its impact on silkworm. | 42 |
| 4.3.1. | Nutrient contents of leaves during fertilizer supplementation to S1 mulberry leaves. | 42 |
| 4.3.1.1. | Total protein content. | 42-43 |
| 4.3.1.2. | Total carbohydrate content. | 43 |
| 4.3.1.3. | Moisture content. | 43-44 |
| 4.3.2. | Effect of fertilizer supplementation on rearing performance of different breeds over seasons | 44 |
| 4.4.2.1. | Weight of 10 mature larvae. | 44-45 |
| 4.3.2.2. | Single cocoon weight. | 45-46 |
| 4.3.2.3. | Single shell weight. | 46-47 |
| 4.3.2.4. | Shell ratio. | 47-48 |
| 4.3.2.5. | Absolute silk content. | 48-49 |
| 4.3.2.6. | Effective rate of rearing in weight. | 49-50 |
| 4.3.2.7. | Effective rate of rearing in weight. | 50-51 |
| 4.3.2.8. | Fecundity. | 51 |

| | | |
|----------------------|--|--------------|
| 4.3.3. | Effect of fertilizer supplementation on cocoon quality. | 51 |
| 4.3.3.1. | Filament length. | 51-52 |
| 4.3.3.2. | Reelability. | 52 |
| 4.3.3.3. | Denier. | 52-53 |
| 4.3.4. | Effect of fertilizer supplementation on consumption and utilization of food in different breeds over seasons. | 53 |
| 4.4.4.1. | Consumption indices. | 53-55 |
| 4.3.4.2. | Nutritional efficiencies. | 55-56 |
| 4.3.5. | Total leaf yield. | 56 |
| 4.4. | Impact of different combinations of two mulberry varieties as food on consumption and utilization and rearing performance of silkworm larvae. | 56 |
| 4.4.1. | Consumption and utilization of food. | 56 |
| 4.4.1.1. | Consumption indices. | 56-57 |
| 4.4.1.2. | Nutritional efficiencies. | 57-58 |
| 4.4.2. | Rearing performance. | 58 |
| CHAPTER - V | | |
| 5. | DISCUSSION | 59-69 |
| 5.1. | Screening. | 59-61 |
| 5.2. | Selection of the best variety from the initially screened three varieties after systematic evaluation. | 61-65 |
| 5.3. | Effect of fertilizer enrichment of mulberry crop on leaf quality and its impact on rearing performance , fecundity , quality of silk and nutritional efficiencies. | 65-69 |
| 5.4. | Effect of combination of leaves of two different varieties. | 69 |
| CHAPTER - VI | | |
| 6. | SUMMARY | 70-73 |
| CHAPTER - VII | | |
| 7. | BIBLIOGRAPHY. | 74-88 |