

FIGURE INDEX

Fig. No.	Title	Page
1	Percentage of hemoglobin in mouse blood after intravenous injection of EEA and ethanol	29
2	Count of RBC in mouse blood after i.v. injection of EEA and ethanol	30
3	Total count of WBC after intravenous injection of EEA and ethanol	30
4	Survivality of lymphocytes <i>in vitro</i> in presence and absence of EEA	31
5	<i>In vitro</i> blastogenic transformation of T and B cells in presence and absence of EEA	32
6	Pattern of incorporation of ³ H-thymidine by lymphocytes treated for different hours with EEA	33
7	DNA histograms representing data of one experiment of cell cycle analysis of splenic lymphocytes upon <i>in vitro</i> treatment with EEA and alcohol	35
8	Inhibitory effect of topical application of EEA on delayed type hypersensitive (DTH) reaction induced with 25 µl 2,4-dinitro-fluorobenzene (DNFB)	36
9	Indurations of DTH swelling, induced with 50 µl DNFB, and inhibition after topical application of EEA and alcohol	37
10	Photographs showing effect of topical application of EEA and alcohol on resensitized paw of DTH mice, induced by 50 µl DNFB, during 9 day period of study	38
11	Changes in paw size of DTH mice, induced with 25 µl DNFB, and prior i.v. application of EEA and alcohol	39
12	Changes in paw size of DTH mice, induced with 50 µl DNFB, and after i.v. application of EEA and alcohol	40
13	Percentage of different leukocytes at the inflammation site (resensitized paw) with EEA treatment (1 hr prior to resensitization)	41

14	Effect of intravenously administered EEA and alcohol on count of CD4 ⁺ T cells from spleen of mice after 24, 48 and 72 hrs of induction of DTH with 25 μ l DNFB	41
15	Level of serum TNF- α from DTH mice (induced with 25 μ l DNFB) at three different hours in presence and absence of EEA	42
16	<i>Expression of certain inflammation related genes in T cells obtained from DTH mice, and with and without EEA treatment, containing 9 mice in each group. Agarose gel electrophoresis of cDNA PCR products amplified with different gene specific primers</i>	43
17	Graphical representation of quantified PCR products for the expression of different genes in splenic T cells of DTH mice with EEA treatment compared to necessary controls	44
18	Healing of linear skin incision in mice in presence and absence of EEA	45
19	Healing of burnt skin in mice with and without EEA treatment	46
20	EEA inhibits solid tumor growth and increases longevity of tumor bearing mice upon topical application of the extract	47
21	Percentage of cytotoxic response by EEA treated lymphocytes after 6hrs incubation with labeled (⁵¹ Cr) tumor target cells	48
22	Expression of certain immunologically active genes in T cells from mice with and without EEA treatment, containing 9 mice in each group. Agarose gel electrophoresis of cDNA PCR products amplified with different gene specific primers	49
23	Graphical presentation of quantification of the expression of the immunologically active genes in T cells from mice with and without EEA treatment, containing 9 mice in each group. The DNA bands were quantified against 30 ng of lambda DNA through ImageAide, Spectronics Corporation, NY	50

24	No. of plaque forming cells (PFCs) in 10^6 lymphocytes from spleen of mice treated with EEA in Primary and Secondary immune response.	52
25	Estimation of serum IgG level (pg/ml) by ELISA in secondary immune response with EEA treatment, alcohol (control) and no treatment (only SRBC)	52
26	Inhibition of superoxide radical generation by EEA	53
27	EEA inhibits lipid peroxidation in mouse splenocytes	54
28	Inhibition of hydroxyl radical generation by EEA	55
29	Generation of nitric oxide in murine lymphocytes in presence of EEA and alcohol	56
30	HPLC analysis of total EEA extract showing a major peak at 468 nm	57
31	IR spectroscopic analysis of the major fraction from EEA	58
32	Mass spectroscopic analysis of the major constituent isolated from EEA	59
33	<i>In vitro</i> blastogenic transformation of T and B cells after treatment with the major fraction from EEA and compared to that of EEA total extract	60
34	Inhibition of superoxide radical by the fraction of EEA and compared to that of EEA total extract	61
35	Inhibition of lipid peroxidation by fraction from EEA and compared to total EEA extract	61
36	Inhibition of hydroxyl radical generation by fraction from EEA and total EEA	62
37	Generation of NO [•] in mouse splenocytes treated with total EEA and fraction from EEA	62
38	Effect of EEA on inhibition of DTH reaction and repair mechanism presented schematically	68
39	Summation of inflammation related events leading to tissue repair or promotion of tumor	70

40 Immunopharmacological properties of EEA compared with certain other plant extracts. Indices represent the level of efficacy of the extracts as ratio over control; the indices presented at the top of the column for easy representation. Any index ratio of 1.2 or more has been considered to have positive effect. Data for the natural products from the plant species other than *E. adenophorum* compiled on the basis of data in published literature.