

# **REFERENCES**

- Aguelon, M and Dunez, J. (1984). Immunoenzymatic techniques for the detection of *Phoma exigua* in infected potato tissues. *Ann. Appl. Biol.* **105**: 463-469.
- Alam, M., Sattar, A. and Janardhanan, K. K. (1991). Changes in phenol and peroxidase in the leaves of Java citronella infected with *Curvularia andropoyonis*. *Biol. Planta.*, **33**: 211-215.
- Alba, A.P.C. and De Vay, J.E. (1985). Detection of cross-reactive antigens between *Phytophthora infestans* (Mont.) de Bary and Solanum species by indirect enzyme-linked immunosorbent assay. *Phytopathology* **112**: 97-104.
- Alba, A.P.C., Guzzo, S.D., Mahlow, M.F.P. and Moreas, W.B.C. (1983). Common antigens in extracts of *Hemilia vastatrix* uredinospores and of *Coffea arabica* leaves and roots. *Fitopathologia Brasileria* **8**: 473-483.
- Amos, R.E. and Burrell, R.G. (1966). Serological differentiation in *Ceratocystis*. *Phytopathology* **57** : 32-34.
- Antoniw, J. F. White, R.F., Barbara, D.J., Jones, P. & Longley, A. (1985). The detection of PR (b) – protein and TMV by ELISA in systemic and localized infections of tobacco. *Plant Mol. Biol* **4**: 50-60.
- Arie- Tsutomu, (1998). Immunology-based diagnostics for soil borne diseases. *J. Pest. Sci.* **23** (3): 349-356.
- Barlansky, R.H. Lee, R.F., Timmer, L.W., Purcifull, D.E. & Raju, B.C. (1982). Immunofluorescent detection of xylem limited bacteria *in situ*. *Phytopathology* **72**: 144-48.
- Benhamou, N., Ouellette, G.B., Gardiner, R.B. and Day, A.W. (1986). Immunocytochemical localization of antigen binding sites in the cell surface of two ascomyete fungi using antibodies produced against fimbriae from *Ustilago violacea* and *Rhodotorula rubra*. *Can.J.Microbiol.* **32**: 871-883.
- Benson. B.M. (1991). Detection of *Phytophthora cinnamomi* in Azalea with commercial serological assay kits. *Plants Dis.* **75** : 478-482.
- Bhattacharya, M. K. and Ward, E. W.B. (1987). Temperature induced susceptibility of soybeans to *Phytophthora megasperma* f.sp. *glycinear*; phenylalanine

ammomia lyase and glyceollin sensitivity of the pathogen. *Physiol. Mol. Plant Pathol.*, **31**: 407-419.

- Biswas, K.K. and Sen, C. (2000).** Management of stem rot of groundnut caused by *Sclerotium rolfsii* through *Trichoderma harzianum*. *Indian Phytopathol.* **53**: 290-295.
- Brill, L.M., Mc Clary, R. and Sinclair, J.B. (1994).** Analysis of two ELISA formats and antigen preparations using polyclonal antibodies against *Phomopsis longicolla*. *Phytopathology* **84** : 173-179.
- Brill, L.M., Mc Clary, R. and Sinclair, J.B. (1994).** Analysis of two ELISA formats and antigen preparations using polyclonal antibodies against *Phomopsis longicolla*. *Phytopathology* **84** : 173-179.
- Bruezzese, E. and Hasan, S. (1991).** A whole clearing and staining technique For host specificity studies of rust fungi. *Plant Pathol.* **32**: 335-338.
- Burrell, R.G., Clayton C.W., Gallegly M.E. and Lilly, V.G. (1966).** Factors affecting the antigenicity of the mycelium of three species of *Phytophthora*. *Phytopathology* **56** : 423-426.
- Chakraborty, B. N. (2005).** Antimicrobial proteins in Plant Defence. In: *New Perspectives in the Frontiers of Chemical Research*. (Ed.S.S. Chakravorti) Royal Society of Chemistry, EIS, Kolkata pp. 470-483.
- Chakraborty, B.N. (1988)** Antigenic disparity. In: *Experimental and Conceptual Plant Pathology* (Eds. R.S. Singh, U.S. Singh W.M. Hess and D.J. Weber) P.477, Oxford and IBH publishing Co. Pvt. Ltd; New Delhi.
- Chakraborty, B. N. and Chakraborty, U. (2003).** Immunodetection of plant pathogenic fungi. In: *Frontiers of Fungal Diversity in India* ( Eds. G. P. Rao, C. Manoharachari, D. J. Bhat, R. C. Rajak and T. N. Lakhanpal ) International Book Distributing Co. Lucknow p. 23-42.

- Chakraborty, B.N., Chakraborty, U., Das, R., Basu, P. and Saha, A. (1996). Serological relationship between *Glomerella cingulata* ( Stoneman ) Spauld & Schrenk and *Camellia sinensis* (L.) O. Kuntze. *J.Plantation Crops* 24 : 205-211.
- Chakraborty, B.N., Chakraborty, U., Deb, D., Das, J., and Sengupta, D. (2002a). Commercialization of *Trichoderma*-based biofungicides in the management of tea root disease. In: Proceedings of National Symposium on “*Diversity of Microbial Resource and their Potential Application*,” Department of Botany, University of North Bengal, p: 37-38.
- Chakraborty, U., Chakraborty, B. N. and Kappor, M. (1993). Changes in the levels of peroxidase and phenylalanine ammonia-lyase in *Brassica napus*. cultivars showing variable resistance to *Leptosphaeria maculans*. *Folia Microbiol.*, 38: 491-496.
- Chakraborty, B.N, Chakraborty, U, Saha, A, Das, R and Basu, P. (1995). Detection and management of blister blight of tea. In: *Management of threatening plant diseases of National importance*. 227-235.
- Chakraborty, B.N., Chakraborty U., Sengupta, D., Deb, D. and Das. J. (2002). Development of immunodiagnostic Kits for detection of *Ustilina zonata* in the soil and tea root tissues. *J. Basic Appl. Mycol.* 1: 58-61.
- Chakraborty, B. N., Das Biswas, R. and Sharma, M. (2004). Multicomponent coordinated defence strategies in tea plants against *Helopeltis theivora* and *Exobasidium vexans*. *Journal of Plantation Crops*, 32: 289-297.
- Chakraborty, B.N., Das, R., Basu, P. and Chakraborty U. (2000). Immunodiagnosis of *Glomerella cingulata* causing brown blight of tea. Proceedings of International Conference on Integrated Plant Diseases Management for Sustainable Agriculture. *Indian Phytopathological Society IARI, New Delhi*, Vol. 1. 482-483.
- Chakraborty, U., Das, R., Basu, P., Guha, S. and Chakraborty, B. N. (2002). Serological cross reactivity between *Glomerella singulata* and *Camellia sinensis*. *J. Plants. Crops.*, 55(1): 1-7.

- Chakraborty, U., Deb, D., Das, J., Sengupta, D. and Chakraborty, B. N. (2001). Development of polyclonal antibody based immunoassay for detecting *Fomes lamoensis* causing brown rot disease of tea. *J. Mycol. Plant Pathol.* **31(1)**: 118-119.
- Chakraborty, B.N. and Purkayastha, R.P. (1983). Serological relationship between *Macrophomina phaseolina* and soybean cultivars. *Physiol. Plant Pathol.* **23**: 197-205.
- Chakraborty, B.N. and Purkayastha, R.P. (1987). Alternation in glyceollin synthesis and antigenic patterns after chemical induction of resistance in soybean to *Macrophomina phaseolina*. *Can. J. Microbiol.* **33**: 825-840.
- Chakraborty, B. N., and Saha, A. (1994). Detection and cellular location of cross reactive antigens shared by *Camellia sinensis* and *Bipolaris carbonum*. *Physiol. Mol. Plant Pathol.* **44** : 403-416.
- Chakraborty, B.N., and Saha, A. (1995). Detection of antifungal compounds in tea leaves after infection with *Bipolaris carbonum* In: *Recent advances in Phytopathological Research*. (Eds. A. K. Roy and K. K. Sinha). P. 205-211. M.D. Publication, New Delhi.
- Chakraborty, U., Sarkar, B. and Chakraborty, B. N. (2003). Protection of soybean root rot by *Bradirhizobium japonicum* and *Trichoderma harzianum* associated changes in enzyme activities and phytoalexins production. *J. Mycol. Pl. Pathol.*, **33(1)**: 21-25.
- Chakraborty, B.N., Sarkar, B. and Chakraborty, U.(1997). Detection of cross-reactive antigens shared by *Fusarium oxysporum* and *Glycine max* by indirect ELISA and their cellular location in root tissues. *Folia Microbiol.***42**: 607-612.
- Chakraborty, B.N. and Sharma, M. (2007) Serological detection and immunogold localization of cross-reactive antigens shared by *Camellia sinensis* and *Exobasidium vexans*. *Journal of Applied Microbiology*. doi:10.1111/j.1365-2672.2007.03459.x

- Charudattan, R. (1970).** Studies on strain of *Fusarium vasinfectum* Atk. II. In vitro production of toxin and enzymes and immunology. *Phytopath. Z. Physiol. Mol. Plant Pathol.* **58**: 31-41.
- Charudattan, R. and DeVay, J.E. (1972).** Common antigen among varieties *Gossypium hirsutum* and isolates of *Fusarium* and *Verticillium* species. *Phytopathology* **62**: 230-234.
- Charudattan, R. and DeVay, J.E.(1981).** Purification and partial characterization of an antigen from *Fusarium oxysporum* f. *vasinfectum* that cross-reacts with antiserum to cotton (*Gossypium hirsutum*) root antigens. *Physiol. Plant Pathol.* **18** P: 289-295.
- Clark, M.F. (1981).** Immunosorbant assays in plant pathology Annu. Rev. *Phytopathol.* **19**: 83-106.
- Clark, M. F., and Adams, A.N. (1997).** Characteristics of the microplate method of enzyme linked immunosorbant assay (ELISA). *J. Gen. Virology* **33**: 165-167.
- Clausen, J. (1988).** Laboratory techniques in Biochemistry and Molecular Biology. **Vol. I part III.** (Eds. By. R. H. Burdon and P. H. Van. Knippenberg). pp. 64-65.
- Coleman, M. J., Mainzer, J. and Dickerson, A. G. (1992).** Characterization of a fungal glycoprotein that elicits a defense response in French bean. *Physiol. Mol. Plant. Pathol.* **40**: 333-351.
- Copper, C. Issac, S., Jones, M. G. Crowther, T., Smith, B. M. and Collin, H. A. (2004).** Morphological and biochemical responses of carrots to *Pythium violae*, causative agent of cavity spot. *Physiol. Mol. Plant. Pathol.* **64**: 27-35.
- Cristinzio, G., Marziano, F. and Giannattasio, M. (1988).** Agglutination response of the conidia of eight *Fusarium* species to lectins having different sugar binding specifications. *Plant Pathol.* **37**: 120-124.

- Daayf, F., Schmidt, A. and Belanger, P. R (1995). The effect of plant extracts of *Reynoutria sachalinensis* on powdery mildew development and leaf physiology of long English cucumber. *Plant Dis.* 79: 577-580.
- Daayf, F., Schmidt, A. and Belanger, P. R (1997). Evidence of phytoalexin in cucumber leaves infected with powdery mildew following treatment with leaf extracts of *Reynoutria sachalinensis*. *Plant Physiol.* 11'3: 719-727.
- Darshika, P. and Daniel, M. (1992). Changes in the chemical content of *Adhatoda* and *Trianthema* due to fungal disease. *Indian J. Pharm.* 54:73-75.
- Day, A.W. Gardiner, R.B.; Svirrcev, A.M. and McKeen, W.E. (1986). Detection of fungal fimbriae by proten-a-gold immunocytochemical labeling in host plants infected with *Ustilago heufleri* or *Peronospora hyoscyami f. sp tabacina* *Can.J.Microbiol* 32 : 77-584.
- De Vay, J. E., Wakeman, J. J., Kavanagh, J. A. and Charudattan, R. (1981). The tissue and cellular location of a major cross-reactive antigen shared by cotton and soil-borne fungal parasites. *Physiol. Plant Pathol.*, 18: 59-66.
- DeVay, J.E., Wilson, Wakeman, R.J. and Kavanagh, J. A. (1981 b). Occurrence of common antigenic determinants between potatoes and *Phytophthora infestans* in relation to host parasite compatibility. In : Proc. **Inter Symp. On Phytophthora : Its Biology, Ecology and Pathology**, D.C. Erwin, S. Bartniti- Garcia, P. Tasao. Ed. 43, Dept, of Plant Pathology. University of California, Riverside.
- Dewey, F.M., Barret, D.K., Vose, I.R. and Lamb, C.J. (1984). Immunofluorescence microscopy for the detection and identification of propagules of *Phaelous schweinitzii* in infested soil. *Phytopathology* 74: 291-296.
- Dhingra, O.D. and Sinclair J.B. (1985). **Basic Plant Pathology Methods**. CRC Press, Inc.Boca Raton, Florida.
- Duncan, J.M. (1980). A technique for detecting red stele (*Phytophthora frageriae*) infection of strawberry stocks before planting. *Plant disease.*, 64: 1023-1025.

- D'Silva, I. and Heath, M. C. (1997).** Purification and characterization of two novel hypersensitive response including specific elicitors produced by the cowpea rust fungus. *J. Biol. Chem.* **272**: 3924-3927.
- Etenbarian, H. R. (1989).** Studies on quantitative changes in phenolic compound on barley varieties during the development of *Puccinia hordei* and the relationship between these substances and brown rust resistance in barley. *Physiol. Mol. Plant Pathol.* **33**: 88-89.
- Fuhrmann, B; Roquebert, M.F., Hoegaerden and Strosberg (1989).** Immunological differentiation of *Penicillium*. *Can. J. Microbiol.* **35**: 1043-1047.
- Gardiner, R.B., McKeen, W.E., Lawrence, T.M., Smith, R.J. and Day, A.W. (1989).** Inhibition of *Botrytis cinerea* spore germination by immunoglobulins. *Can. J. Bot.* **67**: 922-927.
- Gawande, S. J., Chimote, V. P., Shukla, A. and Gadewar, A. V. (2006).** Serological and molecular diagnosis of plant pathogens In: *Plant Protection in New Millennium* (Eds: A. V. Gadewar and B. P. Singh) **1**: 563-588.
- Gawande, V. L., Patil, J. V., Naik, R. M. and Kale, A. A. (2002).** Plant biochemical defence against powdery mildew (*Erysiphe polygoni* DC) disease in mungbean (*Vigna radiata* L. Wilezek). *J.Plant Biol.* (29)(3): 337-341.
- Gerik, J.S. and Huisman, O.C. (1988).** Study of field grown cotton-roots infected with *Verticillium dahliae* using an immunoenzymatic staining technique. *Phytopathology* **78**: 1174-1178.
- Ghosh, S. and Purkayastha, R.P. (1990).** Analysis of host parasite cross-reactive antigens in relation to *Myrothecium* infection of soybean. *Ind. J. Expt. Biol.* **28**: 1-5.
- Gupta, V., Kumar, A., Sing, A. and Gagr, G.K. (2000).** Immunodetection of teliospores of karnal bunt (*Telitia indica*) of wheat using fluorescent staining test. *Plants Cell Biotechno. Mol. Biol.* **1**: 81-86.



- Gwinn, K.D., Collins, S. and Reddick, B. B. (1991). Tissue Print- immunoblot: an accurate method of the detection of *Acremonium coenophialum* in tall fescue. *Phytopathology* 81: 747-748.
- Hansen, M.A. and Wick R.L. (1993). Plant disease diagnosis: Present status and future prospects. *Adv. Plant Pathol.* 10: 64-126.
- Harborne, J. B. (1999). The comparative biochemistry of phytoalexin induction in plants. *Biochem. Sys. Ecol.* 27: 335-367.
- Hardham, A.R. and Suzaki, E. (1989). Glycoconjugates on the surface of spores of the pathogenic fungus *Phytophthora cinnamoni* studied using fluorescence and electron microscopy and flow cytometry. *Can. J. Microbiol.* 36: 189-192.
- Harold, H., Burdsall, JR., Mark, B. and Mark, E.C. (1990). Serologically differentiation of three species of *Armillaria* and *Lentinula edodes* by enzyme linked immunosorbant assay using immunized chickens as a source of antibodies. *Mycologia*, 82(4) : 415-423.
- Harrison J.G., Barker, H., Lowe, R and Rees E. A. (1990). Estimation of amounts of *Phytophthora infestans* mycelium in leaf tissue by enzyme linked immunosorbent assay. *Plant Pathol.* 39: 274-277.
- Haslam, E. (1999). Che fars senza polifenoli. In: *Plant Polyphenols* (Eds.G. G. Gross, R.W. Hemingway and T.Y.Yoshida) Kluwer Academic Publishers New York. NY. PP-15-40.
- Holtz, B. A., Karu A.E., and Weinhold, A.R. (1994). Enzyme linked immunosorbent assay for detection of *Thielaviopsis basicola*. *Phytopathology* 84: 977-983.
- Horn, K L and Jagicza, A (1973). Fluorescent Antibody staining of *Fusarium culmorum*. *Acta Phytopathologica Aca. Sci. Hungaricae* 8: 357-363.
- Iannelli, D, Capparelli, R. Cristinzio, G. Marziano, F., Scala, F. and Noviello, C. (1982). Serological differentiation among formae speciales and physiological races of *Fusarium oxysporum*. *Mycologia* 74 : 313-319.

- Ishizaki, H., Nakamura, and Wheat, R.W. (1981).** Serological cross reactivity between *Sporothrix schenckii* and various unrelated fungi. *Mycopathologia* **73**: 65-68.
- Jain, N.K. (1991).** Indian tea in Retrospect and prospect and the impact R.D. In: World International Symposium on Tea science, Japan, 45-57.
- Jones, C. J. and Hartley, S. E. (1999).** Protein competition model of phenolic allocation. *Oikos* **8**: 27-44.
- Joshi, U. N., Gupta, P. P., Gupta, V. and Kumar, S. (2004).** Biochemical factors in cluster bean that impart *Alternaria* blight resistance. *J. Mycol. Pl. Pathol.* **34**: 581-583.
- Kalim, S., Luthra, Y. P. and Gandi, S. K. (2000).** Influence of bavistin seed Treatment on morpho-physiological and biochemical parameters of cowpea roots susceptible to rhizoctonia species. *J. Mycol. Pl. Pathol.* **30**: 375-379.
- Kalyansundaram, R. and Charudattan, R. (1969).** Serological studies in the genus *Fusarium* : A comparison of strains. *Phytopath. Z.* **64**: 28-31.
- Kauss, H., Franke, R., Krause, K., Conrath, U., Jeblick, W., Grimming, B. and Matern, U. (1993).** Conditioning of parsley (*Petroselinum crispum*L.)suspension cells increase elicitor induced incorporation of cellwall phenolics. *Plant Physiol.* **102**: 259-446.
- Keen, N, T. Legrand, M. (1980).** Surface glycoproteins: evidence that they may function as the race specific phytoalexin elicitors of *Phytophthora megasperma* f. sp. *glycinea* *Physiol. Plant Pathol.* **17**: 175-192.
- Kemp, G., Boha, A.M., Klosppers, F. J. and Pretorius, Z.A. (1999).** Disease development and  $\beta$ -1, 3- glucanase expression following leaf rust infection in resistant and susceptible near-isogenic wheat seedlings. *Physiol. Mol. Plant Pathol.* **55** : (45-52).
- Kitagawa, T.; Sakamoto, Y.; Furumi, K and Ogura, H (1989).** Novel Enzyme immunoassay for specific detection of *Fusarium oxysporum* f.sp.

*cucumerinum* and for general detection of various *Fusarium* species. *Phytopathology* 79: 162-165.

**Kumar, G. R., Ganesh, R. and Pathak, V. N. (1990).** Some biochemical changes in the shoots of pearl millet infected downy mildew. *Adv. Plant. Sci.*, 5: 103-108.

**Kumar, A., Lakhchaura, B.D., Singh, A. and Garg, G.K., (2000).** Development of seed immunoblot binding assay for detection of Karnal bunt of wheat. *Proceedings-Indian Phytopathological Society. Vol. 1* : 424-425.

**Kumar, S., Mandakini, N. and Shridhar, R. (1997).** Cell wall bound phenolics resistance of rice plants to blast. *J. Mycol. Pl. Pathol.* 27. (1): 1-5.

**Kumar, S., Singh, H. N. and Shukla, R. S. (1995).** Host-pathogen interactions influencing the susceptibility of *Drechslera rostrata* causing leaf blight disease of *Costus speciosus*. In: *Global Conf. Adv. Res. on Plant Dis. Management.*, p-43. Udaipur. India.

**Laemmli, U.K. (1970).** Cleavage of structural proteins during assembly of the head of the bacteriophage T4. *Nature* 227: 680-685.

**Lange, L.; Heide, M. and Olson, L.W. (1989).** Serological detection of *plasmodiophora brassica* by dot immunobinding and visualization of the serological reaction by Scanning electron microscop. *Phytopathology* 79: 1066-1071.

**Lange, L.; Heide, M. and Olson, L.w. (1989).** Serological detection of *plasmodiophora brassicae* by dot immunobinding and visualization of the serological reaction by Scanning electron microscop. *Phytopathology* 79: 1066-1071.

**Laxlt, A.M., Madrid, E.A., Beligni, M.V. and Lamattina, L. (1998).** Anti-idiotypic antibodies can mimic the ability of *Phytophthora infestans* to induce the accumulation of transcripts associated with defence responses in potato. *Physiological and Molecular Plant Pathology* 53 : 135-148.

**Linfield C.A. (1993).** A rapid serological test for detecting *Fusarium oxysporum f.sp.marcissi* in Narcissus. *Ann.Appl.Biol.* 123 : 685-693.

- Lowry, O.H.; Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with folin phenol reagent. *J. Biol. Chem.* **193**: 265-275.
- Luthra, Y. P. (1989). Total phenols and their oxidative enzymes in Sorghum leaves resistant and susceptible to *Ramulispora sorghicola*. *Phytopathology*. **20**: 105-150.
- Lyons, N.F. and White, J.G. (1992). Detection of *Pythium violae* and *Pythium sulcatum* in carrots with cavity spot using competition ELISA. *Ann. Appl. Biol.* **120**: 235-244.
- Lyons, P. C., Wood, K. V. and Nicholson, R. L. (1990). Caffeoyl ester accumulation in corn leaves inoculated with fungal pathogen. *Phytochemistry*, **29**: 97-101.
- MacDonald, J.D. and Duniway J.M. (1979). Use of fluorescent antibodies to study the survival of *Phytophthora megasperma* and *P. cinnamomi* zoospores in soil. *Phytopathology* **69**: 436-441.
- Mackie, A. J., Roberts A.M., Green, J.R. and Callow, (1993). Glycoproteins recognized by monoclonal antibodies UB7, UB8 and UB10 are expressed early in the development of pea powdery mildew haustoria. *Physiol. Mol. Plant Pathol.* **43**: 135-146.
- Mahadevan, A. and Sridhar, R. (1982). *Methods in Physiological Plant Pathology* 2<sup>nd</sup> Edn. Sivakami Publication, Madras.
- Mahadevan, A. and Ulaganathan, K. (1991). *Techniques Molecular in Plant Pathology* Publication. Madras pp 219.
- Mansfield, J. W. (1990). Recognition and response in plant-fungus interactions. In: *Recognition and Response in Plant Virus Interactions*. (Ed.R.S.S. Fraser), Springer-Verlag, Berlin. pp 31-52.
- Marshall, M. R. and Partridge J. E. (1981). Immunochemical identification of *Fusarium moniliforme* ribosomes from diseased corn (*Zea mays* L.) stalk tissue. *Physiol.Plant Pathol.* **19**:227-288.

- Masago, H., Yoshikawa, M., Matsumoto, T. Katsumoto M., Ogata, H and Kishibe, Y. (1989). Differentiation and identification of *Phytophthora* species by Two-Dimensional Electrophoresis. *Ann. Phytopath. Spc. Japan.* **55**: 336-343.
- Matern, U. and Kneusel, R. E. (1988). Phenolic compounds in plant disease resistance. *Phytoparasitica*, **16**: 153-170.
- Merz, W.G., Burrell, R.G. and Gallegly, M.E. (1968). A surgical comparison of six heterothallic species of *Phytophthora*. *Phytopathology* **59**: 367-370.
- Miler, S.A., Madden, L.V. and Schmitthenner, A.F. (1997). Distribution of *Phytophthora* spp. In field soils determination by immunoassay. *Phytopathology* **87** (1): 1001-107.
- Mohan, S.B. (1988). Evaluation of antisera, raised against *Phytophthora fragariae* for detecting the red core disease of strawberries by Enzyme linked immunosorbent assay (ELISA). *Plant Pathology* **37**: 206-216.
- Mohr, P. G. and Cahill, D. M. (2000). Relative roles of glyceolin, lignin and the hypersensitive response and the influence of ABA in compatible and incompatible interactions of soybeans with *Phytophthora sojae*.
- Mukhopadhyay, A.N. (1994). Biocontrol of soil borne plant pathogens current status, future prospects and potential limitations. *Ind. phyto pathol* **47**(2): 119-126.
- Mukhopadhyay, A.N. Shrestha, S.M. and Mukherjee, P.K. (1992). Biological seed treatment for control of soil borne plant pathogens. *FAO Pl Prot. Bull.* **40**: 21-30.
- Nachmlas, A., Buchner, and Krikn, J. (1982). Comparison of protein lipopolysaccharide complexes produced by pathogenic and non pathogenic strains of *Verticillium dahliae* Kleb. From potato. *Physiol. Plant Pathol*, **20**: 213-221.
- Nemec, S., Jabaji- Hare, S. and Charest, P.M. (1991). ELISA and immunocytochemical detection of *Fusarium solani* produced naphthazarin toxins in citrus trees in Florida. *Phytopathology* **81**: 1497-1503.

- Nicholson, R. L., Hipskid, J. and Hanan, R. M. (1990). Protection against phenol toxicity by the spore mucilage of *Colletotrichum graminicola*, an aid to secondary spread. *Physiol. Mol. Plant. Pathol.* **35**: 243-252.
- Niemann, G. J., Van der Kerk, A., Niessen, M. A. and Versluis, K. (1991). Free and cell-wall bound phenolics and other constituents from healthy and fungus infected carnation (*Dianthus caryophyllus* L.) stems . *Physiol. Mol. Plant Pathol.* **38**: 417-432.
- Nozzolillo, C., Isabelle, P., Das, G. (1990). Seasonal changes in the phenolics constituents of jack pine seedlings (*Pinus banksiana*) in relation to the purpling phenomenon. *Can. J. Bot.* **68**: 2010-2017.
- Oke, O. A. (1988). Changes in chemical constituents of tobacco leaves infected with *Cassicola corynespora* and *Colletotrichum nicotianae*. *J. Phytopathol.* **122**: 181- 185.
- Ouchterlony, O. (1967). In: Weir (ed.) Handbook of experimental immunology, Weir (ed.) (Blackwell Inc. London ) p.655.
- Purkayastha, R. P. (1994). Phyto-immunology an emerging discipline of plant science..*Everyman's Science* **29**(2): 41-44.
- Purkayastha, R. P. and Ghosh, A. (1987). Immunological studies on root rot of groundnut (*Arachis hypogea* L. ). *Can. J. Microbiol.* **33** : 647-651.
- Purkayastha, R.P and Banerjee R. (1989). Immunoserological studies on cloxacillin induced resistance of soybean against anthracnose. *Plant Dis.Prot.* **97**: 350-259.
- Purkayastha, R.P. and Ghosal, A. (1985). Analysis of cross-reactive antigens of *Acrocyldrium oryzae* and rice in relation to sheath rot disease. *Physiol. Plant Pathol.* **27**: 245-252.
- Purkayastha, R.P. and Pradhan, S. (1994). Immunological approach to study the etiology of *Sclerotium* rot disease of Groundnut. Proc. *Indian Nat. Sci. Acad.* **B-60**: 157-165.

- Rafin, C., Nodet, P. and Tirilly, Y. (1994).** Immunoenzymatic staining procedure for *Pythium* species with filamentous non-inflated sporangia in soil less culture. *Mycol.Res.* **98**: 535-541.
- Ranganathan, V. and Natesan, S. (1987).** Manuring of Tea Revised Recommendation, Hand Book of Tea Culture, Section 11, p.1-27.
- Ransom, R. F., Hipskind, J., Leite, B., Nicholson, R. L. and Dunkle, L. D. (1992).** Effects of elicitors from *Colletotrichum graminicola* on the response of soeghum to *Periconia circinata* and its pathotoxin. *Physol. Mol. Plant Pathol.*, **41**: 75-84.
- Rataj-Guranowska, M., and Wilko, B. (1991).** Comparison of *Fusarium oxysporum* and *Fusarium oxysporum* var. *redolens* by analyzing the isozyme and serological patterns. *Phytopathology* **132**: 287-293.
- Reddy, M.K. and Ananthanarayanan, T.V. (1984).** Detection of *Ganoderma lucidum* in betelnut by the fluorescent antibody technique. *Trans. Br. Mycol. Soc.* **82**: 559-561.
- Ricci, P., Trentin, F., Bonnet, P., Venard, P., Mouton-Perronnet and Bruneteau, M. (1992).** Differential production of parasiticein, an elicitor of necrosis and resistance in tobacco, by isolates of *Phytophthora parasitica*. *Plant Pathol.* **41**: 298-307.
- Roberts, A.M., Mackie, A.J., Hathaway, V., Callow, J.A. and Green, J.R. (1993).** Molecular differentiation in the extra haustorial membrane of pea powdery mildew haustoria at early and latestages of development. *Physiological and Molecular Plant Pathol.* **43**: 147-160.
- Sahabuddin, M. and Anwar, M. N. (1992).** Metabolic changes in the jute plant due to fungal infection. *Bangladesh. J. Bot.* **21**: 205-212.
- Saxena, A. K., Sharma, A. K., Nityagopal, N. and Bakshi, A. K. (1986).** Changes in total phenolics and sugars in relation to *Puccinia recondata* (brownrust) of some wheat varieties. *J. Res. Punjab. Agric. Univ.* **24**: 359-62.

- Shane, W.W. (1991). Prospects for Early Detection of *Pythium* blight epidemics on turfgrass by Antibody- Aided Monitoring. *Plant Dis.* **75** : 921-925.
- Sharma, I. M. (2003). Relationship between phenyl propanoid metabolism and resistance to *Pythium ultimum* Trow in apple Rootstocks. *J. Mycol. Pl. Pathol.* **33**: 114-118.
- Sharma, M. and Chakraborty, B. N. (2005). Hexaconazole and calixin mediated defense strategies of tea plants against *Exobasidium vexans* Masee. *Journal of Mycology and Plant Pathology*, **35(3)**: 417-431.
- Sharma, M. and Sharma, S. K. (2001). Biocontrol of *Dematophora necatri* causing white rot of apple. *Plant Dis. Res.*, **16**: 40-45.
- Sharma, M. and Chakraborty, B. N. (2005). Hexaconazole and calixin mediated defense strategies of tea plants against *Exobasidium vexans*. Masee. *J. Mycol. Pl. Pathol.* **34(3)**: 742-760.
- Sinha, A.K. (1995). Possible role of phytoalexin inducer chemicals in plant disease control. In: *Handbook of Phytoalexin Metablism and Action* (Eds. M Daniel and R.P. Purkayastha), Marcel Dekker, Inc. pp. 555-591.
- Srivastava, A.K. and Arora, D. K. (1997). Evaluation of a polyclonal antibody immunoassay for detection, quantification of *Macrophomina phaseolina*. *Plant Pathol.* (Oxford) **46**: 735-794.
- Sukhlal, R. and Purohit, S. D. (2003). Accumulation of phenolics and changes in activity of oxidative enzymes in maize-infected with *Helminthosporium* species. *J. Mycol. Pl. Pathol.* **33**: 236-239.
- Takenaka, S. (1992). Use of immunological methods with antiribosome serums to detect snow mold fungi in wheat plants. *Phytopathology* **82**: 896-901.
- Tore, G. D. and Tosi, L. (1989). Alteration and composition of phenolics compounds and amino acids in tobacco seedlings infected with *Thielaviopsis basicola*. *Ann. Fac. Agrar. Univ. Stud. Perugia.*, **43**: 279- 288.
- Townsend, R & Archer, D.B. (1983). A fibril protein antigen specific to Spiroplasma *J. Gen. Microbiol.* **129**: 199-206.



- Van Etten, H. D., Mathenes, D.E. and Swith, D. A. (1982).** Metabolism of phytoalexins. **In: *Phytoalexins*** (Eds. J.A. Bailey and J.W. Mansfield) pp. 181-217, Blackie, Glasgow.
- Viswanathan, R., Padnanaban, P., Mohanraj, D., and Jothi, R. (2000).** Indirect ELISA technique for the detection of the red rot pathogen in sugarcane (*Saccharum* spp. *hybrid*) and resistance screening. ***Ind. J. Agric. Sci.*, 70:** 308-31.
- Wakeham, A. J. and White, J.G. (1996).** Serological detection in soil of *Plasmodiophora brassica* resting spores. ***Physiol. Mol. Plant Pathol.* 48:** 289-303.
- Walez, I., Pacsa, A.S.I Emody and Szabo, L. Gy, (1985).** Detection of *Sclerotinia sclerotiorum* in sunflower by Enzyme-Linked Immunosorbant assay (ELISA). ***Trans. Br. Mycol. Soc.* 85:** 485-488.
- Walsh, J.A., Merz, U. and Harrison, J.G. (1996).** Serological detection of spore balls of *Spongospora subterranean* and quantification in soil. ***Plant Pathology* 45:** 884-895.
- Werder, J. and Kern, H. (1985).** Resistance of maize to *Helminthosporium carbonum* changes in host phenolics and their antifungal activity. ***Plant Dis. Protect.* 92:** 477-484.
- Werres, S. and Steffens, C. (1994).** Immunological techniques used with fungal plant pathogens: aspects of antigens, and antibodies and assays for diagnosis. ***Ann. Appl. Biol.* 125:** 615-617.
- Werres, S. and Steffens, C. (1994).** Immunological techniques used with fungal plant pathogens: aspects of antigens, and antibodies and assays for diagnosis. ***Ann. Appl. Biol.* 125:** 615-617.
- Wycoff, K.L and Ayers A.R. (1991).** Localization of carbohydrate antigens in the walls of *Phytophthora megasperma* f. sp. *Glycinea* by monoclonal antibodies. ***Physiol. Mol Plant Pathol.* 39:** 95-109.

- Yoshikawa, M. and Sugimoto, K. (1993). A specific binding site on soybean membranes for a phytoalexin elicitor released from fungal cell walls by Beta-1-3 endoglucanase. *Plant Cell Physiol.* **34**: 1229-1247.
- Young, C.S. and Andrews, J.H. (1990). Inhibition of pseudothecial development of *Venturia inaequalis* by the Basidiomycete *Athelia bombacina* in apple leaf litter. *Phytopathology* **80**: 536-542.
- Young, C.S. and Andrews, J.H. (1990). Inhibition of pseudothecial development of *Venturia inaequalis* by the Basidiomycete *Athelia bombacina* in apple leaf litter. *Phytopathology* **80**: 536-542.
- Yuen, G.Y., Kim, K.N. and Horst, G.L. (1994). Use of ELISA and isolation for determining the distribution of *Rhizoctonia solani* and other *Rhizoconia* spp. In asymptomatic creeping bentgrass. *Crop Protection* **13(4)**: 296-300.
- Yuen, G.Y., Xia, J.Q. and Satula, C.L. (1988). A sensitive ELISA for *Pythium ultimum* using polyclonal and species-specific monoclonal antibodies. *Plant Dis.* **89**: 1029-1032.

