

Preface

The nitrono moiety can be regarded as a 3 centered dipolar 4π system, which enables 1,3-dipolar cycloaddition reactions with different dipolarophilic reagent to occur. 1,3-dipolar cycloadditions are found to be susceptible to both electronic and steric influences. 1,3-dipolar cycloaddition reaction between a nitrono and an olefinic dipolarophiles is an efficient method for the synthesis of isoxazolidine systems.

The question of reactivity and substituent effects in 1,3-dipolar cycloaddition reaction has been rationalized successfully using frontier molecular orbital theory which provides relative interaction energies of frontier orbitals between 1,3-dipole and dipolarophiles. The electron attracting or electron releasing moiety influences the atomic orbital co-efficiency and have a significant influence on the regioselectivity of the reaction.

Furthermore, the cycloadducts have found numerous applications in synthesis through reductive cleavage of the N-O bond to give γ -amino alcohols. Asymmetric induction in nitrono olefin cycloaddition has been achieved through the incorporation of chirality in both the dipole and the dipolarophiles. Recently, significant advances have been made by the use of water as the solvent to influence the rate, regioselectivity and stereoselectivity of cycloaddition reactions. The present work entitled "*Synthesis and 1,3 dipolar cycloaddition reactions of N-Phenyl- α -Chloro Nitrono*" reports newly discovered α -chloro nitrono from chlorohydrin and its cycloaddition reactions with different olefins and alkynes leading to the formation of regio and stereoselective products. An important application of the nitrones and the cycloadducts are aldehyde synthesis and antimicrobial activities. An efficient methodology for the synthesis of aldehydes from alkyl halides using α -chloro nitrones as an oxidizing reagent with an excellent yield has also been discussed along with novel α -N-methyl/phenyl furan derivatives (obtained as side product during aldehyde & ketone synthesis) as dipolarophile in the regioselective synthesis of 5-spiro isoxazolidines using α -chloro nitrones at RT with an excellent yield.

The following chapters fulfill these ideas:-

Chapter I This chapter deals with the general theoretical approach of different 1,3-dipoles and their stabilities and general nature of intra and intermolecular 1,3-dipolar cycloaddition reactions of nitrones. Special emphasis has been given on HOMO-LUMO approach in this regard. Attempts have been made in this chapter to cover a complete review of the literature and latest developments up to January 2010 in a rather comprehensive manner.

Chapter II It deals with the experimental section. In this section, experimental procedures are discussed along with the method of formation of different nitrones, cycloaddition reaction with different olefins and alkynes leading to the formation of cycloadducts. The cycloadducts are explained along with their reaction conditions in different solvents.

Chapter III This chapter deals with results and discussion and achievements of the work done. Spectral interpretation *viz.* ^1H NMR, ^{13}C NMR, MS, IR, HRMS and elemental analysis have been discussed in detail.

Chapter IV This chapter is focused on the future perspective of the work done and is referred to as *scope and objectives* of the present work.