

List of Tables

Table	Title	Page No.
1	Site characteristics of selected populations of <i>Swertia chirayita</i> in Sikkim	71
2	Phytosociological attributes of <i>Swertia chirayita</i> and its associates in Sikkim	73
3	Diversity and dominance pattern of <i>Swertia chirayita</i> pockets in Sikkim	77
4	Types and intensity of existing disturbances in different populations of <i>Swertia chirayita</i> in Sikkim Himalaya	78
5	Morphological differences amongst populations of <i>Swertia chirayita</i> (rosette) in Sikkim	97
6	Morphological differences amongst populations of <i>S. chirayita</i> (reproductive) in Sikkim	98
7	Pearson's correlation coefficient (r) and significance of correlation (P) between the means of observed parameters of <i>S. chirayita</i> with altitude and humus depth in Sikkim	99
8	Result of MANOVA showing morphological variation amongst populations of <i>S. chirayita</i> in Sikkim	100
9	Pearson's correlation matrix amongst populations of <i>S. chirayita</i> (rosette) in Sikkim	101
10	Pearson's correlation matrix amongst populations of <i>S. chirayita</i> (reproductive) in Sikkim	102
11	Summary of Canonical Discriminant Function of the growth parameters among populations of <i>S. chirayita</i> (rosette) in Sikkim	103
12	Summary of Canonical Discriminant Function of the growth parameters among populations of <i>S. chirayita</i> (reproductive) in Sikkim	104
13	Results of two way ANOVA showing effect of temperature and microhabitats and their interaction on various germination parameters in <i>Swertia chirayita</i>	119
14	Seed characteristics of different populations of <i>Swertia chirayita</i> in Sikkim	128
15	Result of Multivariate ANOVA depicting the effect of population, storage period and their interaction on seed germination and mean germination time in <i>Swertia chirayita</i>	128
16	Site characteristics of the populations of <i>Swertia chirayita</i> in Sikkim	140
17	Result of MANOVA characteristics by populations, treatments and their interaction on <i>Swertia chirayita</i>	141
18	Result of MANOVA characteristics by populations, treatments and their interaction on <i>Swertia chirayita</i>	142
19	Correlation matrix (Pearson Product-Moment Correlation Coefficient) between different growth parameters in <i>Swertia chirayita</i> after 24 weeks of sowing at <i>ca.</i> 2000m asl	143
20	Seed characteristics of different plant population of <i>Swertia</i>	158

	<i>chirayita</i> collected in Sikkim	
21	Result of Three way ANOVA showing the effect of populations, storage conditions and storage periods and their interaction on seed germination and mean germination time in <i>Swertia chirayita</i>	158
22	Summary of one way MANOVA depicting the effect of different growing conditions on the growth and biomass of <i>Swertia chirayita</i>	176
23	Pearson's correlation coefficient amongst different growth parameters in <i>Swertia chirayita</i>	176
24	Seed characteristics of <i>Swertia chirayita</i> grown under different conditions at <i>ca.</i> 2000m asl	194
25	Effect of different chemical treatments on seed germination in <i>Swertia Chirayita</i> , sourced from six <i>ex-situ</i> conditions	195
26	Effect of different chemical treatments on Mean Germination Time in <i>Swertia Chirayita</i> , sourced from six <i>ex-situ</i> conditions	196
27	Results of MANOVA showing effect of different seed sources, treatments and their interaction on seedling emergence variables in <i>Swertia chirayita</i>	206
28	Index of emergence rate in <i>Swertia chirayita</i> , using a modified Timson Index of germination velocity	206
29	Results of MANOVA showing effect of seed sources, treatments and their interaction on different growth parameters in <i>Swertia chirayita</i>	207
30	Correlation matrix (Pearson Product-Moment Correlation Coefficient) between different growth parameters in <i>Swertia chirayita</i>	208
31	Total investment (in Indian Rupee, Rs*) per hectare on the basis of present market rate for the cultivation of <i>Swertia chirayita</i> at Pangthang (<i>ca.</i> 2000m asl), Sikkim (India)	226
32	Result of ANOVA characterized by the effect of growing conditions in <i>Swertia chirayita</i> cultivated at Pangthang (<i>ca.</i> 2000m asl), Sikkim (India)	226
33	Total production (per hectare) at the end of third year of cultivation of <i>Swertia chirayita</i> on the basis of projected and actual plant survival percent, at Pangthang (<i>ca.</i> 2000m asl), Sikkim (India)	227
34	Economic viability (per hectare) at the end of third year of cultivation of <i>Swertia chirayita</i> on the basis of projected and actual plant survival percent, at Pangthang (<i>ca.</i> 2000m asl), Sikkim (India)	228