

LIST OF TABLES

<u>CHAPTERS</u>	<u>TABLE CAPTIONS</u>	<u>PAGE NO.</u>
Chapter-II	<i>Table II.1. Comparison between organic solvents and ionic liquids.</i>	64
Chapter-IV	<i>Table IV.1: Values of Density (ρ), Viscosity (η) and Relative Permittivity (ϵ_r) of ACN, THF, 1,3-DO and their binary mixtures where w_1, w_2 and w_3 are the mass fraction of ACN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	252
	<i>Table IV.2: Molar conductance (Λ) and the corresponding concentration (c) of Bu_4PBF_4 in ACN, THF, 1,3-DO and their binary mixtures where w_1, w_2 and w_3 are the mass fraction of ACN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	253
	<i>Table IV.3: Limiting molar conductivity (Λ_0), the association constant (K_A), the distance of closest approach of ions (R), Standard Deviations δ of Experimental Λ from Equation (IV.1), Walden Product ($\Lambda_0\eta$) and free energy change (ΔG°) of Bu_4PBF_4 in ACN and its binary mixtures with THF, 1,3-DO, where w_1 are the mass fraction of ACN at $T = 298.15$ K</i>	255
	<i>Table IV.4: Limiting Ionic Conductance (λ_0^\pm), Ionic Walden Product ($\lambda_0^\pm\eta$), Stokes' Radii (r_s) and Crystallographic Radii (r_c) of Bu_4PBF_4 in ACN and its binary mixtures with THF, 1,3-DO, where w_1 are the mass fraction of ACN at $T = 298.15$ K</i>	256
	<i>Table IV.5: The calculated limiting molar conductance of ion-pair (Λ_0), limiting molar conductance of triple-ion (Λ_0^T), slope and intercept of Equation (IV.15) of Bu_4PBF_4 in THF, 1,3-DO and their binary mixture, where w_2 and w_3 are mass fraction of THF and 1,3-DO at $T = 298.15$ K</i>	257

	<i>Table IV.6: Salt concentration at the minimum conductivity (c_{min}) along with the ion-pair formation constant (K_p), triple-ion formation constant (K_T) of Bu_4PBF_4 in THF, 1,3-DO and their binary mixture, where w_2 and w_3 are mass fraction of THF and 1,3-DO at $T = 298.15$ K</i>	257
	<i>Table IV.7: Salt concentration (c_{min}) at the minimum conductivity (Λ_{min}), the ion-pair fraction (α), triple-ion fraction (α_T), ion-pair concentration (C_p) and triple-ion concentration (C_T) of Bu_4PBF_4 in THF, 1,3-DO and their binary mixture, where w_2 and w_3 are mass fraction of THF and 1,3-DO at $T = 298.15$ K</i>	258
Chapter-V	<i>Table V.1: Values of density (ρ), viscosity (η) and relative permittivity (ϵ_r) of studied solvents at $T = 298.15$ K</i>	295
	<i>Table V.2: Molar conductance (Λ) and the corresponding concentration (c) of the studied ILs in different solvents at $T = 298.15$ K</i>	296
	<i>Table V.3: Limiting molar conductivity (Λ_0), the association constant (K_A), the distance of closest approach of ions (R), standard deviations δ of experimental Λ, Walden product ($\Lambda_0\eta$) and Gibb's energy change (ΔG°) of ILs in different studied solvents at $T = 298.15$ K</i>	297
	<i>Table V.4: Limiting ionic conductance (λ_o^\pm), ionic Walden product ($\lambda_o^\pm\eta$), Stokes' radii (r_s) and crystallographic radii (r_c) of imidazolium based ionic liquids in different studied solvents at $T=298.15$ K</i>	298
	<i>Table V.5: A-coefficient and solvation number (n_s) of ionic liquids in different studied solvents at 298.15K</i>	299
	<i>Table V.6: Diffusion coefficient (D_i) and ionic mobility (i_i) of ionic liquids in different studied solvents at 298.15K</i>	299
	<i>Table V.7: The calculated limiting molar conductance of ion-pair (Λ_0), limiting molar conductance of triple-ion (Λ_0^T), slope and intercept of equation (V.17) of</i>	300

	<i>ionic liquids in methylamine solution (CH_3NH_2) at $T = 298.15 \text{ K}$</i>	
	<i>Table V.8: Salt concentration at the minimum conductivity (c_{\min}) along with the ion-pair formation constant (K_p), triple-ion formation constant (K_T) for ionic liquids in methylamine solution (CH_3NH_2) at $T = 298.15 \text{ K}$</i>	300
	<i>Table V.9: Salt concentration (c_{\min}) at the minimum conductivity (Λ_{\min}), the ion-pair fraction (α), triple-ion fraction (α_T), ion-pair concentration (C_p) and triple-ion concentration (C_T) of ionic liquids in methylamine solution (CH_3NH_2) at $T = 298.15 \text{ K}$</i>	300
	<i>Table V.10: Interionic distance parameter for ion-pair (a_p) and for triple-ion (a_T) in methylamine solution (CH_3NH_2) at $T = 298.15 \text{ K}$</i>	301
	<i>Table V.11: Stretching frequencies of the functional groups present in the pure solvent and change of frequency after addition of ILs in the solvents.</i>	301
Chapter VI	<i>Table VI.1: Molarity (c), density (ρ), viscosity (η), apparent molar volume (ϕ_v), and $(\eta_r - 1)/c^{1/2}$ for Citric Acid in different aqueous Cetrimonium Bromide solution at different temperatures</i>	321
	<i>Table VI.2: Limiting partial molar volume (ϕ_v^0), adjustable parameters A_v and B_v, and values of viscosity A and B coefficients for Citric Acid in different aqueous Cetrimonium Bromide solutions with standard deviations σ and limiting partial molar volume of (water + Cetrimonium Bromide) at different temperatures.</i>	324
	<i>Table VI.3: Values of various coefficients of Eq. VI.4 for Citric Acid in different aqueous Cetrimonium Bromide solutions</i>	325
	<i>Table VI.4: Values of $(\bar{V}_2^0 - \bar{V}_1^0)$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for Citric Acid in different aqueous Cetrimonium Bromide solutions at different temperatures</i>	326
Chapter VII	<i>Table VII.1: Sample description</i>	344

	<i>Table VII.2: Values of Density (ρ), Viscosity (η) and Ultrasonic speed (u) of AN, THF, 1,3-DO and their binary mixtures where w_1, w_2 and w_3 are the mass fraction of AN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	345
	<i>Table VII.3: Experimental values of Densities (ρ), Viscosities (η), and Ultrasonic speeds (u) of $[Bu_4PBF_4]$ in AN, THF, 1,3-DO and their binaries where w_1, w_2 and w_3 are the mass fraction of AN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	346
	<i>Table VII.4: Molality (m), apparent molar volume (ϕ_V), $(\eta/\eta_0-1)/m^2$, adiabatic compressibility (β_S) and apparent molal adiabatic compressibility (ϕ_K) with their standard deviation (σ) of $[Bu_4PBF_4]$ in AN, THF, 1,3-DO and their binaries where w_1, w_2 and w_3 are the mass fraction of AN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	349
	<i>Table VII.5: Limiting apparent molar volumes (ϕ_V^0), experimental slopes (S_V^*), A-, B-coefficients, limiting partial adiabatic compressibility (ϕ_K^0), and experimental slope (S_K^*) of $[Bu_4PBF_4]$ in AN, THF, 1,3-DO and their binaries where w_1, w_2 and w_3 are the mass fraction of AN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	352
	<i>Table VII.6: Limiting ionic apparent molar volumes (ϕ_{\pm}^0), ionic viscosity B-coefficients (B_{\pm}), of $[Bu_4P]^+$ and $[BF_4]^-$ in AN, THF, 1,3-DO and their binaries where w_1, w_2 and w_3 are the mass fraction of AN, THF and 1,3-DO respectively at $T = 298.15$ K</i>	353
Chapter VIII	<i>Table VIII.1: Densities (ρ), Viscosities (η), Refractive Index (n_D) and Speed of sound (u) of Methanoic acid at 298.15 K</i>	370

	<i>Table VIII.2: Experimental values of Densities (ρ), Viscosities (η), Refractive Index (n_D) and Speed of sound (u) of Glycine, Alanine, Valine and Leucine in Methanoic acid at 298.15 K</i>	370
	<i>Table VIII.3: Molality (m), apparent molar volume (ϕ_V), $(\eta/\eta_0-1)/m^2$, molar refraction (R), adiabatic compressibility (β_S) and apparent molal adiabatic compressibility (ϕ_K) of Glycine, Alanine, Valine and Leucine in Methanoic acid at 298.15 K</i>	371
	<i>Table VIII.4: Limiting apparent molar volumes (ϕ_V^0), experimental slopes (S_V^*), A, B coefficients, limiting partial adiabatic compressibility (ϕ_K^0), and experimental slope (S_K^*) of Glycine, Alanine, Valine and Leucine in Methanoic acid at 298.15 K</i>	373
Chapter IX	<i>Table IX.1: Values of density (ρ), viscosity (η) and ultrasonic speed (u) of aqueous β-CD in different mass fraction (w_1), at 293.15K to 308.15K</i>	402
	<i>Table IX.2: Experimental values of density (ρ), viscosity (η), and ultrasonic speed (u) of amino acids in different mass fraction of aqueous β-CD (w_1) at 293.15K to 308.15K respectively</i>	403
	<i>Table IX.3: Molality (m), apparent molar volume (ϕ_V), $(\eta_r-1)/\sqrt{m}$, and apparent molar adiabatic compressibility (ϕ_K) of amino acids in different mass fraction of aqueous β-CD (w_1) at 293.15K to 308.15K respectively</i>	408
	<i>Table IX.4: Limiting apparent molal volumes (ϕ_V^0), experimental slopes (S_V^*), viscosity A, B-coefficients, limiting partial molal adiabatic compressibilities (ϕ_K^0), and experimental slopes (S_K^*) of amino acids in different mass fraction of aqueous β-CD (w_1) at 293.15K to 308.15K respectively</i>	416

	Table IX.5: Contributions of zwitter ionic group (NH_3^+ , COO^-), CH_2 group, and the other alkyl chains to the limiting apparent molar volume, ϕ_1^0 , for amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	419
	Table IX.6: Values of ϕ_V^0 (aqueous), $\Delta\phi_V^0$, ϕ_V^0 (elect), ϕ_K^0 (elect), and hydration number (n_H) for amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	420
	Table IX.7. Contributions of zwitter ionic group (NH_3^+ , COO^-), CH_2 group, and the other alkyl chains to the limiting apparent molar volume transfer $\Delta\phi_V^0$, in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	423
	Table IX.8: Values of empirical coefficients (a_0 , a_1 , and a_2) of Equation 14 for amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15K to 308.15K respectively	425
	Table IX.9. Limiting apparent molal expansibilities (ϕ_E^0) for amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15K to 308.15K respectively	425
	Table IX.10: Contributions of zwitter ionic group (NH_3^+ , COO^-), CH_2 group, and the other alkyl chains to the B-coefficient in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	426
	Table X.11: Values of $\frac{dB}{dT}$, A_1 , and A_2 coefficient for the amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	427
	Table IX.12: Values of ϕ_1^0 , ϕ_V^0 (aqueous), $\mu^{0\#}$, $T\Delta S^\#$, $\Delta H^\#$, for amino acids in different mass fraction of aqueous β -CD (w_1) at 293.15 to 308.15K respectively	428