

LIST OF FIGURES

- Fig. 1: Representative examples of different types of siderophores and their natural producers
- Fig. 2: Representative examples of mixed types of siderophores and their natural produce
- Fig. 3: Representative examples of different siderophores produced by members of the genus *Pseudomonas*
- Fig. 4: Map showing present study area in India
- Fig. 5: Map showing locations of collection of rhizosphere soil samples in Darjeeling and Jalpaiguri Districts of West Bengal
- Fig. 6: Isolation of bacteria from soil: serially diluted sample spread onto soil extract agar produced isolated bacterial colonies
- Fig. 7: Isolation of pure culture of bacteria by streaking on nutrient agar plates
- Fig. 8: Screening of bacteria in CAS agar plates to identify siderophore producing strains
- Fig. 9: Inhibitory effect of selected bacterial isolate on the growth of *Fusarium equiseti* evident by the dual culture test in PDA plates
- Fig. 10: Inhibitory effect of selected bacterial isolate on the growth of *Lasiodiplodia theobromae* evident by the dual culture test in PDA plates
- Fig. 11: Inhibitory effect of selected bacterial isolate on the growth of *Rhizoctonia solani* evident by the dual culture test in PDA plates
- Fig. 12: Inhibitory effect of selected bacterial isolate on the growth of *Collectotrichum gloeosporioides* evident by the dual culture test in PDA plates
- Fig. 13: Inhibitory effect of selected bacterial isolate on the growth of *Alternaria alternata* evident by the dual culture test in PDA plates
- Fig. 14: Inhibitory effect of selected bacterial isolate on the growth of *Fusarium solani* evident by the dual culture test in PDA plates
- Fig. 15: Inhibitory effect of selected bacterial isolate on the growth of *Fusarium graminearum* evident by the dual culture test in PDA plates
- Fig. 16: Biochemical characterization of selected bacterial isolates
- Fig. 17: Biochemical characterization of selected bacterial isolates
- Fig. 18: Biochemical characterization of selected bacterial isolates

- Fig. 19: Agarose gel electrophoresis of (a) extracted genomic DNA of different bacterial strains; (b) 1500 bp long PCR product; (c) 560 bp length PCR product; (d) Blue white screening of transformed *E. coli* JM109 cells
- Fig. 20: Phylogenetic tree generated by the neighbour-joining method
- Fig. 21: Production of extracellular lytic enzymes by selected antagonistic isolates
- Fig. 22: Inhibition of mycelial growth by crude acetone extract obtained from strain AS04 culture supernatant of (a) *F. solani* (b) *F. equiseti*
- Fig. 23: a) Amount of IAA produced by the selected antagonistic isolates; (b) Amount of siderophore produced by selected antagonistic isolates
- Fig. 24: Scanning Electron Microscopic observation of the interaction zone between fungus and antagonistic bacteria grown in PDA plates in dual cultures
- Fig. 25: characterization of siderophore: (a) Tetrazolium salt test; (b) Arnow's test
- Fig. 26: (a) Siderophore production in presence of various medium supplements; (b) The effects of incubation time on growth and siderophore production by *Pseudomonas putida* strain AS04.
- Fig. 27: Optimization of siderophore production by *Pseudomonas putida* strain AS04 at various incubation temperatures
- Fig. 28: Spectral scan of partially purified siderophore of *Pseudomonas putida* strain AS04 showing a peak at 430 nm
- Fig. 29: Tubes with increasing colour intensity, containing Amberlite XAD-2 fractions collected and tested for their siderophore content
- Fig. 30: Detection of hydroxamate siderophore as intense brown spots using Thin Layer Chromatography
- Fig. 31: Study of growth kinetics by turbidometry of *Pseudomonas putida* strains; (c) Study of *in vitro* antagonism by *P. putida* strains AS01 and AS04 against *F. solani* in PDB dual culture assay
- Fig. 32: Suppression of wilt disease in brinjal seedlings by the isolated *Pseudomonas putida* strains under sterile soil conditions
- Fig. 33: Disease Index (DI) of wilt in brinjal seedlings inoculated by *F. solani* and treated by isolated antagonistic *Pseudomonas putida* strains AS01 and AS04
- Fig. 34: Percent Efficacy of Disease Control (PEDC) of wilt in brinjal seedlings by the isolated antagonistic *Pseudomonas putida* strains AS01 and AS04 under sterile and unsterile conditions