

# **CHAPTER VIII**

## **ROLE OF GERMINATION INDUCED PEPTIDE POOL IN PLANT TISSUE CULTURE**

## 8.1 INTRODUCTION

The growth and development of higher plants can be considered to be characterized by the execution of cell division, expansion and differentiation along two axes: the apical-basal and radial patterning. In evolutionary terms, the apical-basal axis of development can be considered to have a strong selective advantage based upon plant competition for light, water and nutrients. Because plants are sessile organisms, their success in a particular environment will depend on their ability to integrate a complex range of external and internal information that may vary from time to time.

It is now fully recognized that plants exhibit greater morphological and developmental plasticity than animals. This conclusion has emerged as a result of integrating the data from molecular biological and genetic approaches with data gained from whole-plant physiological investigations (Trewavas and Knight, 1994). This organogenic plasticity in plants is coordinated by complex interplay between diverge signaling systems, leading to its immediate translation in the rate and plane of cell division, and cell expansion. The totipotency and plasticity of plant cells and tissues can be vividly exhibited when they are cultured *in vitro* with changing hormonal profiles. Basically plant tissue culture relies on the fact that more plant cells have the ability to regenerate a whole plant. Importantly totipotent cells must be able to differentiate not only into any cell in the organism, but also into extra-embryonic tissue associated with the organism. During development, the activities of plant hormones such as auxins, cytokinins, ethylenes, gibberellins and abscisic acid depend on cellular context and exhibit interactions that can be either synergistic or antagonistic. For example, auxin can suppress cytokinin biosynthesis (Nordstrom *et al.*, 2004), auxin and cytokinin can act synergistically to induce ethylene biosynthesis (Vogel *et al.*, 1998) and ethylene can modify auxin responses and meristem function (Souter *et al.*, 2004; Stepanova and Alonso, 2005).

To date, researchers have identified four major group of peptide-ligand-receptor pairs in plants (Ryan *et al.* 2002), which are involved in a variety of developmental processes, such as wound responses, cellular dedifferentiation, meristem organization and self-incompatibility. However, these must only be part of the story, because plant genome

sequencing has revealed many genes predicted to encode small peptide ligands and receptor-like kinases, whose function remain to be uncovered (Shiu and Bleecker, 2001). Furthermore, induced mutation in prohormone processing proteases has been shown to disrupt plant growth and development.

In this context peptide fractions isolated from different germination hours of *Vigna radiata* cv. Sonali B1 were analyzed for their bioactivity related to morphogenesis, cell division and dedifferentiation. Now, proper embryo development and germination are the learning phase for any plant to face the challenge of its adverse world. These two phenomena set the 'programming' of upcoming metabolic machinery, which will last till the death with minimum 'tuning'. 'Germination' is that crucial period where autotrophic plant also behaves heterotrophically, which is an excellent manifestation of its selfishness and dependence, and slowly empowers its machinery for their future 'autotrophic' nature. This 'switch over' is controlled by a number of intrinsic factors, triggered by internal and external signal perceptions. This report is an attempt towards the understanding of changing pattern of peptide bioactivity isolated from different germination periods (i.e. – hrs of germination) of sonamung [*Vigna radiata* (Wilczek) cv. *sonali* B1] and also the interaction of these peptides with different hormones controlling growth and development.

## **8.2 MATERIAL AND METHODS**

### **8.2.1 Plant Materials for Germination**

Seeds of dicotyledonous plant material, mung bean, *Vigna radiata* (B1 *sonali* var.) were collected from Central Pulses Research Institute (C.P.R.I.), Berhampur, West Bengal, India. Seeds were weighed (each set of 250g.) and allowed to culture in sterile petri-plates with absorbent cotton supplied with modified Hoagland solution with one-half strength major nutrients and full-strength micronutrients. The nutrient solution was aerated continuously and renewed weekly. Experiments were conducted inside a controlled environment growth chamber with following conditions: 14-h light period with a light intensity of  $350\mu\text{mol m}^{-2} \text{s}^{-1}$ , 25 °c / 20 °c day / night temperature, and 80% relative humidity. The plant materials were grown for different germination hours like 0h, 8h, 24h, 32h, 40h, 48h, 56h, 64h, 72h, 5days, 6days and 7days respectively and treated as

different sets for definite experimental objective. Likewise certified seeds of rice, wheat and chickpea were also processed aseptically and cultured as stated above.

### **8.2.2 Isolation and purification of low molecular weight peptides**

Aqueous extraction and purification of peptides from different plant materials up to the stage of ultrafiltration were performed in the same way as mentioned in Chapter III Section 3.2.2.

### **8.2.3 Paper chromatography**

100 µl (1 g fresh weight equivalent peptide) of each isolated peptide solution (peptide solutions of 8h, 24h, 32h, 40h, 48h, 56h, 64h, 72h, 5days and 6 days of germination) was loaded onto Whatman No-1 chromatography paper (size-46 cm x 57 cm, thickness-0.16 mm), and separated by descending chromatography with Isopropanol: Ammonia: Water(10:1:1 v/v).After removal of solvents the paper was sprayed with freshly prepared Ninhydrin location reagent (0.2g ninhydrin was dissolved in 100 ml methanol) (Plummer, 2003) on paper. The paper was placed in oven at 70°C for a very short period. When the bluish-purple spots were developed then it was taken out of the oven and documentation was done.

### **8.2.4 Study of mitotic index and cell division**

For the present work, root apical meristems of *Allium cepa* L. ( $2n = 16$ ) were used. Approximately equal sized and healthy looking bulbs of *A. cepa* were collected from local market; surface sterilized, then washed thoroughly and placed with basal side facing downward for getting water source for root germination. When the roots of *A. cepa* L. were about 2 cm long, they were aseptically transferred to 100 µg/ml of isolated peptides from one week old germinating mung bean (*V. radiata*) seedlings and incubated for 12h at 25° C. Control roots were simultaneously treated with water to compare with peptides. Analysis of mitotic index were performed as described in details in Chapter VII Section 7.2.4

## **8.2.5 Plant materials for embryo and pith culture**

### **8.2.5a Carrot pith culture:**

Carrot taproots purchased from the local market were used as plant materials for the study. The carrots were chosen according to their tenderness, smoothness and freshness. After removing 5 cm sections from both ends, the carrots, were peeled and washed followed by immersion in 70 % ethanol for 20 min. Sterilization was carried out in 3% sodium hypochlorite solution for 15 min, to which two drops of Tween 20 were added. This was followed by rinsing three times in sterile distilled water. About 2 mm thick discs were cut from the carrots in such a way so as to include the xylem, phloem and the cambium. The cubes were inoculated onto the culture medium, maintaining polarity.

### **8.2.5b Excised and cotyledonary embryo culture of *Cicer arietinum*:**

Seeds of Kabuli type chickpea (*Cicer arietinum* L.) were surface sterilized as above, seed coats were removed aseptically and cotyledonary embryo or excised embryo proper were used for tissue culture.

### **8.2.5c Culture medium and culture conditions:**

The culture medium used for all experiments was based on MS medium (Murashige and Skoog, 1962) to which 0.8% Difco Bacto Agar and different concentration of peptides were added, but no plant growth regulators. The control consisted of basal MS medium supplemented with 2, 4-D and BAP. Medium pH was adjusted to 5.8 with KOH before adding the agar. All media were sterilized by autoclaving at 121° C and 101 KPa for 20 min. Cultures were incubated at 24 ± 2° C and exposed for 16 h per day to an illumination of 16  $\mu\text{molm}^{-2}\text{s}^{-1}$  provided by white fluorescent lamps.

## **8.2.6 Determination of peroxidase and total protein**

Crude Peroxidase extracts were prepared by grinding a fresh tissue in 0.1 M phosphate buffer (pH 7.0). The homogenate was centrifuged at 15,000g for 15 min. The total Guaiacol Peroxidase activity of supernatant was determined spectrophotometrically at

470 nm (Krsnik-Rasol, 1991). Enzyme activity is expressed as  $A_{470}$  (g. fresh weight)<sup>-1</sup> min<sup>-1</sup>.

The total protein in the supernatant was determined by the method of Lowry *et al.* (1951) using Bovine Serum Albumin Fraction-V as standard.

### 8.3 RESULT AND DISCUSSION

In spite of the importance attained by plant tissue culture and of many studies that have been conducted on related developmental process, there are still many aspects of growth and dedifferentiation that are not fully understood. Recent progress achieved on understanding the interaction between exogenously added plant growth regulators over the concentration of endogenous hormones, together with the involvement of sensitivity of the tissues to particular hormone groups, might help clarifying the occurrence of divergent patterns in callus induction and somatic embryogenesis (Jimenez 2005). In the present investigation, callus induction in chickpea was observed onto MS media containing different concentrations and combination of 2,4-D, NAA, IAA, BAP and Kinetin within 20 days of incubation of cotyledonary embryo explants depending upon the concentration and combination of hormones. Callus induction was noticed in all media formulations. But there was a wide range of variation in percentage of callus formation and average fresh weight of callus. The highest percentage of callus induction (90%) was observed on MS media containing 3.0 mg/l 2, 4-D and 1 mg/l BAP. The results are in concordance with Huda *et al.* (2006) where they showed that 2, 4-D and BAP were the most ideal hormonal combination in MS medium for induction of callogenesis in *Cicer arietinum* L. Highest callus growth in terms of fresh weight (0.576 g) was observed in MS medium fortified with 2 mg/l 2,4-D, 0.5 mg/l BAP and 100 µg/ml of rice peptides. Colour of the calli was mostly light brown to whitish green and light green. It was observed that only light green calli produced shoot buds. From this study it was observed that 2,4-D without cytokinin could induce callus but for better proliferation auxin (2,4-D, NAA and IAA) and cytokinin were required. Maximum shoot buds were proliferated from MS media containing 3 mg/l 2,4-D + 2 mg/l BAP. The shoot buds first appeared as nodular growth from cotyledonary axils after 22 days of culture and at the end of one month these nodules increased in size and produced leaf primordia.



Figure 8.1



Figure 8.2



Figure 8.3



Figure 8.4



Figure 8.5



Figure 8.6



Figure 8.7



Figure 8.8



Figure 8.9



Figure 8.10




Figure 8.11



Figure 8.12



Figure 8.13






Figure 8.14

Figure 8.1 = Treatment with BAP; Figure 8.2 = Treatment with BAP + 24h Peptides; Figure 8.3 = Treatment with BAP+ 32h Peptides; Figure 8.4 = Treatment with BAP + 40h Peptides; Figure 8.5 = Treatment with BAP + 48h Peptides; Figure 8.6 = Treatment with BAP + 56h Peptides; Figure 8.7 = Treatment with BAP + 64h Peptides; Figure 8.8 = Treatment with BAP + 72h Peptides; Figure 8.9 = Treatment with BAP + 5 day Peptides; Figure 8.10 = Treatment with BAP + 6 day Peptides; Figure 8.11A = Normal culture; Figure 8.11B = Treatment with BAP + 6 day Peptides; Figure 8.12 = Normal carrot pith culture; Figure 8.13 = Culture with 6 day Peptides ; Figure 8.14 = One dimension paper chromatography of germinating mung bean



**Figure 8.15** Increased fresh weight of cotyledonary embryo after application of phytohormones and peptides of *V. radiata* isolated from different germination hours




**Figure 8.16** Number of multiple shooting of cotyledonary embryo after application of *V. radiata* peptides isolated from different germination stages (24h – 144h days after soaking)

No remarkable enhancement of callogenesis was observed from our experiments if the peptides of rice, wheat, mung and chickpea were incorporated in both auxin and cytokinin deprived medium. In general, percentage of callus induction was strikingly lower when only auxin or cytokinin is present in the medium (also without peptides). In this condition, when the above mentioned peptides were combined with either of these two hormones (2, 4-D or BAP) prominent increase of callus tissue was noticed and peptides isolated from one week old *Vigna radiata* was most responsive. As the bioactivity of *Vigna radiata* peptides were observed to be most prominent among the four, only *Vigna radiata* peptides isolated from different germination hours of growing seedlings were taken for further bioassays. Induction of callus gradually increased in MS media fortified with 3 mg/l of 2,4-D and 100 µg/ml of peptides isolated from 8 to 64 hours and the maximum response was obtained at 2,4-D + 64 h peptides where callogenesis was observed in 86% of the explants. The minimum response of 58% callogenesis was observed at 2, 4-D + 5 days peptides combination whereas the callus induction was only 49% when only 2, 4-D alone was present in MS medium. So the results of the present study conclusively demonstrated that the induction of callogenesis was amplified significantly when peptides of germinating mung bean were associated with auxin. Restricted callogenesis was observed at cotyledonary nodes when the same peptides were combined with BAP (Figure 8.11 A & B). In BAP combined peptide medium, the calli were fragile, light green and embryogenic but no callogenesis was documented when BAP was only present. But peptides of *V. radiata* isolated from germination stages could efficiently enhance the weight of cotyledonary embryo (Figure 8.15). Peptides in BAP combining medium enhanced the potency of regeneration of multiple shoots (Figure 8.1 – 8.10) and maximum percentage of multiple shooting was observed at 1 mg/l BAP + 72 h peptides with almost 79% regeneration capacity (Figure 8.16). Maximum multiple shooting was only 53% when BAP alone was treated with explants (Figure 8.16). So in true sense germination induced peptides improved the bio-efficacy of both the classical hormone 2,4-D and BAP. But peptide alone has no capacity to induce and proliferate callogenesis and multiple shooting in *Cicer arietinum*. Also it was further documented that BAP (1 mg/l) combined peptide (100 µg/ml.) medium reduced the number of multiple shoots per plant, though the same combinations enhanced



Interphase



Prophase; Early (1), Late (2)



Metaphase [Equatorial View]



Metaphase [Polar View]



Anaphase

Figure 8.17 Cell division stages of *Allium cepa* L. root tip




Figure 8.18 Mitotic index of *Allium cepa*

the percentage of regeneration of multiple shoots from cotyledonary explants when compared with BAP (1 mg/l) alone.


Legumes generally and chickpea particularly is supposed to be recalcitrant and difficult to manipulate with *in vitro* cultures. The alteration in morphogenesis that was achieved due to application of plant peptides with either auxin or cytokinin may be considered significant because it was required to overcome the recalcitrant behaviour of chickpea. Though in chickpea callogenesis was not accomplished only by peptides due to this obstacle, in carrot disc culture restricted callogenesis (Figure 8.13) was observed in lower frequency (16.82%) by the application of wheat peptide alone in MS medium without the combination of any other hormones. Generally in carrot optimum callogenesis was observed in combination with 3 mg/l of 2, 4-D and 1 mg/l of BAP (Figure 8.12), as established from our experimental results and the findings were also corroborated with the views of other developmental biologists (Jimenez *et al.* 2005). No callus induction has been documented in carrot root disc culture in MS without any hormones or peptides.

Form the above discussion it may be concluded that isolated peptides somehow mimicked or influenced the callus inducing patterns of chickpea or carrot, regulated by auxin. The results were again reasonably fascinating when the peptide fractions of different germination hours (0 h. to 6 days) were subjected for incubation with mitotically dividing meristematic zones of *Allium cepa* (Figure 8.17) for observing their direct effect on cell division. Like callus induction mitotic index of *Allium cepa* root tip was also increased with the application of *Vigna* peptides isolated from germinating and post-germinating phases of 0 h to 7 days of seedling age. Mitotic index was maximized with peptides representing 64 h of seedling age and after 3 days; the effect of isolated peptides was gradually retarded. When the bioactivities of 64 h peptides were further elucidated with paper chromatographic separations (Figure 8.14), it was observed that the mitotic index drastically increased with two fractions ( $R_f$  zone1 and  $R_f$  zone3) but the total improvement was the reflection of enhancement of prophase. This is again fairly interesting because it shows that a preliminary signal induced the mitotically active cells to enter in dividing phase (M phase) but due to lack of some components, most of them


were specifically arrested in prophase (Figure 8.18). In contrast, application of peptides separated from R<sub>f</sub> zone 4 reduced the whole mitotic index of the cells.

Cell cycle is a critical and disciplined phenomenon, which generally shows no bypasses towards the decisive result. The basic machinery of cell cycle is highly conserved. In particular, many cellular events during this cycle progression are controlled by cyclin-dependent kinases (CDKs). Plants possess a unique class of CDKs (B-type CDKs) with preferential protein accumulation at G<sub>2</sub>/M-phases and control the entry into mitosis from G<sub>2</sub> of interphase (Fobert *et al* 1996, Segers *et al* 1996). The regulations of cell cycle control of animals are subsequently different in the embryonic and somatic cells (Eckardt, 2001). Sauter *et al.* (1998) hypothesized that zygotic regulation of the first cell cycle may contribute to a greater adaptive ability of plants during early embryogenesis, unlike animal embryos in which the cell division is more rigidly fixed before fertilization. This analysis also might lead to the prediction that embryonic tissues of plants are more totipotent than animals with greater morphogenic plasticity which might be due to the contribution of diffusible peptides in addition to classical hormones. Endogenous bioactive peptides, *i.e.* plant-derived peptides that evoke specific cellular responses, provide the most direct evidence for a general role of peptides in the regulation of plant growth and development. Already the peptides like POLARIS, Phytosulfokine, ENOD 40, Rapid Alkalinization Factor (RALF) and Clavata 3 (CLV3) have pivotal role in plant growth and development (Schaller, 2001). On the other hand, different physiological upsets, like wound or pathogen (e.g. – *Agrobacterium tumifaciens*) invasion, can stimulate mitotic division.

As these peptides of ultra low molecular weight range were functional from extracellular medium and showed a direct effect on mitotic index, they must have diffusible properties, which were thought to be much better suited to penetrate the rigid cell walls between adjacent cells as compared to large peptide hormones. Research in recent years, however, has indicated that peptides may be widely used as chemical signals in plants as well. Like Phytosulfokine these peptides led the cells towards the entry into mitotic division phases but unlikely unable to complete the rest. One thing we must take in care that the source of peptides is mainly developing embryo and the test material is somatic tissue. Therefore, it is quite logical to say that these peptides



**Figure 8.19** Peroxidase activity in cotyledonary embryo after application of *V. radiata* peptides isolated from different germination stages (24h – 144h days after soaking)



**Figure 8.20** Protein concentration of cotyledonary embryo after application of *V. radiata* peptides isolated from different germination stages (24h – 144h days after soaking)

promoted the cells towards the divisional phase by creating immense impact over cell division in developing embryo. As a matter of fact, the peptides of R<sub>f</sub> zone-4 have negative regulation on both cell division and amylase synthesis. So, it is reasonably justified to assume that living system is so disciplined and balanced that it always maintains a feedback inhibition within them. Though it is somewhat surprising for the same molecule to inhibit both of the above actions but not impossible. Our investigations with germinating sonamung support the above view.

Like all developmental processes, callogenesis and organogenesis was also associated with alteration of oxidative stress. Peroxidase activity as well as thiol and ascorbic acid contents can be considered as important metabolic markers of oxidative stress during development of somatic embryo (Norgaard, 1997). When the peptides were tested for regeneration capacity and callogenesis in *Cicer* tissue, it was observed that peroxidase activity increased considerably (Figure 8.19) along with enhanced protein content (Figure 8.20) in the callus at the end of the culture. But the highest peroxidase activity was determined in the embryogenic line at 6 days of culture in the presence of growth regulator BAP and peptides at a concentration of 100 µg / ml. The minimum peroxidase activity was estimated in habituated tissue grew in presence of 2,4-D and BAP and formed white swollen aggregates of shoots and hypocotyls with strongly reduced potential for somatic embryogenesis. An increase in peroxidase activity was also recorded in those flasks, where tissue became looser and after few days embryoids were noticed. After formation of globular to heart shaped embryoids, peroxidase activity was dropped 3 to 4 times, as compared with callus tissues from which they originated. It was also detected that peroxidase activity was enhanced during multiple shooting formations in cytokinin peptide combined medium. When the tissue specific activity of peroxidase was studied, it was noticed that the peroxidase activity was about three times higher in bushy multiple shoots than that of single leaf or shoot. This is in agreement with Krsnik-Rasol (1991) where they claimed that bushy transformants and crown gall tissue extracts were 6 to 7 times higher level of enzyme activity than in normal leaves.

As a conclusion, these results shed light on the bioactivity of peptides which play a significant role in callogenesis and morphogenic alteration. These data show that peroxidase activity can also be associated with organogenesis under the influence of

peptides of ultra low range. Further investigations will be required to establish the molecular mechanisms of regulation of peptides controlling morphogenesis and cell division.

## REFERENCES

- Eckardt NA. 2001. A calcium-regulated gatekeeper in phloem sieve tubes. *Plant Cell*. 13:989-992
- Fobert PR, Gaudin V, Lunness P, Coen ES, Doonan JH. 1996. Distinct classes of cdc2-related genes are differentially expressed during the cell division cycle in plants. *Plant Cell*. 8:1465-1476
- Huda MS, Wilding JP, Pinkney JH. 2006. Gut peptides and the regulation of appetite. *Obes. Rev.* 7:163-182
- Jimenez VM. 2005. Involvement of plant hormones and plant growth regulators on in vitro somatic embryogenesis. *Plant Growth Reg.* 47(2-3):91-110
- Krsnik-Rasol M. 1991. Peroxidase as a developmental marker in plant tissue culture. *Int. J. Dev. Biol.* 35(3):259-63
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with the *Folin phenol* reagent. *J. Biol. Chem.* 193:265-275
- Murashige T, Skoog F. 1962. A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Plant Physiol.* 15:473-497
- Nordstrom A, Tarkowski P, Tarkowska D. 2004. Auxin regulation of cytokinin biosynthesis in *Arabidopsis thaliana*: a factor of potential importance for auxin-cytokinin-regulated development. *Proc. Natl. Acad. Sci. USA.* 101:8039-8044
- Norgaard JV. 1997. Somatic embryo maturation and plant regeneration in *Abies nordmanniana*. *Plant Sci.* 124:211-221
- Ryan CA, Pearce G, Scheer J, Moura DS. 2002. Polypeptide hormones. *Plant Cell*. 14:251-264
- Sauter M, von Wiegen P, Lorz H. 1998. Cell cycle regulatory genes from maize are differentially controlled during fertilization and first embryonic cell division. *Sex. Plant Reprod.* 11:41-48
- Schaller MD. 2001. Biochemical signals and biological responses elicited by the focal adhesion kinase. *Biochem. Biophys. Acta.* 1540:1-21

- Segers G, Gadisseur I, Bergounloux C, de Almeida Engler J, Jacqmird A, Van Montagu M, Inze D. 1996. The *Arabidopsis* cyclin-dependent kinase gene *cddbAt* is preferentially expressed during S and G2 phases of the cell cycle. *Plant J.* 6:01-12
- Shiu SH, Bleecker AB. 2001. Receptor-like kinases from *Arabidopsis* form a monophyletic gene family related to animal receptor kinases. *Proc. Natl Acad. Sci. USA.* 98:10763-10768
- Souter MA, Pullen M, Topping JF, Zhang X, Lindsey K. 2004. Rescue of defective auxin-mediated gene expression and root meristem function by inhibition of ethylene signalling in sterol biosynthesis mutants of *Arabidopsis*. *Planta.* 219:773-783
- Stepanova AN, Alonso JM. 2005. Ethylene signaling and response pathway: A unique signaling cascade with a multitude of inputs and outputs. *Physiol. Plant.* 123:195-206
- Trewavas A, Knight M. 1994. Mechanical signaling, calcium and plant form. *Plant Mol. Biol.* 26:1329-1341
- Vogel JP, Woeste KW, Theologis A, Kieber JJ. 1998. Recessive and dominant mutations in the ethylene biosynthetic gene *ACS5* of *Arabidopsis* confer cytokinin insensitivity and ethylene overproduction, respectively. *Proc. Natl. Acad. Sci. USA.* 95:4766-4771