

## LIST OF TABLES

- Table 1.1:** Major plant drugs for which no synthetic one is available.
- Table 1.2:** Diversity of ethnomedicinal plants in Darjeeling Himalayas.
- Table 1.3:** Threatened medicinal plants of the Darjeeling Himalayas.
- Table 1.4:** List of species of *Swertia* and their geographical distribution in Darjeeling and Sikkim Hills.
- Table 1.5:** Ethnomedicinal uses of some species of *Swertia*.
- Table 1.6:** Important phases occurring in the life cycle of *Swertia chirayita*. Data were recorded from 10 uniformly grown plants.
- Table 1.7:** Meteorological data recorded during the year 2009 at Darjeeling.
- Table 1.8:** Meteorological data recorded during the year 2010 at Darjeeling.
- Table 1.9:** Meteorological data recorded during the year 2011 at Darjeeling.
- Table 1.10:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in total chlorophyll (T.C.), chlorophyll a (C. a) and chlorophyll b (C. b) [mg/g fresh weight] contents in leaves of chirata.
- Table 1.11:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in protein (Pr.) and free amino acids (F.A.A.) [mg/g fresh weight] content in leaves of chirata.
- Table 1.12:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in soluble (Sol.) and insoluble (Insol.) carbohydrate [mg/g fresh weight] content in leaves of chirata.
- Table 1.13:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in RNA and DNA [µg/g fresh weight] content in leaves of chirata.
- Table 1.14:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in lipid (L) [mg/g fresh wt] and total phenol (T.P.) [µg/g fresh wt] content in leaves of chirata.
- Table 1.15:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in bitter principles content (in %) of root (R), stem (S) and leaf (L) of chirata plant.

**Table 1.16:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in plant height (P. H., cm) and stem circumference (S. C., cm) of chirata plant.

**Table 1.17:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of leaves (L) and number of branches (B) per plant of chirata.

**Table 1.18:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of nodes (N) and internodal distance (I. D., cm) of chirata plant.

**Table 1.19:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on number of flowers per plant, total length of the plant (from root-tip to stem-tip), fresh weight and dry weight of chirata plant.

**Table 1.20:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in total chlorophyll (T.C.), chlorophyll a (C.a) and chlorophyll b (C.b) [mg/g fresh weight] content in leaves of chirata.

**Table 1.21:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in protein (Pr.) and free amino acids (F.A.A.) [mg/g fresh weight] content in leaves of chirata.

**Table 1.22:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in soluble (Sol.) and insoluble (Insol.) carbohydrate [mg/g fresh weight] content in leaves of chirata.

**Table 1.23:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in RNA and DNA [µg/g fresh weight] content in leaves of chirata.

**Table 1.24:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in lipid (L in mg) and total phenol (T. P. in µg/ml) content in leaves of chirata.

**Table 1.25:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in bitter principles content (in %) of root (R), stem (S) and leaf (L) of chirata plant.

**Table 1.26:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in plant height (P.H., cm) and stem circumference (S. C., cm) of chirata plant.

**Table 1.27:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of leaves (L) and number of branches (B) per plant of chirata.

**Table 1.28:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of nodes (N) and internodal distance (I. D., cm) of chirata plant.

**Table 1.29:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on number of flowers per plant, total length of the plant (in cm; from root-tip to stem-tip), fresh weight (in gm) and dry weight (in gm) of chirata plant.

**Table 1.30:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in total chlorophyll (T.C.), chlorophyll a (C.a) and chlorophyll b (C.b) [mg/g fresh weight] content in leaves of chirata.

**Table 1.31:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in protein (Pr.) and free amino acids (F.A.A.) [mg/g fresh weight] content in leaves of chirata.

**Table 1.32:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in soluble (Sol.) and insoluble (Insol.) carbohydrate [mg/g fresh weight] content in leaves of chirata.

**Table 1.33:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in RNA and DNA [µg/g fresh weight] content in leaves of chirata.

**Table 1.34:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in lipid (L in mg) and total phenol (T. P. in µg/ml) content in leaves of chirata.

**Table 1.35:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in bitter principles content (in %) of root (R), stem (S) and leaf (L) of chirata plant.

**Table 1.36:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in plant height (P.H., cm) and stem circumference (S.C., cm) of chirata plant.

**Table 1.37:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of leaves (L) and number of branches (B) per plant of chirata.

**Table 1.38:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on changes in number of nodes (N) and internodal distance (I. D., cm) of chirata plant.

**Table 1.39:** Effect of GA<sub>3</sub> (100 and 200 µg/ml), IAA (100 and 200 µg/ml), KIN (100 and 200 µg/ml), MH (100 and 200 µg/ml), SADH (100 and 200 µg/ml) and ABA (10 and 15 µg/ml) on number of flowers per plant, total length of the plant (in cm; from root-tip to stem-tip), fresh weight (in gm) and dry weight (in gm) of chirata plant.