

## LIST OF FIGURES

Fig I.1	Active site structure of sMMO in its reduced form.	<b>5</b>
Fig I.2	Active site structure of To4MO, a non-heme multicomponent oxygenase containing diiron (III) active site in its resting state (left) and a putative 'complex Q' which is active oxidising species [49].	<b>7</b>
Fig I.3	Active site structure of Rieske dioxygenases. The Rieske [2Fe-2S] cluster is linked to the oxygenase component via an H-bonding network by an aspartate residue.	<b>8</b>
Fig I.4	A family of non-heme based iron(II) system including N,N,O donor state.	<b>12</b>
Fig I.5	Diiron(III) core structure that used as the mimic of sMMO.	<b>14</b>
Fig II.1	<sup>1</sup> H NMR spectra of the ligand bpmen; Inset: Expanded aromatic region.	<b>26</b>
Fig II.2	UV-Vis spectra of the complexes ( <b>1-3</b> ) in acetonitrile at room temperature (concentration of the complexes = 1.0 mM).	<b>27</b>
Fig II.3	UV-Vis spectra of the complex <b>1</b> in acetonitrile at room temperature (concentration of the complex = 1.0 mM).	<b>28</b>
Fig II.4	ESI-MS spectra of the complex [Fe(bpmen)(μ-O)(μ-OH)Fe(bpmen)](ClO <sub>4</sub> ) <sub>3</sub> ( <b>2</b> ).	<b>29</b>
Fig II.5	ESI-MS spectra of the complex [Fe(bpmen)(μ-O)(μ-CH <sub>3</sub> COO)Fe(bpmen)](ClO <sub>4</sub> ) <sub>3</sub> ( <b>3</b> ).	<b>29</b>
Fig II.6	Ortep plot of the complex <b>1</b> [Fe(bpmen)(μ-O)FeCl <sub>3</sub> ] showing 50% probability ellipsoids.	<b>30</b>
Fig II.7	UV-vis spectral changes in the reaction of 1.0 mM [Fe <sub>2</sub> (μ-O)(μ-OH)(bpmen) <sub>2</sub> ]( <b>2</b> ) and H <sub>2</sub> O <sub>2</sub> (10 mM) in presence of 100 mM benzyl alcohol in acetonitrile at 298K.	<b>37</b>
Fig II.8	<sup>1</sup> H NMR of the ligand bpmen in CDCl <sub>3</sub> .	<b>40</b>
Fig II.9	<sup>1</sup> H NMR of the ligand bpmen in CDCl <sub>3</sub> (Expanded aromatic region).	<b>40</b>
Fig III.1	UV-Vis spectra of the complex <b>2</b> in acetonitrile at room temperature (concentration of the complex = 2.5X 10 <sup>-4</sup> mM).	<b>51</b>
Fig III.2	Competitive experiments for the oxidation of different olefin pairs by <b>2</b> /H <sub>2</sub> O <sub>2</sub> .	<b>57</b>
Fig III.3	UV-vis spectral changes obtained upon the addition of H <sub>2</sub> O <sub>2</sub> (30 mM) to a solution of complex <b>2</b> (1 mM) and <i>tert</i> -butyl acrylate (20 mM) in neat acetonitrile at 298 K. Inset: Change of absorbance of the 480 and 550 nm bands monitored with time	<b>61</b>
Fig III.4.	UV-vis spectrum of the complex <b>2</b> (red line) and that of the species obtained at the end of the aromatic hydroxylation (blue line) in dry acetonitrile at 298K. Electronic spectra of <b>2</b> and <b>2'</b> in the visible region in dry acetonitrile at 298 K.	<b>62</b>
Fig III.5	UV-vis spectral changes obtained upon the addition of H <sub>2</sub> O <sub>2</sub> (30 mM) to a solution of complex <b>2</b> (1 mM), AcOH (1 mM) and <i>tert</i> -butyl acrylate (20 mM) in neat acetonitrile at 298 K. Inset: Change of absorbance of the 480 and 550 nm bands monitored with time	<b>62</b>
Fig III.6	X-band EPR spectrum obtained after the addition of 10 mM H <sub>2</sub> O <sub>2</sub> to an acetonitrile	<b>63</b>

	solution of <b>2</b> (1.0 mM) at 298 K followed by rapid freezing of the solution.	
Fig III.7	ESI-MS spectrum obtained immediately after the addition of 10 mM H <sub>2</sub> O <sub>2</sub> to an acetonitrile solution of <b>2</b> (1.0 mM) at 298 K (Above). The experimental and theoretical isotope distribution patterns for the peaks at m/z 342 and 361 (Below).	64
FigIV.1	UV-visible changes for the reaction of <b>2</b> (1.0 mM), benzene (300 mM) and H <sub>2</sub> O <sub>2</sub> (10 mM) in acetonitrile at 298K. Inset: Absorbance changes at 650 nm with time after the addition of H <sub>2</sub> O <sub>2</sub> .	79
Fig IV.2	Decay of the 650 nm band with time. The final spectrum corresponding to the [(bpmen)(H <sub>2</sub> O)Fe(μ-O)Fe(OH)(bpmen)] <sup>3+</sup> ( <b>2'</b> ) complex is shown in red. Inset: The rate of formation of the iron(III) phenolate species in acetonitrile at 298K as a function of H <sub>2</sub> O <sub>2</sub> concentration.	80
Fig IV.3	Experimental and theoretical isotope distribution patterns.	81
Fig IV.4	Rate of formation of phenolate as a function of benzene concentration	82
Fig IV.5	Plot of first-order rate constants against 1/T to determine activation parameters.	82
Fig IV.6	UV-vis spectrum of the complex <b>2</b> (1.25 x 10 <sup>-4</sup> M) (red line) and that of the species obtained at the end of the aromatic hydroxylation (blue line) in dry acetonitrile at 298K. Inset: Electronic spectra of <b>2</b> (1 mM) and <b>2'</b> in the visible region in dry acetonitrile at 298 K.	83
Fig IV.7	Electronic changes observed after the addition of additional H <sub>2</sub> O <sub>2</sub> (10 mM) to the final diiron(III) complex (black line).	84
Fig IV.8	X-band EPR spectrum obtained after the addition of 10 mM H <sub>2</sub> O <sub>2</sub> to an acetonitrile solution of <b>2</b> (1.0 mM) at 298 K (red line). EPR spectrum obtained after the addition of 10 mM H <sub>2</sub> O <sub>2</sub> to an acetonitrile solution of <b>2</b> (1.0 mM) and benzene (300 mM) at 298 K (black line).	87
Fig V.1	Active site structure of sMMO in its reduced form.	96
Fig V.2	Currently accepted mechanism of methane oxidation by sMMO.	96
Fig V.3	<sup>1</sup> H NMR spectra of the ligand <i>iso</i> -bpmen; Inset: Expanded aromatic region.	98
Fig V.4	Electronic spectrum of complex <b>1</b> in acetonitrile	99
Fig V.5	ORTEP diagram of complex <b>1</b> showing 50% probability thermal ellipsoids and the labelling scheme for selected atoms. The atoms with the asterisks (*) are symmetry generated. Hydrogen atoms and the counter anions are removed for clarity.	100
Fig V.6	Packing arrangement in <b>1</b>	102
Fig VI.1	Electronic spectrum of complex <b>1</b> in acetonitrile	118
Fig VI.2	ORTEP plot of an independent cation ( <b>1</b> <sup>3+</sup> ) at 50% probability level. The perchlorate anions were omitted for clarity. The atoms marked with # are symmetry generated.	119
Fig VI.3	Packing diagram of <b>1</b> -ClO <sub>4</sub>	121
Fig VI.4	Structure of iron (II) complexes of TPA and BPMEN	124

---