

LIST OF TABLES

Table No.	Title of tables	Page No.
1.	Effect of BDE on the body weight, relative spleen weight, number of splenocytes, number of plaques formed, hemagglutination titre value and number of peritoneal macrophages of mice after 15 days (S1) and 45 days (S2) of treatment	58
2.	Effect of BDE on the body weight, relative spleen weight, number of splenocytes, number of plaques formed, hemagglutination titre value and number of peritoneal macrophages of mice after 90 days (S3), 135 days (S4) and 180 (S5) days of treatment.	58
3.	Serum concentrations of different Th1 (IL-2 and IFN- γ) and Th2 (IL-4 and IL-10) cytokines of mice that are treated with different doses of BDE for 15 (S1) and 45 (S2) days	62
4.	Serum concentrations of different Th1 (IL-2 and IFN- γ) and Th2 (IL-4 and IL-10) cytokines of mice that are treated with different doses of BDE for 90 (S3), 135 (S4) and 180 (S5) days	62
5.	Effect of BDE on body weight and relative weight of testis and other sexual organs after subacute and subchronic doses	68
6.	Effect of BDE on body weight and relative weight of testis and other sexual organs after chronic doses	69
7.	Effect of BDE on different biochemical parameters of testis and other sexual organs after sub-acute (15 and 45 days) and sub-chronic doses (90 days)	71
8.	Effect of BDE on different biochemical parameters of testis and other sexual organs after chronic doses (135 and 180 days)	72
9.	Histomorphometric parameters of seminiferous tubules	75
10.	Effects of the BDE on fertility of Swiss albino mouse after different days of treatment	76
11.	Effects of the BDE on fecundity of Swiss albino mouse after different days of treatment	76
12.	Scavenging of reactive oxygen species, iron chelating and lipid peroxidation	89

	inhibition activity (IC ₅₀ values) of <i>Diplazium esculentum</i> (MDE) and reference compounds.	
13.	Qualitative analysis of the phytochemicals of <i>D. esculentum</i>	90
14.	Acute toxicity study of BDE on body weight and relative organ weight of Swiss albino mouse	91
15.	Chronic toxicity study (180 days) of BDE on body weight and relative organ weight of Swiss albino mouse	92
16.	Biochemical measurements in serum of mice fed with BDE for 15, 45 and 90 days	93
17.	Biochemical measurements in serum of mice fed with BDE for 135 and 180 days	94
18.	Effect of CDE and BDE on the HA titre value in mice after 180 days of treatment	98
19.	Effect of CDE and BDE on body weight and relative weight of male reproductive organs after chronic dose (180 days)	102
20.	Effect of CDE and BDE on different biochemical parameters of male reproductive organs after 180 days of treatment	103
21.	Comparison of the biochemical parameters in serum of mice fed with CDE and BDE for 180 days	105

LIST OF FIGURES

Figure No.	Title of Figures	Page No.
1.	Principle of the cholinergic anti-inflammatory pathway	8
2.	Paradigm for the immunomodulatory activity of plant-derived medicines	12
3.	The distribution of <i>Diplazium esculentum</i>	26
4.	<i>Diplazium esculentum</i> in its natural habitat and collected frond (the main edible part)	27
5.	Effect of BDE on the primary cultured splenocytes after 24 h of incubation.	59
6.	Effect of BDE on the primary cultured splenocytes after 48 h and 72 h of incubation.	60
7.	MTT splenocytes proliferation assay demonstrates dose-dependent increase in the percentage inhibition of splenocytes proliferation in the BDE treated splenocytes.	60
8.	The hemolytic activity of BDE was increased significantly in a dose-dependent manner in case of mouse erythrocytes.	61
9.	Effect of different concentrations (0-200 µg/ml) of BDE on different Th1 (IL-2 and IFN-γ) and Th2 (IL-4 and IL-10) cytokine production by Con A induced splenocytes.	63
10.	Hypo-osmotic swelling test (HOST) of spermatocytes demonstrates dose-dependent decrease in the percentage of live sperm in the mice treated with BDE for 15, 45, 90, 135 and 180 days.	65
11.	MTT reduction assay of spermatocytes demonstrates dose-dependent increase in the percentage inhibition of sperm viability in the mice treated with BDE for 15, 45, 90, 135, and 180 days.	67
12.	Histology of the testis of control mice	73
13.	Histology of the testis of mice treated with 320 mg/kg bw of BDE for 135 days	74
14.	Histology of the testis of mice treated with 320 mg/kg bw of BDE for 180 days	74
15.	Acetylcholinesterase activity in BDE treated mice	77

16.	Acetylcholinesterase inhibitory activity of <i>D. esculentum</i> extract	78
17.	NADH oxidase inhibitory activity of <i>D. esculentum</i> extract	79
18.	Absorbance value of the MDE in the linoleic acid emulsion using FTC method	80
19.	Total antioxidant activity of MDE and reference compound trolox on decolourization of ABTS radical cation	81
20.	DPPH radical scavenging activity of <i>D. esculentum</i> extract	82
21.	Hydroxyl radical scavenging activities of MDE and the reference compound mannitol	82
22.	Scavenging effect of MDE and the standard quercetin on superoxide radical	83
23.	Nitric oxide inhibition by MDE and the standard curcumin	84
24.	Effect of MDE and sodium pyruvate on the scavenging of H ₂ O ₂	84
25.	The peroxynitrite anion scavenging activity of MDE and the standard gallic acid	85
26.	Effects of MDE and the standard lipoic acid on the scavenging of singlet oxygen	86
27.	Hypochlorous acid scavenging activities of MDE and the standard ascorbic acid	86
28.	Effects of <i>D. esculentum</i> plant extract and EDTA on Fe ₂ ⁺ -ferrozine complex formation	87
29.	Reducing Power Assay of MDE and the standard ascorbic acid	88
30.	Lipid peroxidation inhibition by MDE and the standard trolox	88
31.	Histological architecture of liver in Control (A) and BDE treated mouse (B)	95
32.	Histological architecture of kidney in Control (A) and BDE treated mouse (B)	95
33.	Comparison between the numbers of splenocytes of CDE vs BDE treated mice after 180 days of treatment	96
34.	Comparison between the numbers of plaque forming cells in the spleen of CDE and BDE treated mice	97
35.	Result shows dose-dependent decrease in the number of CDE and BDE treated	98

	splenocyte after 24 hours of incubation	
36.	Result shows dose-dependent decrease in the number of splenocyte after 48 hours of incubation	99
37.	Result shows dose-dependent decrease in the number of splenocyte after 72 hours of incubation	99
38.	MTT assay showing the effect of CDE and BDE on splenocytes	99
39.	Comparison between the hemolytic activity of CDE and BDE on human erythrocytes	100
40.	Comparison between serum IgM concentration of CDE and BDE treated mice	100
41.	Comparison of the percentage of live sperm between CDE and BDE treated mice	101
42.	MTT assay showing the effect of CDE and BDE on sperm	101
43.	Results indicated dose-dependent decrease in the rate of hydrolysis of acetylthiocholine iodide substrate by acetylcholinesterase (CDE vs BDE)	104

LIST OF APPENDICES

APPENDIX – 1 (Kits used in experiments).....	152
APPENDIX – 2 (Photographs of some of the experiments conducted).....	161
APPENDIX – 3 (Publications).....	165