

LIST OF TABLES

- Table 1.** Nutrient content of rice per 100 gram portion
- Table 2.** Some common rice diseases
- Table 3.** Rice cultivars and its localization
- Table 4.** Morphological characterization of rice cultivars
- Table 5.** Disease index showing the establishment of natural disease
- Table 6.** Effects of antifungal compound from rice leaf extracts on radial growth of *D. oryzae*
- Table 7.** Pathogenicity test of *D. oryzae* on three rice cultivars following detached leaf inoculation technique
- Table 8.** Pathogenicity test of *D. oryzae* on three cultivars of rice plant following whole plant inoculation technique
- Table 9.** ELISA values showing reaction of anti-*D. oryzae* with antigens of healthy and artificially inoculated rice plants after every 24hrs
- Table 10.** Dot immunobinding assay of healthy and artificially inoculated leaf antigen of rice plants using PAb of *D. oryzae*.
- Table 11.** Genbank Accession Numbers and Geographic locations of the Ex-Type strains of *D. oryzae* that showed homology with the isolate R1.DO.01
- Table 12.** Population count of AM Fungi in rhizosphere of fifteen different rice cultivars and percentage colonization in root.
- Table 13.** Morphological and microscopical characters of dominant AMF spores
- Table 14.** *In vitro* pairing of PGPR isolates with foliar pathogen of rice- *Drechslera oryzae* for evaluation of antifungal activities.
- Table 15.** *In vitro* antagonistic tests of selected PGPR isolate against brown spot pathogen *D. oryzae*
- Table 16.** Total sugar content of rice leaves following treatments with PGPR
- Table 17.** Total chlorophyll content of rice leaves following treatments with PGPR
- Table 18.** Protein content of rice leaves following treatments with PGPR
- Table 19.** Total phenol content of rice leaves following treatments with PGPR
- Table 20.** Evaluation of Disease index for brown spot in rice plants following treatments with PGPR
- Table 21.** Peak results of Phytocassanes extracts from leaves of Black nuniya following treatment with *Bacillus altitudinus* (NAIMCC-B01485) and pathogen challenge

- Table 22.** Total sugar content of rice leaves following treatments with PGPF
- Table 23.** Protein content of rice leaves following treatments with PGPF
- Table 24.** Total phenol content of rice leaves following treatments with PGPF
- Table 25.** Evaluation of Disease index for brown spot in rice plants following treatments with PGPF
- Table 26.** Peak results of Phytocassanes from leaf extracts of rice plant (cv. Black Nuniya) following treatment with *T. harzianum* (NAIMCC-F-03288) and pathogen challenge
- Table 27.** Growth promotion of field grown rice plants following application of AMF (*Glomus fasciculatum*) and pathogen challenge
- Table 28.** Change in biochemical activity in rice cultivars following application of AMF (*Glomus fasciculatum*) and pathogen challenge
- Table 29.** Percent Disease Index in rice plants following treatment and pathogen challenge.
- Table 30.** Growth promotion in rice cultivars following application of bioinoculants
- Table 31.** Changes in biochemical components in rice cultivars following treatment with bioinoculants against challenged inoculation with the pathogen
- Table 32:** Effects of antifungal compound (Phytocassanes) from rice leaf extracts of (cv. Black Nuniya) following treatment with bioinoculants [*T. harzianum* (NAIMCC-F-03288), *G. fasciculatum* and *B. altitudinus* (NAIMCC-B01485)] and inoculation with *D. oryzae*
- Table 33.** Peak Value of Standard Phenolics
- Table 34:** Peak results of Phenolics extracts from rice leaf of (cv. Black Nuniya) following treatment with bioinoculants and challenge inoculated with foliar fungal pathogen (*D. oryzae*)
- Table 35.** Peak results of Phytocassanes extracts from rice leaves (cv. Black nuniya) following treatment with bioinoculants and challenge inoculation with foliar fungal pathogen (*D. oryzae*)

LIST OF FIGURES

- Figure 1:** Contribution of State in India's Rice Production
- Figure 2:** Rice cultivars grown in petriplates for viability test.
- Figure 3:** Maintenance of rice seedlings in glass house
- Figure 4:** Maintenance of rice cultivars in the field
- Figure 5:** Fifteen different rice cultivars showing morphology
- Figure 6:** Initial screening of rice cultivars grown in petriplates. (A) Germination ability, (B) Protein content and (C) Growth rate.
- Figure 7:** Growth rate of fifteen rice cultivars grown in field
- Figure 8:** Rice cultivars grown in experimental plot showing natural infection (Brown Spot) caused by *D. oryzae*
- Figure 9:** PAL (Phenylalanine ammonia lyase) (A) and Peroxidase (B) activities in healthy and naturally infected rice leaves.
- Figure 10:** Chitinase (A) and β -1,3-glucanase (B) activities in healthy and naturally infected rice leaf samples
- Figure 11:** Total phenol content of healthy and naturally infected rice cultivars.
- Figure 12:** Radial growth pattern of fungi in different media (7-10 days old).(A) PDA,(B)Oats meal agar,(C) Richard's agar.
- Figure 13:** Microscopic view of fungal isolate from infected rice leaves. (A)Conidium (40X), (B) Germinating conidium (40X), (C) Conidiophore bearing conidia (10X),(D) Enlarged view of (C 40X)
- Figure 14:** Pathogenicity test of rice cultivars against *D. oryzae*. (A&B)Whole plant C&D) Detached leaf
- Figure 15:** Serological assays of *Drechslera oryzae*; (A) Dot- blot , (B) Immunofluorescence of young mycelia treated with PAbs of *D. oryzae* and labelled with FITC, (C) (SDS-PAGE and (D) Western blot analysis of *D oryzae*
- Figure 16:** Chromatogram and sequence of 18S rDNA region *Drechslera oryzae* strain RI.DO.01 deposited in NCBI Genbank
- Figure 17:**18S r DNA sequence alignments of *Drechslera oryzae*(KT768092) with other exotypes isolate. The conserved regions of the gene are demonstrated in different colour.
- Figure 18:** Phylogenetic placement of *Drechslera oryzae*KT768092with other ex-type strains obtained from NCBI Gen Bank Database.

- Figure 19:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Brimful). (A) *Glomus* sp.; (B) *Glomus* sp.;(C) *Glomus* sp.;(D) *Glomus fasciculatum*; (E) *Glomus mosseae* (juvenile); (F) *Scutellospora* sp.;(G) *Glomus* sp.;(H) *Glomus mosseae* (mature).
- Figure 20:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Tulai panji).(A) *Glomus* sp. ;(B) *Glomus* sp.;(C) *Glomus* sp.;(D) ruptured spore of *Glomus* sp.;(E) *Scutellospora* sp.;(F) mature *Glomus* sp.;(G) *Glomus constrictum*
- Figure 21:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (UBKV-5).(A) *Acaulospora* sp.;(B) ruptured *Gigaspora* with conspicuous hyphal attachment;(C) *Scutellospora* sp.;(D) *Glomus* sp.;(E)*Scutellospora* sp.;(F) *Scutellospora* with hyphal attachment (magnified)
- Figure 22:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (UBKV-4).(A)*Scutellospora* sp.;(B) *Gigaspora* sp.;(C) *Glomus* sp.;(D)*Glomus* sp.;(E) Hyphal attachment of *Gigaspora* sp.;(F) *Scutellospora* sp.
- Figure 23:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Black Nuniya).(A) *Glomus* sp.;(B) *Glomus* sp.;(C) *Scutellospora* sp.;(D) *Glomus* sp.;(E) *Glomus* sp.; (F) *Acaulospora* sp.; (G) *Glomus* sp.;(H) *Glomus* sp.; (I) *Glomus* sp.
- Figure 24:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Champasari).(A) *Gigaspora* sp.;(B) *Glomus* sp.;(C) Ruptured spores of *Scutellospora* sp.;(D) *Scutellospora* sp.(matured)
- Figure 25:**Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Adde).(A) *Glomus constrictum*;(B) *Glomus constrictum*; (C) *Glomus mosseae*; (D) *Acaulospora* sp.; (E) *Acaulospora* sp.;(F) *Acaulospora* sp.;(G,H&I) *Glomus mosseae* in different angles
- Figure 26:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (UBKV-1).(A) *Gigaspora* sp.;(B) *Glomus* sp.;(C)*Glomus* sp.;(D)*Glomus* sp.
- Figure 27:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Swarna masuri).(A) *Scutellospora* sp.; (B) Ruptured *Glomus* sp.;(C) *Glomus* sp. with hyphae
- Figure 28:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Maiti). (A) *Scutellospora* sp.;(B) *Scutellospora* sp.;(C) *Gigaspora margarita*

- Figure 29:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Attheu).(A) *Glomus* sp.;(B) *Scutellospora* sp.;(C) *Glomus* sp.;(D) *Acaulospora* sp.;(E) *Glomus constrictum*;(F) Ruptured AMF.
- Figure 30:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Sanomasuri).(A) *Scutellospora* sp.;(B) *Scutellospora* sp.;(C) *Glomus* sp.;(D) *Gigasporarosea*.
- Figure 31:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Loknath 505).(A) *Glomus* sp. juvenile;(B) *Entrophospora* sp.;(C) *Glomus* sp.;(D) *Scutellospora* sp. with hyphal attachment
- Figure 32:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Kaberi 9090).(A) *Glomus* sp.;(B) *Glomus* sp.;(C) *Glomusbadium* ;(D) *Glomus badium*
- Figure 33:** Compound microscopic observation of Arbuscular Mycorrhizal Fungal spores obtained from rice cultivar (Gouraknath 509).(A) *Glomus fasciculatum*;(B) *Glomus multicaule*;(C) *Glomus* sp.;(*Glomus* sp.)
- Figure 34:** Root colonization of rice cultivars. (A-C) Brimful; (D-F) Black nuniya; (G-I) Champasari and (J-L) Attheu. (Ih- Intracellular hyphae, Vs- Vesicle, Ar- Arbuscule).
- Figure 35:** Root colonization of rice cultivars. (A-C) Maiti; (D-F) Sano masuri; (G-I) Tualai panji and (J-L) Swarnamasuri. (Ih- Intracellular hyphae, Irs- Intra radicle spore, Vs- Vesicle, Ar- Arbuscule).
- Figure 36:** Root colonization of rice cultivars. (A-C) UBKV-1; (D-F) UBKV-4; (G-I) UBKV-5 and (J-L) Adde. (Ih- Intracellular hyphae, Vs- Vesicle, Ar- Arbuscule)
- Figure 37:** Root colonization of rice cultivars. (A-C) Kaberi 9090; (D-F) Loknath 505; (G-I) Gouraknath 509. (Ih- Intracellular hyphae, Ca- Coiled arbuscule, Vs- Vesicle, Ar- Arbuscule)
- Figure 38:** Mass multiplication of AMF spores in maize plant (A-C) Pots; (D) Experimental field
- Figure 39:** Histopathological study of AMF colonized maize roots.(A) Vesicles and intracellular hyphae, (B) Spores bearing hyphae,(C) Vesicles,(D) Vesicle and Intracellular hyphae.(Vs- Vesicle, Ih-Intracellular hyphae, Sp-Spore)
- Figure 40:** *In vitro* antifungal activities of foliar fungal pathogen *Drechslera oryzae* against selected PGPR isolates. Inhibition of *Drechslera oryzae* in dual plate culture assay by NAIMCC-B01485 (B), NAIMCC-B01483 (C), NAIMCC-B01486 (D), NAIMCC-B01488 (E), NAIMCC-B01489 (F) and Control (A)

- Figure 41:** *In vitro* antagonistic test of foliar fungal pathogen (*D. oryzae*) against different PGPF. (B) *T. harzianum* (NAIMCC-F-03289), (C) *T. harzianum* (NAIMCC-F-03288), (D) *T. harzianum* (NAIMCC-F-03290), (E) *T. asperellum* (NAIMCC-F-03291), (F) *T. asperellum* (NAIMCC-F-03292), (G) *T. asperellum* (NAIMCC-F-03293) and (A) Control (*D. Oryzae*)
- Figure 42:** Growth promotion of rice plants following treatment with different PGPR. C-Untreated control (UI), T1-*Bacillus pumilus* (NAIMCC-B01483), T2-*Bacillus altitudinis* (NAIMCC-B01484), T3-*Bacillus altitudinis* (NAIMCC-B01485), T4-*Enterobacter cloacae* (NAIMCC-B01486), T5-*Bacillus pumilus* (NAIMCC-B01487), T6-*Bacillus pumilus* (NAIMCC-B01488), T7- *Burkholderia symbiont* (NAIMCC-B01489), T8-*Bacillus aerophilus* (NAIMCC-B01490), T9- *Paenibacillus polymyxa* (NAIMCC-B01491), T10-*Bacillus methylotrophicus* (NAIMCC-B01492).
- Figure 43:** Activity of defense enzymes in rice leaf samples (A) Peroxidase and (B) Chitinase. C-Untreated control, T1-*Bacillus pumilus* (NAIMCC-B01483), T2-*Bacillus altitudinis* (NAIMCC-B01485), T3-*Bacillus altitudinis* (NAIMCC-B01484), T4-*Enterobacter cloacae* (NAIMCC-B01486), T5-*Bacillus pumilus* (NAIMCC-B01487), T6-*Bacillus pumilus* (NAIMCC-B01488), T7- *Burkholderia symbiont* (NAIMCC-B01489), T8-*Bacillus aerophilus* (NAIMCC-B01490), T9- *Paenibacillus polymyxa* (NAIMCC-B01491), T10-*Bacillus methylotrophicus* (NAIMCC-B01492).
- Figure 44:** Activity of defense enzymes in rice leaf samples (A) PAL (Phenyl alanine ammonia lyase) and (B) β -1,3-Glucanase. C-Untreated control, T1-*Bacillus pumilus* (NAIMCC-B01483), T2-*Bacillus altitudinis* (NAIMCC-B01484), T3-*Bacillus altitudinis* (NAIMCC-B01485), T4-*Enterobacter cloacae* (NAIMCC-B01486), T5-*Bacillus pumilus* (NAIMCC-B01487), T6-*Bacillus pumilus* (NAIMCC-B01488), T7- *Burkholderia symbiont* (NAIMCC-B01489), T8-*Bacillus aerophilus* (NAIMCC-B01490), T9- *Paenibacillus polymyxa* (NAIMCC-B01491), T10-*Bacillus methylotrophicus* (NAIMCC-B01492).
- Figure 45:** HPLC profile of Phytocassanes obtained from leaf extracts of rice plant (cv. Black nuniya) showing different peaks of Phytocassanes following treatment with *Bacillus altitudinis* (NAIMCC-B01485) and pathogen challenge. (UH= Untreated Healthy, UI= Untreated Inoculated, TH= Treated Healthy and TI= Treated Inoculated)
- Figure 46:** Growth promotion of rice cultivars following treatment with different PGPF. (C= Untreated Control, T1= *T. harzianum* NAIMCC-F-03288, T2= *T. harzianum* NAIMCC-F-03289, T3= *T. harzianum* NAIMCC-F-03290, T4= *T. asperellum* NAIMCC-F-03291, T5= *T. asperellum* NAIMCC-F-03292 and T6= *T. asperellum* NAIMCC-F-03293)
- Figure 47:** Defense enzyme activity (A) Phenylalanine ammonia lyase and (B) Peroxidase of rice cultivars following treatment with PGPF and inoculated

with pathogen (*D. oryzae*). C-Untreated Control, T1-T6=Treated Inoculated [T1-*T.harzianum* (NAIMCC-F-03288), T2-*T.harzianum* (NAIMCC-F-03289), T3-*T.harzianum* (NAIMCC-F-03290), T4-*T.asperellum* (NAIMCC-F-03291), T5-*T.asperellum* (NAIMCC-F-03292) and T6-*T.asperellum* (NAIMCC-F-03293)]

Figure 48: Defense enzyme activity (A) Chitinase (B) β -1, 3-Glucanase of rice cultivars following treatment with PGPF and pathogen challenge. C- Untreated Control, T1-*T.harzianum* (NAIMCC-F-03288), T2-*T.harzianum* (NAIMCC-F-03289), T3-*T.harzianum* (NAIMCC-F-03290), T4-*T.asperellum* (NAIMCC-F-03291), T5-*T.asperellum* (NAIMCC-F-03292) and T6-*T.asperellum* (NAIMCC-F-03293)

Figure 49: HPLC analysis of Phytocassanes from leaf extracts of rice plant (cv. Black Nuniya) treated with *T. harzianum* (NAIMCC-F-03288) and pathogen challenge. (UH- Untreated Healthy, UI- Untreated Inoculated, TH- Treated Healthy and TI- Treated Inoculated)

Figure 50 : Root colonization of rice following application of AMF (*G. fasciculatum*).

Figure 51: Changes in the level of defense enzyme in rice cultivars treated with *G. fasciculatum* following artificial inoculation with *D. oryzae*.(A) PAL, (B) Peroxidase, (C) Chitinase and (D) β -1,3 Glucanase. (UH- Untreated Healthy, UI- Untreated Inoculated, TH- Treated Healthy and TI- Treated Inoculated)

Figure 52: Percent disease index in rice cultivars following treatment and artificial inoculation with *D. oryzae*. (UI- Untreated inoculated, T1- *Bacillus altitudinus* (NAIMCC-B01485) + *G. fasciculatum*, T2- *T. harzianum* (NAIMCC-F-03288)+ *G. fasciculatum*, T3= *T. harzianum* (NAIMCC-F-03288)+ *Bacillus altitudinus* (NAIMCC-B01485) and T4= *T. harzianum*(NAIMCC-F-03288)+ *G. fasciculatum* + *Bacillus altitudinus* (NAIMCC-B01485)

Figure 53: Defense enzyme activity of rice cultivars against *D. oryzae* following dual and combined application of bioinoculants. (A) PAL (Phenylalanine ammonia lyase) and (B) Peroxidase. (UI- Untreated inoculated, T1- *Bacillus altitudinus* (NAIMCC-B01485) + *G. fasciculatum*, T2- *T. harzianum* (NAIMCC-F-03288)+ *G. fasciculatum*, T3= *T. harzianum* (NAIMCC-F-03288)+ *Bacillus altitudinus* (NAIMCC-B01485) and T4= *T. harzianum*(NAIMCC-F-03288)+ *G. fasciculatum* + *Bacillus altitudinus* (NAIMCC-B01485)

Figure 54: Defense enzyme activity of rice cultivars against *D. oryzae* following dual and combined application of bioinoculants. (A)Chitinase and (B) β -1,3Glucanase. (UI- Untreated inoculated, T1- *Bacillus altitudinus* (NAIMCC-B01485) + *G. fasciculatum*, T2- *T. harzianum* (NAIMCC-F-03288)+ *G. fasciculatum*, T3= *T. harzianum* (NAIMCC-F-03288)+ *Bacillus*

altitudinus (NAIMCC-B01485) and T4= *T. harzianum*(NAIMCC-F-03288)+ *G. fasciculatum* + *Bacillus altitudinus* (NAIMCC-B01485)

Figure 55: SDS-PAGE (A) and Peroxizyme analysis (B) of leaf proteins of rice cultivar (Black nuniya) following treatment with bioinoculants and challenged inoculation with *D. oryzae*

Figure 56: Radial growth bioassay of Ethyl acetate extracts of untreated control and treated inoculated rice leaves (cv. Black Nuniya). (A) Untreated Control and(B) Treated with bioinoculants [*T. harzianum* (NAIMCC-F-03288), *R. fasciculatus* and *B. altitudinus* (NAIMCC-B01485)] and inoculation with *D. oryzae*.

Figure 57: Cellular localization of glucanase in leaf tissue of rice (cv. Black Nuniya) following combined treatment of bioinoculants and challenge inoculation with fungal pathogen, probed with PAb of glucanase and labelled with FITC. (A) Control and (B&C) Treated.

Figure 58: Cellular localization of chitinase in leaf tissue of rice (cv. Black Nuniya) following combined treatment of bioinoculants and challenge inoculation with fungal pathogen, probed with PAb of chitinase and labelled with FITC. (A) Control and (B&C) Treated.

Figure 59: Standard phenolics detected through HPLC analysis. (A) 1: Gallic acid, 2: 3,4-dihydroxybenzoic acid(DHBA) , 3: Catechol, 4: Caffeine, 5: Caffeic acid, 6: Ferulic acid, 7: Salicylic acid.(B) 1: Phloroglucinol 2: Pyrogallol 3: Catechin 4: Chlorogenic acid 5: Vanillic acid 6: Cinnamic acid.

Figure 60: HPLC analysis of phenolic acid content from leaf extracts of (cv. Black Nuniya). (A) Control and (B) Treated with bioinoculants against pathogen challenge

Figure 61: HPLC analysis of Phytocassanes from leaf extracts of (cv. Black Nuniya) treated with bioinoculants and challenge inoculation with foliar fungal pathogen (*D. oryzae*). UH= Untreated Healthy, UI= Untreated Inoculated, TH= Treated Healthy and TI= Treated Inoculated.

LIST OF APPENDICES

APPENDIX A: List of thesis related publications

APPENDIX B: List of Abbreviations

APPENDIX C: List of chemicals