

LIST OF TABLES

| Table | Page No. |
|---|----------|
| 3.1. Provenance and purity of the cosolutes used. | 84 |
| 3.2. Provenance and purity of the food additives and drugs used. | 85 |
| 4.1. Densities (ρ) and viscosities (η) of different aqueous solutions of TBAHS at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 95 |
| 4.2. Molalities (m), densities (ρ), viscosities (η), and apparent molar volumes (ϕ_V) of ascorbic acid in aqueous TBAHS solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 97-101 |
| 4.3. Partial molar volumes (ϕ_V^0) and the slopes (S_V^*) of Eq. 2 for L-ascorbic acid in different aqueous TBAHS solutions with corresponding standard deviations at different temperatures. | 102-103 |
| 4.4. Partial molar expansibilities (ϕ_E^0), its temperature dependence ($\partial\phi_E^0/\partial T$) _p and the slopes (S_E) of Eq. 5 for ascorbic acid in aqueous TBAHS solutions at $T = 298.15 \text{ to } 318.15 \text{ K}$. | 108 |
| 4.5. Viscosity B -coefficient of L-ascorbic acid with the correlation coefficients (R^2), standard deviations (σ) for linear regression of Eq. 6 along with the solvation number (S_n) in aqueous TBAHS solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 109-110 |
| 4.6. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for L-ascorbic acid in different in aqueous TBAHS solutions at different temperatures. | 113-115 |
| 5.1. Molalities (m), densities (ρ), viscosities (η), and apparent molar volumes (ϕ_V) of thiamine hydrochloride in aqueous TBAHS solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 121-125 |
| 5.2. Standard partial molar volumes (ϕ_V^0), the slopes (S_V) and corresponding standard deviations (σ) for thiamine hydrochloride in aqueous solutions of TBAHS at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 127 |
| 5.3. Standard partial molar expansibilities (ϕ_E^0) for thiamine hydrochloride in aqueous solutions of TBAHS at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 134 |
| 5.4 Viscosity B -coefficients of thiamine hydrochloride with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous TBAHS solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 136-137 |
| 5.5. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $\Delta H_2^{0\#}$ and $T\Delta S_2^{0\#}$ for thiamine hydrochloride in aqueous TBAHS solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 141-143 |

| Table | Page No. |
|---|----------|
| 6.1. Densities (ρ) and viscosities (η) of different aqueous nicotinic acid solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 148 |
| 6.2. Molalities (m), densities (ρ), viscosities (η), and apparent molar volumes (ϕ_V) of TBAHS in aqueous nicotinic acid solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 150-154 |
| 6.3. Standard partial molar volumes (ϕ_V^0), the slopes (S_V^*) and corresponding standard deviations (σ) for TBAHS in aqueous nicotinic acid solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 155 |
| 6.4. Standard partial molar expansibilities (ϕ_E^0) for TBAHS in aqueous nicotinic solutions acid at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 157 |
| 6.5. Viscosity B -coefficients of TBAHS with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) in aqueous nicotinic acid solutions at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 164 |
| 6.6. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\neq}$, $\Delta\mu_2^{0\neq}$, $\Delta H_2^{0\neq}$ and $T\Delta S_2^{0\neq}$ for TBAHS in aqueous nicotinic solutions acid at $T = (298.15 \text{ to } 318.15) \text{ K}$. | 166-168 |
| 7.1. Densities (ρ) and viscosities (η) of different aqueous solutions of TBAHS at $T = (288.15 \text{ to } 318.15) \text{ K}$. | 174 |
| 7.2. Molalities (m), densities (ρ), viscosities (η) and apparent molar volumes (ϕ_V) of pyridoxine hydrochloride in different aqueous TBAHS solutions at $T = (288.15\text{- } 318.15) \text{ K}$. | 176-181 |
| 7.3. Partial molar volumes and the experimental slopes of Eq. (2) for pyridoxine hydrochloride in different aqueous TBAHS solutions with corresponding standard deviations at $T = (288.15\text{- } 318.15) \text{ K}$. | 182-183 |
| 7.4. Limiting partial molar expansibilities for pyridoxine hydrochloride in different aqueous TBAHS solutions at $T = (288.15\text{- } 318.15) \text{ K}$. | 191 |
| 7.5. Viscosity B -coefficients of pyridoxine hydrochloride with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of TBAHS at $T = (298.15\text{- } 318.15) \text{ K}$. | 192-194 |
| 7.6. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for pyridoxine hydrochloride in different aqueous TBAHS solutions at $T = (288.15\text{- } 318.15) \text{ K}$. | 196-198 |
| 8.1. Density (ρ) and viscosity (η) for different mixtures of $[\text{P}_{4,4,4,1}][\text{CH}_3\text{SO}_4]$ and water at $T = (303.15 - 323.15) \text{ K}$. | 205 |

| Table | Page No. |
|---|----------|
| 8.2 Molalities (m), densities (ρ), viscosities (η) and apparent molar volumes (ϕ_V) of glycine and L-alanine in different aqueous $[P_{4,4,4,1}][CH_3SO_4]$ solutions at $T = 303.15$ to 323.15 K. | 207-215 |
| 8.3. Standard partial molar volumes (ϕ_V^0), the slopes (S_V^*) and corresponding standard deviations (σ) for glycine and L-alanine in aqueous solutions of $[P_{4,4,4,1}][CH_3SO_4]$ at $T = (303.15$ to $323.15)$ K. | 217-218 |
| 8.4. Standard partial molar expansibilities (ϕ_E^0) for glycine and L-alanine in aqueous solutions $[P_{4,4,4,1}][CH_3SO_4]$ at $T = (303.15$ to $323.15)$ K. | 224 |
| 8.5. Viscosity B -coefficients of glycine and L-alanine with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of $[P_{4,4,4,1}][CH_3SO_4]$ at $T = (303.15$ to $323.15)$ K. | 226-228 |
| 8.6. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\neq}$, $\Delta\mu_2^{0\neq}$, $\Delta H_2^{0\neq}$ and $T\Delta S_2^{0\neq}$ for glycine/L-alanine in aqueous solutions of $[P_{4,4,4,1}][CH_3SO_4]$ at $T = (303.15$ to $323.15)$ K. | 230-233 |
| 9.1. Densities (ρ) and viscosities (η) of different aqueous solutions of L-ascorbic acid at $T = (298.15$ to $318.15)$ K. | 239 |
| 9.2. Molalities (m), densities (ρ), viscosities (η), apparent molar volumes (ϕ_V) and apparent specific volumes ($\phi_{V,sp}$) of lactose monohydrate in aqueous solutions of L-ascorbic acid at $T = (298.15$ to $318.15)$ K. | 241-245 |
| 9.3. Standard partial molar volumes (ϕ_V^0), the slopes (S_V^*) and corresponding standard deviations (σ) for lactose monohydrate in aqueous L-ascorbic acid solutions at $T = (298.15$ to $318.15)$ K. | 247 |
| 9.4 Standard partial molar expansibilities (ϕ_E^0) for lactose monohydrate in aqueous solutions of L- ascorbic acid at $T = (298.15$ to $318.15)$ K. | 253 |
| 9.5. Viscosity B -coefficients of lactose monohydrate with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of L-ascorbic acid at $T = (298.15$ to $318.15)$ K. | 257-258 |
| 9.6. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\neq}$, $\Delta\mu_2^{0\neq}$, $\Delta H_2^{0\neq}$ and $T\Delta S_2^{0\neq}$ for lactose monohydrate in aqueous solutions of L-ascorbic acid at $T = (298.15$ to $318.15)$ K. | 260-262 |
| 10.1. Density (ρ) and viscosity (η) for different mixtures of Na-Malonate and water at $T = (298.15 - 318.15)$ K. | 268 |

| Table | Page No. |
|--|----------|
| 10.2. Molalities (m), densities (ρ), viscosities (η) and apparent molar volumes (ϕ_v) of paracetamol in different aqueous sodium malonate solutions at $T = 298.15$ to 318.15 K. | 270-274 |
| 10.3. Partial molar volumes and the experimental slopes of Eq. (2) for paracetamol in different aqueous sodium malonate solutions with corresponding standard deviations at $T = (298.15- 318.15)$ K. | 275-276 |
| 10.4. Limiting partial molar expansibilities for paracetamol in different aqueous sodium malonate solutions at $T = (298.15- 318.15)$ K. | 283 |
| 10.5 Viscosity B -coefficients of paracetamol with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of sodium malonate at $T = (298.15- 318.15)$ K. | 284-285 |
| 10.6. Refractive indices (n_D), molar refractivities (R_M), apparent molar refractivities (R_D) and molar polarizabilities (α_p) as a function of molarities (m) of paracetamol in sodium malonate solution. | 287-289 |
| 10.7. Values of $\phi_{v,2}^0 - \phi_{v,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for paracetamol in different aqueous sodium malonate solutions at $T = (298.15- 318.15)$ K. | 291-293 |
| 11.1. Density (ρ) and viscosity (η) for different mixtures of uracil and water at $T = (298.15 - 318.15)$ K. | 300 |
| 11.2. Molalities (m), densities (ρ), viscosities (η) and apparent molar volumes (ϕ_v) of betaine hydrochloride in different aqueous uracil solutions at $T = 298.15$ to 318.15 K. | 301-305 |
| 11.3. Partial molar volumes and the experimental slopes of Eq. (2) for betaine hydrochloride in different aqueous uracil solutions with corresponding standard deviations at $T = (298.15-318.15)$ K. | 306-307 |
| 11.4. Limiting partial molar expansibilities for betaine hydrochloride in different aqueous uracil solutions at $T = (298.15-318.15)$ K. | 312 |
| 11.5. Viscosity B -coefficients of betaine hydrochloride with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of uracil at $T = (298.15$ to $318.15)$ K . | 314-315 |
| 11.6. Values of $\phi_{v,2}^0 - \phi_{v,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for betaine hydrochloride in different aqueous uracil solutions at $T = (298.15-318.15)$ K. | 318-320 |
| 12.1. Density (ρ), viscosity (η) and speed of sound (u) for different mixtures of D-glucose and water at $T = (298.15 - 318.15)$ K. | 326 |

| Table | Page No. |
|--|----------|
| 12.2. Molalities (m), densities (ρ), viscosities (η) and apparent molar volumes (ϕ_V) of metformin hydrochloride in different aqueous D-glucose solutions at $T = 298.15$ to 318.15 K. | 328-332 |
| 12.3 Partial molar volumes and the experimental slopes of Eq. (2) for metformin hydrochloride in different aqueous D-glucose solutions with corresponding standard deviations at $T = (298.15-318.15)$ K. | 333-334 |
| 12.4. Limiting partial molar expansibilities for metformin hydrochloride in different aqueous D-glucose solutions at $T = (298.15-318.15)$ K. | 338 |
| 12.5. Viscosity B -coefficients of metformin hydrochloride with the correlation coefficients R^2 , standard deviations σ for linear regression of Eq. (7) along with the solvation number S_n in aqueous solutions of D-glucose at $T = (298.15$ to $318.15)$ K. | 340-341 |
| 12.6. Molalities (m), speed of sound (u), adiabatic compressibility (β_s) and apparent molar adiabatic compression (ϕ_k), limiting apparent molar adiabatic compressibility and experimental slopes (S_K^*) of metformin hydrochloride in different aqueous D-glucose solutions at $T = 298.15$ to 318.15 K. | 343-350 |
| 12.7. Values of $\phi_{V,2}^0 - \phi_{V,1}^0$, $\Delta\mu_1^{0\#}$, $\Delta\mu_2^{0\#}$, $T\Delta S_2^{0\#}$, $\Delta H_2^{0\#}$ for metformin hydrochloride in different aqueous D-glucose solutions at $T = (298.15-318.15)$ K. | 353-355 |

LIST OF FIGURES

| Figure | Page No. |
|--|----------|
| 1.1. Schematic representation of preferential solvation, A: binary mixture of solvent 1 and solvent 2; B: ideal solvation of the solute; C: preferential solvation of the solute by solvent 1. | 9 |
| 1.2. Dissolution of an ionic crystal in a solvent. | 11 |
| 1.3. Structures of used drugs as solutes. | 19 |
| 1.4. Structures of used food additives as solutes and cosolutes. | 20 |
| 2.1. Dependence of the total volume of a solution on the volume of the pure solvent and the apparent molar volume (ϕ_V) of the solute. | 35 |
| 2.2. Passage of a molecule from one equilibrium position to another position in the same layer of a liquid with the creation of a suitable hole. | 59 |
| 2.3. Potential energy barrier for viscous flow with and without shearing force. | 59 |
| 2.4. One-dimensional Gibbs energy diagram for an equilibrium reaction, $A \leftrightarrow B$ in the solvents I and II. | 67 |
| 2.5. Three tautomeric forms of 1,3-dicarbonyl compounds. | 68 |
| 2.6. Reactants A and B react through the activated complex $(AB)^\ddagger$ to products C and D. | 70 |
| 2.7. One-dimensional Gibbs energy diagram for the reaction $A+B \rightleftharpoons C+D$ through the activated complex $(AB)^\ddagger$ in solution. | 70 |
| 2.8. One-dimensional Gibbs energy diagram for a chemical reaction in three different solvents I, II and III. | 71 |
| 3.1. Schematic representation of the water distillation set. | 83 |
| 3.2. Digital electronic analytical balance (Mettler Toledo, AG 285). | 86 |
| 3.3. Anton Paar densitometer (DMA 4500M). | 87 |
| 3.4. Capillary viscometers: A suspended-level Ubbelohde viscometer. Inset: Cannon type Ubbelohde viscometer. | 88 |
| 3.5. Abbe Refractometer (Cyberlab, MA01527, USA). | 89 |
| 3.6. Jasco V-530 double beam UV-VIS Spectrophotometer. | 90 |
| 3.7. Multifrequency ultrasonic interferometer. | 91 |
| 3.8. Systronics digital pH meter. | 92 |

| Figure | Page No. |
|--|----------|
| 4.1. Molecular structure of: a) L-Ascorbic acid and b) Tetrabutylammonium hydrogen sulfate. | 95 |
| 4.2. Dependence of standard partial molar volumes (ϕ_V^0) for L-ascorbic acid on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 104 |
| 4.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for L-ascorbic acid on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 106 |
| 4.4. Plots of viscosity B -coefficients of transfer ($\Delta_t B$) for L-ascorbic acid on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 111 |
| 5.1. Molecular structure of (a) thiamine hydrochloride and (b) TBAHS. | 120 |
| 5.2. Dependence of standard partial molar volumes (ϕ_V^0) for thiamine hydrochloride on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 128 |
| 5.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for thiamine hydrochloride on the molarity of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 130 |
| 5.4. Different possible ionic group interactions in aqueous solution of thiamine hydrochloride and TBAHS. | 131 |
| 5.5. Changes in the UV-Visible absorption spectra of thiamine hydrochloride in aqueous TBAHS ($1 \cdot 10^{-4} \text{ mol} \cdot \text{L}^{-1}$) solution at 298.15 K with the following molarities of thiamine hydrochloride ($10^{-4} \text{ mol} \cdot \text{L}^{-1}$): 1, 0.000; 2, 0.009; 3, 0.019; 4, 0.029; 5, 0.039; 6, 0.049; 7, 0.059; 8, 0.069; 9, 0.079; 10, 0.089; 11, 0.099; 12, 0.109; 13, 0.118; 14, 0.128; 15, 0.138; 16, 0.148. | 132 |
| 5.6. Changes in the UV-Visible absorption spectra of thiamine hydrochloride ($1 \cdot 10^{-4} \text{ mol} \cdot \text{L}^{-1}$) in different aqueous TBAHS solution at 298.15 K with the following molarities of TBAHS ($10^{-5} \text{ mol} \cdot \text{L}^{-1}$): 1, 0.000; 2, 0.099; 3, 0.196; 4, 0.291; 5, 0.385; 6, 0.476; 7, 0.566; 8, 0.654; 9, 0.740; 10, 0.826; 11, 0.909; 12, 0.991; 13, 1.071; 14, 1.150; 15, 1.228; 16, 1.304; 17, 1.379; 18, 1.453; 19, 1.525; 20, 1.597; 21, 1.667; 22, 1.735; 23, 1.803; 24, 1.870; 25, 1.935. Inset: 1, thiamine hydrochloride ($1 \cdot 10^{-4} \text{ mol} \cdot \text{L}^{-1}$) in water; 2: average of absorption spectra (1 to 25) at 298.15 K. | 133 |
| 5.7. Plots of viscosity B -coefficients of transfer ($\Delta_t B$) for thiamine hydrochloride on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 139 |

| Figure | Page No. |
|---|----------|
| 6.1. Molecular structure of (a) nicotinic acid and (b) TBAHS. | 148 |
| 6.2. Dependence of standard partial molar volumes (ϕ_V^0) for TBAHS on the molarity of nicotinic acid (c_{NA}) in aqueous solutions at $T = (298.15-318.15)$ K. | 156 |
| 6.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for TBAHS on the molarity of nicotinic acid (c_{NA}) in aqueous solutions at $T = (298.15-318.15)$ K. | 159 |
| 6.4. Different possible structural forms of nicotinic acid in aqueous solution: a positively charged form (AH_2^+), two of isoelectric forms (AH^+ and AH^0) and a negatively charged form (A^-). | 160 |
| 6.5. Changes in the UV-Visible absorption spectra of 4.0×10^{-5} mol.dm ⁻³ nicotinic acid in ternary solutions (TBAHS + nicotinic acid + H ₂ O) with the following molarities of TBAHS: 1, 2.0×10^{-5} mol.dm ⁻³ ; 2, 4.0×10^{-5} mol.dm ⁻³ ; 3, 6.0×10^{-5} mol dm ⁻³ ; 4, 8.0×10^{-5} mol.dm ⁻³ . | 161 |
| 6.6. Changes in the UV-Visible absorption spectra of nicotinic acid (4.0×10^{-5} mol.dm ⁻³) in ternary solutions (TBAHS+ nicotinic acid + H ₂ O) with the following molarities of nicotinic acid: 1, 2.0×10^{-5} mol.dm ⁻³ ; 2, 4.0×10^{-5} mol .dm ⁻³ ; 3, 6.0×10^{-5} mol.dm ⁻³ ; 4, 8.0×10^{-5} mol.dm ⁻³ . | 162 |
| 6.7. Different possible ionic group interactions in aqueous solution of TBAHS and nicotinic acid. | 163 |
| 7.1. Molecular structure of (a) pyridoxine hydrochloride and (b) TBAHS. | 173 |
| 7.2. Dependence of standard partial molar volumes (ϕ_V^0) for pyridoxine hydrochloride on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (288.15-318.15)$ K. | 184 |
| 7.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for pyridoxine hydrochloride on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (288.15-318.15)$ K. | 186 |
| 7.4. Changes in the UV-Visible spectra of TBAHS ($1. \cdot 10^{-4}$ mol.L ⁻¹) in different aqueous Pyridoxine hydrochloride solutions at 298.15 K with the following molarities of pyridoxine hydrochloride (10^{-4} mol.L ⁻¹): 1, 0.000; 2, 0.0198; 3, 0.0396; 4, 0.0594; 5, 0.0792; 6, 0.0990; 7, 0.1002 at 298.15 K. | 188 |
| 7.5. Changes in the UV-Visible spectra of pyridoxine hydrochloride ($1. \cdot 10^{-4}$ mol.L ⁻¹) in different aqueous TBAHS solutions at 298.15 K with the following molarities of TBAHS (10^{-4} mol.L ⁻¹): 1, 0.000; 2, 0.0198; 3, 0.0396; 4, 0.0594; 5, 0.0792; 6, 0.0990; 7, 0.1002 at 298.15 K. | 189 |
| 7.6. Plots of viscosity B -coefficients of transfer ($\Delta_t B$) for pyridoxine hydrochloride on the molality of TBAHS (m_{TBAHS}) in aqueous solutions at $T = (298.15-318.15)$ K. | 195 |

| Figure | Page No. |
|---|----------|
| 8.1. Molecular structure of: a) Glycine and b) L-alanine and c) $[P_{4,4,4,1}][CH_3SO_4]$. | 205 |
| 8.2. Dependence of standard partial molar volumes (ϕ_V^0) for (a) glycine and (b) L-alanine on the molality of $[P_{4,4,4,1}][CH_3SO_4]$ (m_{IL}) in aqueous solutions at $T = (298.15-318.15)$ K. | 219 |
| 8.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for (a) glycine and (b) L- alanine on the molality of $[P_{4,4,4,1}][CH_3SO_4]$ (m_{IL}) in aqueous solutions at $T = (298.15-318.15)$ K. | 221 |
| 9.1. Molecular structures of (a) lactose monohydrate and (b) L-ascorbic acid. | 239 |
| 9.2. Dependence of standard partial molar volumes (ϕ_V^0) for lactose monohydrate on the molality (m) of l-ascorbic acid in aqueous solutions at $T = (298.15-318.15)$ K. | 248 |
| 9.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for lactose monohydrate on the molality of L-ascorbic acid in aqueous solutions at $T = (298.15-318.15)$ K. | 250 |
| 9.4. Changes in the UV-Visible absorption spectra of lactose monohydrate ($1. 10^{-4}$ mol.L ⁻¹) in different aqueous L-ascorbic acid solutions at 298.15K with the following molarities of L-ascorbic acid (10^{-4} mol.L ⁻¹): 1, 0.000; 2, 0.0099; 3, 0.0196; 4, 0.0291; 5, 0.0385; 6, 0.0476; 7,0.0566; 8, 0.0654; 9, 0.0740; 10, 0.0826; 11, 0.0909; 12, 0.0991; 13, 0.1071. | 254 |
| 9.5 Changes in the UV-Visible absorption spectra of lactose monohydrate in aqueous L-ascorbic acid($1. 10^{-4}$ mol.L ⁻¹) at 298.15 K with the following molarities of lactose monohydrate (10^{-4} mol.L ⁻¹): 1, 0.000; 2, 0.0099; 3, 0.0196; 4, 0.0291; 5, 0.0385; 6, 0.0476; 7,0.0566; 8, 0.0654; 9, 0.0740; 10, 0.0826; 11, 0.0909; 12, 0.0991; 13, 0.1071 Inset: 1, ascorbic acid in water (5.10^{-5} mol.L ⁻¹); 2, average of absorption spectra (1to 13) at 298.15K. | 255 |
| 9.6. Plots of viscosity B -coefficients of transfer (Δ_tB)for lactose monohydrate on the molality of L-ascorbic acid in aqueous solutions at $T = (298.15-318.15)$ K. | 259 |
| 10.1. Molecular structure of (a) paracetamol and (b) sodium malonate. | 267 |
| 10.2. Dependence of standard partial molar volumes (ϕ_V^0) for paracetamol on the molality of sodium malonate (m_{Na_2Mal}) in aqueous solutions at $T = (298.15-318.15)$ K. | 277 |
| 10.3. Plots of standard partial molar volume of transfer ($\Delta_t\phi_V^0$) for paracetamol on the molality of sodium malonate (m_{Na_2Mal}) in aqueous solutions at $T = (298.15-318.15)$ K. | 278 |

| Figure | Page No. |
|--|----------|
| <p>10.4. Changes in the UV-Visible absorption spectra of paracetamol in aqueous sodium malonate ($1 \cdot 10^{-4} \text{ mol.L}^{-1}$) at 298.15 K with the following molarities of paracetamol ($10^{-4} \text{ mol.L}^{-1}$): 1, 0.000; 2, 0.0099; 3, 0.0196; 4, 0.0291; 5, 0.0385; 6, 0.0476; 7, 0.0566; 8, 0.0654; 9, 0.0740; 10, 0.0826; 11, 0.0909; 12, 0.0991; 13, 0.1071; 14, 0.150; 15, 0.1228; 16, 0.1304; 17, 0.138; 18, 0.145; 19, 0.1525; 20, 0.160; 21, 0.167; 22, 0.173; 23, 0.180; 24, 0.187; 25, 0.193; 26, 0.199.</p> | 280 |
| <p>10.5. Changes in the UV-Visible absorption spectra of paracetamol ($5 \cdot 10^{-5} \text{ mol.L}^{-1}$) in different aqueous sodium malonate solutions at 298.15 K with the following molarities of sodium malonate ($10^{-4} \text{ mol.L}^{-1}$): 1, 0.000; 2, 0.099; 3, 0.0196; 4, 0.0291; 5, 0.0385; 6, 0.0476; 7, 0.0566; 8, 0.0654; 9, 0.0740; 10, 0.0826; 11, 0.0909; 12, 0.0991; 13, 0.1071; 14, 0.150; 15, 0.1228; 16, 0.1304. Inset: 1, paracetamol in water ($5 \cdot 10^{-5} \text{ mol.L}^{-1}$); 2, average of absorption spectra (1 to 16) at 298.15 K.</p> | 281 |
| <p>10.6. Plots of viscosity B-coefficients of transfer ($\Delta_t B$) for paracetamol on the molality of sodium malonate ($m_{\text{Na}_2\text{Mal}}$) in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 286 |
| <p>11.1. Molecular structures of (a) betaine hydrochloride and (b) uracil.</p> | 300 |
| <p>11.2. Dependence of standard partial molar volumes (ϕ_V^0) for Betaine hydrochloride on the molality of uracil (m_{Uracil}) in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 309 |
| <p>11.3. Plots of standard partial molar volume of transfer ($\Delta_t \phi_V^0$) for Betaine hydrochloride on the molality of uracil in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 310 |
| <p>11.4. Plots of viscosity B-coefficients of transfer ($\Delta_t B$) for Betaine hydrochloride on the molality of uracil in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 316 |
| <p>12.1 Molecular structure of (a) metformin hydrochloride and (b) D-glucose.</p> | 325 |
| <p>12.2. Dependence of standard partial molar volumes (ϕ_V^0) for Metformin hydrochloride on the molality of D-glucose (m_{Glucose}) in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 335 |
| <p>12.3. Plots of standard partial molar volume of transfer ($\Delta_t \phi_V^0$) for Metformin hydrochloride on the molality of D-glucose in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 336 |
| <p>12.4. Dependence of limiting apparent molar adiabatic compressibility (ϕ_k^0) for Metformin hydrochloride on the molality of D-glucose (m_{Glucose}) in aqueous solutions at $T = (298.15\text{-}318.15) \text{ K}$.</p> | 342 |

12.5. Plots of partial molar adiabatic compression of transfer ($\Delta_t \phi_k^0$) for Metformin hydrochloride on the molality of D-glucose in aqueous solutions at $T = (298.15-318.15)$ K.

351

LIST OF APPENDICES

| Appendix | Page No. |
|--|-----------------|
| I: List of Publications | 365 |
| II: List of Communicated Articles | 366 |
| III: List of Other Publications | 367 |
| IV: Seminar, Symposium, Convention Attended | 368 |