

# LIST OF TABLES

## Chapter 1

**Table 1.1.** Hydrodynamic diameter and zeta potential data for thermally treated colloidal silica NPs. 31

## Chapter 3

**Table 3.1** Transmission Electron Microscopic (TEM) and DLS Results of Colloidal Zinc Chromate prepared in H<sub>2</sub>O/AOT/n-heptane W/O Microemulsion (μE). 72

## Chapter 4

**Table 4.1** Elemental contribution to the nanocomposite from EDS analysis. 90

## Chapter 5

**Table 5.1** Calculated molar absorption coefficients ( $\epsilon_{abs}$ ), molar scattering coefficient( $\epsilon_{sca}$ ), Surface area to volume ratio ( $S/V$ ) and band gaps of NPs of varying shapes (sphere, cylinder, cube) and sizes at their corresponding surface plasmon resonance wavelength maximum ( $\lambda_{max}$ ) 101

**Table 5.2** Discrete electronic excitations of Cu<sub>2</sub>O clusters of different sizes in  $C_{2v}$  symmetry calculated through TDDFT. 113

**Table 5.3** TDDFT data of discrete electronic excitations of Cu<sub>2</sub>O clusters of same sizes in different symmetries. 116

**Table 5.4** TDDFT data of discrete electronic excitations of larger Cu<sub>2</sub>O clusters of different sizes in same symmetry. 118

## Chapter 6

**Table 6.1** Discrete electronic excitation of 11-*cis*-retinal and its gold conjugates with different symmetries calculated using TDDFT. 131

**Table 6.2.** TDDFT data most intense excitation wavelengths of 11-*cis*-retinal, gold clusters of different symmetries and corresponding retinal-gold conjugates with involved transitions and their dipole contribution to transition dipole moments. 136

**Table 6.3.** Thermochemical data of 11-*cis*-retinal and its gold conjugate calculated from vibrational frequency analysis. 137

# LIST OF FIGURES AND SCHEMES

## Chapter 1

- Figure 1.1.**  $\chi(n)$  vs.  $n^{-\beta}$  Plots where  $\beta = 0$ . 3
- Figure 1.2.** Calculated cohesive energies of various size magnesium clusters in their most stable geometry as function of  $N^{-1/3}$ . 4
- Figure 1.3.**  $I-U$  characteristic of ideal single electron transport, where Coulomb blocked is shown as the step function. 7
- Figure 1.4. (a)** Absorption spectra of colloidal metal NPs with different size **(b)** Luminescent colloidal semiconductor NPs. 8
- Figure 1.5.** Magnetic behavior under an applied field. 9
- Figure 1.6.** Transition from superparamagnetic to single to multi-domain regimes. 9
- Figure 1.7.** Mechanistic steps of nanoparticle formation. 15
- Figure 1.8.** Surfactants (long chain) and cosurfactants (short chain alkanols) are essential for stable microemulsion. 16
- Figure 1.9.** Schematic representations of micelle and reverse micelles. 17
- Figure 1.10.** Schematic representation of reverse and normal microemulsions. 17
- Figure 1.11.** Various methods to synthesis different NPs 18
- Figure 1.12.** Schematic representation of steps of formation of NPs in reverse micelle templates. 21
- Figure 1.13.** Growth of AgBr NPs in CTAB colloidal dispersion. 22
- Figure 1.14.** Absorption spectra of surfactant stabilized silver chloride NPs. 23
- Figure 1.15. (a)** Absorption and **(b)** emission spectra of 7-HC in presence of Ludox<sup>R</sup>. 23
- Figure 1.16.** FTIR spectra of Fe<sub>3</sub>O<sub>2</sub> NPs. 24

<b>Figure 1.17.</b> XRD patterns of Co and Fe <sub>3</sub> O <sub>4</sub> nanoparticles.	25
<b>Figure 1.18.</b> Raman spectra of surfactant stabilized AgCl NPs in aqueous media.	25
<b>Figure 1.19.</b> SEM image of monodispersed superparamagnetic single-crystal magnetite NPs with mesoporous structure.	26
<b>Figure 1.20.</b> Transmission electron microscopy image of Fe <sub>3</sub> O <sub>4</sub> NPs.	27
<b>Figure 1.21.</b> HRTEM image showing fringes in the core and shell of a particle.	27
<b>Figure 1.22.</b> AFM image of the transferred films of Au nanoparticles-PMMA hybrids.	28
<b>Figure 1.23.</b> EDX spectrum Nanoparticle grafted on Dendritic $\alpha$ -Fe <sub>2</sub> O <sub>3</sub> .	29
<b>Figure 1.24.</b> The size distribution of recoil-implanted Au NPs.	30
<b>Figure 1.25.</b> Schematic representation of electric double layer formation around NPs.	30

## Chapter 2

<b>Figure 2.1 (a)</b> Otto configuration <b>(b)</b> Kretschmann configuration.	42
<b>Figure 2.2</b> Nanoparticles of various shape and size in solution – the plasmonic resonance determines the colour.	43
<b>Figure 2.3</b> Real and imaginary part of dielectric constant for gold according to the Drude-Sommerfeld free electron model.	44
<b>Figure 2.4</b> Quasi-static model for particle plasmons.	49
<b>Figure 2.5</b> Prolate (left, $a > b = c$ ) and oblate (right, $a < b = c$ ) ellipsoids.	51
<b>Figure 2.6</b> Extinction, absorption and scattering efficiencies ( $Q$ ) of spherical silver NPs (in vacuum) with increasing diameter ( $D$ ).	54
<b>Figure 2.7</b> (left) Change in valence electron density using quantum fluid dynamics, where dark shading indicates a density decrease for the lowest excitation in Na <sub>2</sub> . (right) Integrated electron density along the bond axis using TDDFT for the lowest excitation in Na <sub>2</sub> .	55

## Chapter3

**Figure 3.1. A.** X-ray diffraction pattern of zinc chromate particles prepared in water/AOT/n-heptane microemulsion. 65

**Figure 3.1.B.** FTIR spectra of  $K_2CrO_4$  (A);  $ZnCrO_4$  prepared in bulk water (B) and in water/AOT/n-heptane W/O microemulsion at  $\omega = 20$  (C); 15(D); 10 (E) and 5 (F). 66

**Figure 3.2** Absorption spectra of colloidal  $ZnCrO_4$  in water/AOT/heptane W/O microemulsion at 303K at  $\omega = 15$ . Inset: absorption spectra of  $K_2CrO_4$  at  $\omega = 15$  67

**Figure 3.3** Absorption spectra of colloidal zinc chromate prepared in  $H_2O/AOT/n$ -heptane W/O microemulsion at different  $\omega$  values at 303K. Inset: absorption spectra of  $K_2CrO_4$  ( $1.0 \times 10^{-4} \text{ mol dm}^{-3}$ ) in the same medium at various  $\omega$  values. 68

**Figure 3.4** Variation of absorbance with  $\omega$  for the nanocolloidal dispersion of  $ZnCrO_4$  prepared in water/AOT/n-heptane W/O microemulsion. 69

**Figure 3.5** Plot of  $(\epsilon h\nu)^2$  against  $h\nu$  at fixed concentration of  $ZnCrO_4$  with various  $\omega$  values. 70

**Figure 3.6.** Profile of  $d_h/nm$  vs.  $\omega$ . Inset: (A) Variation of  $d_h$  with the concentration of  $ZnCrO_4$  in microemulsion medium at  $\omega= 5$ . (B)  $d_h(nm)$  profile for a  $0.3 \text{ mmol dm}^{-3}$   $ZnCrO_4$  in microemulsion medium. 71

**Figure 3.7** Plot of  $H_f$  vs.  $\omega$  for the formation of  $ZnCrO_4$  in  $H_2O/AOT/heptane$  W/O microemulsion at 303 K. 73

**Figure 3.8** Transmission electron micrograph of colloidal  $ZnCrO_4$  dispersions prepared in water/AOT/n-heptane W/O microemulsion. 75

**Figure 3.9** Transmission electron micrograph of colloidal  $ZnCrO_4$  dispersions prepared in water/AOT/n-heptane W/O microemulsion. 77

## Chapter 4

**Figure 4.1** X-ray diffraction pattern of  $Fe_3O_4/Co$  nanocomposite. 87

**Figure 4.2** Morphology of nanocomposites at increasing concentration from (a) to (c) 87

**Figure 4.3** HRTEM images of nanocomposites showing the fringes of the Fe<sub>3</sub>O<sub>4</sub>, Co and CoFe structures. Image of the SAED experiment also shows the reflections of the corresponding planes. 88

**Figure 4.4** Image of the spectral lines obtained from EDS analysis. 89

**Figure 4.5 (a)** Concentration vs size and PDI profile of nanocomposite dispersed in water.  
**(b)** Time vs size and PDI profile of nanocomposite dispersed in water 91

**Figure 4.6 (a)** The absorption spectra of Fe<sub>3</sub>O<sub>4</sub>/Co nanocomposite formed at different precursor concentrations. **(b)** The absorption spectra of Fe<sub>3</sub>O<sub>4</sub> NP formed at different precursor concentrations. 92

## Chapter 5

**Figure 5.1** Simulated spectra of (a) absorption (b) scattering and (c) extinction of Cu<sub>2</sub>O nanospheres with diameter 100 nm to 240 nm 100

**Figure 5.2** Variation of calculated band gaps of spheres (red), cylinders (green) and cubes (blue) with increasing size parameters. 102

**Figure 5.3** Simulated spectra of (a) absorption (b) scattering (c) extinction of Cu<sub>2</sub>O nanocylinders with size parameters long axis : diameter from 553 nm : 30 nm to 1185 nm : 80 nm 103

**Figure 5.4** Variation of absorption with shapes [sphere (red), cylinder (green), cube (black)] of NPs having equal volume of a sphere with effective radius 100 nm. 104

**Figure 5.5** Variation of scattering with shapes [sphere(red), cylinder(green), cube(black)] of NPs having equal volume of sphere with effective radius 100 nm. 105

**Figure 5.6** Simulated spectra of (a) absorption (b) scattering (c) extinction of Cu<sub>2</sub>O nanocubes having equal volume of nanospheres of diameter 100 nm to 240 nm. 106

**Figure 5.7** Molecular clusters of Cu<sub>2</sub>O (a) (Cu<sub>2</sub>O)<sub>1</sub>, (b) (Cu<sub>2</sub>O)<sub>2</sub>, (d) (Cu<sub>2</sub>O)<sub>3</sub> in C<sub>2v</sub> symmetry, (c) and (e) are (Cu<sub>2</sub>O)<sub>3</sub> in C<sub>s</sub> and C<sub>3v</sub> symmetry respectively. 109

**Figure 5.8 (a).** TDDFT valence excitation spectra (oscillator strength vs excitation energy) of (Cu<sub>2</sub>O)<sub>1</sub> (black), (Cu<sub>2</sub>O)<sub>2</sub> (red) and (Cu<sub>2</sub>O)<sub>3</sub> (green) clusters having same symmetry (C<sub>2v</sub>). **(b).** The representation of MOs participating in the electronic transitions. 110

**Figure 5.9** Variation of extent of LMCT in terms of % contribution of (a) ligand AOs in occupied MOs and (b) metal AOs in virtual MOs in molecular clusters of  $(\text{Cu}_2\text{O})_1$ (black),  $(\text{Cu}_2\text{O})_2$  (red) and  $(\text{Cu}_2\text{O})_3$  (green). 112

**Figure 5.10 (a)** TDDFT valence excitation spectra (oscillator strength vs excitation energy) of  $(\text{Cu}_2\text{O})_3$  clusters having different symmetries  $C_{2v}$  (red) and  $D_{3h}$  (black). The MOs involved in corresponding transitions are also displayed. **(b)**. The representation of MOs participating in the electronic transitions. 115

**Figure 5.11** Molecular clusters of (a)  $[\text{Cu}_{28}\text{O}_{15}]^{6+}$  and (b)  $[\text{Cu}_{44}\text{O}_{30}]^{6+}$  in  $T_d$  symmetry. 118

**Figure 5.12** TDDFT valence excitation spectra (oscillator strength vs excitation energy) of (a)  $[\text{Cu}_{28}\text{O}_{15}]^{6+}$  and (b)  $[\text{Cu}_{44}\text{O}_{30}]^{6+}$  in  $T_d$  symmetry. 119

## (a) Chapter 6

**Figure 6.1** Photoisomerization of 11-*cis*-retinal to *all-trans* retinal by photo-absorption at 500 nm wavelength. 124

**Figure 6.2. (a)** TDDFT valence excitation spectra (oscillator strength vs excitation energy) of 11-*cis*-retinal molecule. **(b)** The representation of MOs participating in  $\pi$ - $\pi^*$  transition leading to photoisomerization of 11-*cis*-retinal to *all-trans* retinal at 500 nm. 128

**Figure 6.3.** Optimized  $\text{Au}_{14}$  clusters with **(a)**  $O_h$ , **(b)**  $D_{2h}$  and **(c)**  $C_{4v}$  symmetries respectively. 129

**Figure 6.4.** The most intense excitations ( $\lambda_{\text{max}}$ ) for  $\text{Au}_{14}$  clusters having  $O_h$ ,  $D_{2h}$  and  $C_{4v}$  symmetries, are depicted in black, green and red colors respectively. 130

**Figure 6.5.** (a) UV-visible excitation spectra with most intense excitations ( $\lambda_{\text{max}}$ ) for retinal-cluster conjugates with symmetries  $O_h$ ,  $D_{2h}$  and  $C_{4v}$ . (b) The source and destination MOs responsible for the peaks. 133

**Figure 6.6.** The source and destination MOs obtained from the UV-visible excitation energy calculation of the conjugates using CAM-B3LYP long range corrected functional. 134

**Figure 6.7.** An apparent isomerization is displayed as the *cis*-retinal is transformed into *trans*-form in the estimation of the excited states corresponding to the most intense excitations for (a) retinal-gold ( $O_h$ ) and (b) retinal-gold ( $D_{2h}$ ) conjugates. 135

**Figure 6.8.** The symmetric ethylenic stretching frequency of  $\text{C}_{11}=\text{C}_{12}$  double bond. 138

**Figure 6.9.** (A) Frequency of the torsional motion of H–C<sub>11</sub> and H–C<sub>12</sub> bond in the HOOP region of C<sub>11</sub>=C<sub>12</sub> bond. For better clarification the regions spotted by (B), (C), and (D) are amplified individually.

139

## Schemes

### Chapter 1

**Scheme 1.1.** Two new surfaces being created by breaking a rectangular solid into two pieces. Multiplication of atoms ( $10^7$ ) times in nanocrystal as compared to the surface atoms in bulk. 5

**Scheme 1.2.** (A) Density of states in one band of a semiconductor as a function of dimension. (B) Schematic illustration of the density of states in metal and semiconductor clusters. 6

**Scheme 1.3** Typical synthetic methods for nanoparticles for the top-down and bottom-up approaches. 12

**Scheme 1.4.** Schematic representation of (a) epitaxial route (pulsed laser deposition) and (b) colloidal route (one microemulsion method) of synthesis of NPs. 13

**Scheme 1.5.** Schematic representation of the LSPR of NPs with the shape of a sphere (A) and a triangular prism (B). 34

### Chapter 4

**Scheme 4.1** Synthetic steps of Fe<sub>3</sub>O<sub>4</sub>/Co nanocomposite in micellar medium. 85

### Chapter 6

**Scheme 6.1.** Schematic representation of the primary steps in the mechanism of vision. 127



