

LIST OF FIGURES

- Fig. 1.1** (a) Cultivated bottle gourd in farmer's field; (b) Diseased bottle gourd fruits in farmer's field; (c) Diseased samples showing fungal infection in bottle gourd fruits
- Fig. 3.1** Map showing different areas of collection of diseased bottle gourd plant samples in sub-Himalayan West Bengal region
- Fig. 3.2** Regions of collection of infected bottle gourd plant parts from different districts of North Bengal. The green cones showing the collection spots. (a) Khoribari, Darjeeling; (b) Mohitnagar, Jalpaiguri; (c) Naxalbari, Darjeeling; (d) Kamakhyaguri, Alipurduar; (e) Dinhata, Coochbehar; (f) Islampur, Uttar Dinajpur
- Fig. 4.1** Naturally infected bottle gourd plants in the cultivation fields of different areas; (a) from Khoribari, Darjeeling; (b) from Mohitnagar, Jalpaiguri; (c) from Naxalbari, Darjeeling; (d) from Kamakhyaguri, Alipurduar; (e) from Dinhata, Coochbehar; (f) from Islampur, Uttar Dinajpur
- Fig. 4.2** Fungal infection within and outside of the fruits. Bottle gourd fruit rot caused by *Fusarium incarnatum* (a-d) and *Colletotrichum gloeosporioides* (e and f)
- Fig. 4.3** (a-f) Infected bottle gourd fruits in different farmers' fields of the present study area caused either by *Fusarium incarnatum* or by *Colletotrichum gloeosporioides* or by both
- Fig 4.4** Bottle gourd fruit in glass chambers (a) control without scratch; (b) control with scratches done by sterile needle; (c) scratched and inoculated by spore suspension of a pathogen soaked in sterile absorbent cotton wool
- Fig 4.5** Fruits inoculated by *Colletotrichum gloeosporioides*; (a) Control (Gadda-1) after 10 days of inoculation; (b) coalescence started and brown patch found to appear after 10 days of inoculation; (c) control (local variety) after 10 days of inoculation; (d) Deep brown lines found to occur after 10 days of inoculation
- Fig. 4.6** Fruits inoculated by *Fusarium incarnatum*; (a) Control (Gadda-1) after 10 days of inoculation; (b) Brown lines started coalescence after 10 days of inoculation; (c) control (local

variety) after 10 days of inoculation; (d) Deep brown lines found after 10 days of inoculation

- Fig. 4.7** Whole plant inoculation of *Lagenaria siceraria*. (a) Healthy Gadda-1 plant sprayed with sterile distilled water; (b) Healthy Local variety sprayed with sterile distilled water; (c) Infected plant following inoculation of *Colletotrichum gloeosporioides*. Disease symptoms are visible in the margin of leaves; (d) Local variety infected with *Fusarium incarnatum*. Brown necrotic symptoms visible within white mycelial infestation
- Fig. 4.8** Pure culture of both of the pathogenic isolates in PDA plates and slant. (a) and (b) 7 day old culture of *Colletotrichum gloeosporioides* in Petriplate and slants; (c) and (d) 7 day old culture of *Fusarium incarnatum* in Petriplate and slants
- Fig. 4.9** Microscopic field (100X) showing spores of the fungal isolates. (a) *Colletotrichum gloeosporioides* spores attached with mycelium, (b) *Fusarium incarnatum* spores showing macroconidia and microconidia
- Fig. 4.10** Nucleotide sequence identity matrix of *C. gloeosporioides* of the present study and other *Colletotrichum* species following 18S rRNA sequence analysis. Identity percentages are indicated on the right side corner of the matrix
- Fig. 4.11** Nucleotide sequence identity matrix of *C. gloeosporioides* of the present study and *C. gloeosporioides* isolates from different countries following 18S rRNA sequence analysis. Identity percentages are indicated on the right side corner of the matrix
- Fig. 4.12** Phylogenetic tree generated by neighbour joining of different *Colletotrichum* species. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the collection spot of the viruses have been indicated at the end of each branch
- Fig. 4.13** Phylogenetic tree generated by neighbor joining of *C. gloeosporioides* isolates from different countries. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the collection spot of the viruses have been indicated at the end of each branch

- Fig. 4.14** Nucleotide sequence identity matrix of *Fusarium* sp. of the present study and other *Fusarium* species following 18S rRNA sequence analysis. Identity percentages are indicated on the right side corner of the matrix
- Fig. 4.15** Nucleotide sequence identity matrix of *Fusarium* sp. of the present study and *F. incarnatum* isolates from different countries following 18S rRNA sequence analysis. Identity percentages are indicated on the right side corner of the matrix
- Fig. 4.16** Phylogenetic tree generated by neighbour joining of different *Fusarium* species. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the collection spot of the viruses have been indicated at the end of each branch
- Fig. 4.17** Phylogenetic tree generated by neighbour joining of *Fusarium incarnatum* isolates from different countries. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the collection spot of the viruses have been indicated at the end of each branch
- Fig. 4.18** Activity of peroxidase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Cg = *C. gloeosporioides* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Cg Inoculated plants; BABA+I = BABA treated and Cg inoculated plants; GABA+I = GABA treated and Cg inoculated plants]
- Fig. 4.19** Activity of peroxidase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Fi= *F. incarnatum* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plants]
- Fig. 4.20** Detection of induced peroxidase isoforms (POI) on PAGE. C = 'untreated-uninoculated' control plants; I = inoculated plants; BTH = BTH treated plants; BABA = BABA treated plants; GABA = GABA treated plants BTH+I = BTH treated and Inoculated

plants; BABA+I = BABA treated and inoculated plants; GABA+I = GABA treated and inoculated plants.

- Fig. 4.21** Activity of β -1,3-glucanase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Cg = *C. gloeosporioides* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Cg inoculated plants; BABA+I = BABA treated and Cg inoculated plants; GABA+I = GABA treated and Cg inoculated plants]
- Fig. 4.22** Activity of β -1,3-glucanase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Fi= *F. incarnatum* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi Inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plants]
- Fig. 4.23** Detection of β -1,3glucanaseon PAGE; C = ‘untreated-uninoculated’ control plants; I = inoculated plants; BTH = BTH treated plants; BABA = BABA treated plants; GABA = GABA treated plants BTH+I = BTH treated and Inoculated plants; BABA+I = BABA treated and inoculated plants; GABA+I = GABA treated and inoculated plants.
- Fig. 4.24** Activity of chitinase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Cg= *C. gloeosporioides* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi Inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plants]
- Fig. 4.25** Activity of chitinase in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Fi= *F. incarnatum* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi Inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plants]
- Fig. 4.26** Activity of PAL in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Cg= *C. gloeosporioides*

inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi Inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plant]

Fig. 4.27 Activity of PAL in treated and inoculated bottle gourd plants (a) BTH treated; (b) BABA treated; (c) GABA treated; [Control =Untreated and uninoculated; Fi= *F. incarnatum* inoculated; GABA= GABA treated; BABA = BABA treated; BTH= BTH treated; BTH+I = BTH treated and Fi Inoculated plants; BABA+I = BABA treated and Fi inoculated plants; GABA+I = GABA treated and Fi inoculated plants]

Fig. 4.28 Visualization of total RNA from bottle gourd leaves in 1.2% (w/v) agarose gel under UV transilluminator. [C = ‘untreated-uninoculated’ control plants; I = *C. gloeosporioides* inoculated plants; BTH = BTH treated plants; BABA = BABA treated plants; BTH+I = BTH treated and *C. gloeosporioides* inoculated plants; BABA+I = BABA treated and *C. gloeosporioides* inoculated plants]

Fig. 4.29 Agarone gel electrophoresis showing PCR amplified PAL gene product from bottle gourd leaves after purification. [C = ‘untreated-uninoculated’ control plants; I = *C. gloeosporioides* inoculated plants; BTH = BTH treated plants; BABA = BABA treated plants; BTH+I = BTH treated and *C. gloeosporioides* inoculated plants; BABA+I = BABA treated and *C. gloeosporioides* inoculated plants]

Fig. 4.30 Graphical representation of PAL Transcript accumulation. [C = ‘untreated-uninoculated’ control plants; I = *C. gloeosporioides* inoculated plants; BTH = BTH treated plants; BABA = BABA treated plants; BTH+I = BTH treated and *C. gloeosporioides* inoculated plants; BABA+I = BABA treated and *C. gloeosporioides* inoculated plants]

Fig. 4.31 Control of *Colletotrichum gloeosporioides* using different bacterial antagonists

- Culture of control plate (*C. gloeosporioides*)
- Dual culture of *S. marcescens* and *C. gloeosporioides*
- Dual culture of *P. putida* and *C. gloeosporioides*
- Dual culture of *B. amyloliquefaciens* and *C. gloeosporioides*
- Dual culture of *R. radiobacter* and *C. gloeosporioides*

Fig. 4.32 Control of *Fusarium incarnatum* using different bacterial antagonists

- a) Culture of control plate (*F. incarnatum*)
- b) Dual culture of *S. marcescens* and *F. incarnatum*
- c) Dual culture of *P. putida* and *F. incarnatum*
- d) Dual culture of *B. amyloliquefaciens* and *F. incarnatum*
- e) Dual culture of *R. radiobacter* and *F. incarnatum*

Fig. 4.33 Effect of different botanicals on growth of *Colletotrichum gloeosporioides* after 6 days of incubation after treatment with (b) *Azadirachta indica*; (c) *Datura metel* and (d) *Piper beetle* leaf extract using poisoned food technique. (a) Control

Fig. 4.34 Effect of different botanicals on growth of *Fusarium incarnatum* after 6 days of incubation after treatment with (b) *Azadirachta indica*; (c) *Holarrhena antidysenterica* and (d) *Murraya koenigii* leaf extract using poisoned food technique. (a) Control