

## LIST OF FIGURES

| Figure No.   | Title   | Page No. |
|--------------|---|----------|
| Figure- I.1  | Structure of some important biologically active molecules   | 3        |
| Figure- I.2  | Puss-pull synergistic effect  | 4        |
| Figure- I.3  | Pd-PEPPSI-NHC complexes for Suzuki-coupling reaction  | 9        |
| Figure- I.4  | Sulfonate-NHC-Pd(II) complex for Heck reaction  | 11       |
| Figure-II.1  | Somme Biologically active Biaryl Ketone compounds<br>Chemical structure of 4-quinolones   | 35       |
| Figure-II.2  | (A & B) General strategies for the synthesis of biaryl ketones using CuTC and a Pd catalyst, as previously reported. A Nucleophilic addition of $\text{ArB(OH)}_2/(\text{Het})\text{ArBF}_3\text{K}$ in the absence of S-pendant, CuTC, and a Pd catalyst is proposed herein. | 36       |
| Figure-III.1 | Biologically active diaryl ketone moieties  | 66       |
| Figure-IV.1  | Our approach towards the carbonyl insertion techniques  | 95       |
| Figure-V.1   | Structure of Tipifurnib   | 118      |
| Figure-V.2   | Pd-NHC(catalyst-A)  | 119      |