

<b>Serial no.</b>	<b>LIST OF TABLES</b>	<b>Page no.</b>
<b>Table 1.</b>	Major classes of antioxidant with their mechanism of action	17
<b>Table 2.</b>	Some antidiabetic plant species with their proposed mechanism of actions	46
<b>Table 3.</b>	GIS locations of different sampling sites for different herb and spice samples	52
<b>Table 4.</b>	Plant species selected for the present investigation	53
<b>Table 5.</b>	Herb species along with the accession no. of the voucher specimens	53
<b>Table 6.</b>	Experimental design followed in antidiabetic investigation	75
<b>Table 7.</b>	Enumeration of common dietary herbs and spices under survey and collection for the present study	86
<b>Table 8.</b>	Moisture analysis of different herb and spice samples	94
<b>Table 8A.</b>	ANOVA of data presented in Table 8 (Total moisture content)	95
<b>Table 9.</b>	Qualitative detection of major phytochemical groups present in lyophilized hot aqueous extracts of different herbs	96
<b>Table 10.</b>	Qualitative detection of major phytochemical groups present in lyophilized methanolic extracts of selected herbs	97
<b>Table 11.</b>	Qualitative detection of major phytochemical groups present in lyophilized hot aqueous extracts of different spices	98
<b>Table 12.</b>	Qualitative detection of major phytochemical groups present lyophilized in methanolic extracts from selected spices	99
<b>Table 13.</b>	Quantitative estimation of soluble sugar content and reducing sugar content present in different herb and spice samples	100
<b>Table 13A.</b>	ANOVA of data presented in Table 13 (Soluble sugar content)	101
<b>Table 13B.</b>	ANOVA of data presented in Table 13 (Reducing sugar content)	101
<b>Table 14.</b>	Quantitative estimation of soluble protein content and total lipid content present in different herb and spice samples	102
<b>Table 14A.</b>	ANOVA of data presented in Table 14 (Soluble protein content)	102
<b>Table 14B.</b>	ANOVA of data presented in Table 14 (Total lipid content)	103
<b>Table 15.</b>	Quantitative estimation of vitamin C content and vitamin E content present in different herb and spice samples	103
<b>Table 15A.</b>	ANOVA of data presented in Table 15 (Vitamin C content)	104
<b>Table 15B.</b>	ANOVA of data presented in Table 15 (Vitamin E content)	104
<b>Table 16.</b>	Quantitative determination of total polyphenol content present in lyophilized extracts of different herb and spice samples	105

<b>Serial no.</b>	<b>LIST OF TABLES</b>	<b>Page no.</b>
<b>Table 16A.</b>	ANOVA of data presented in Table 16 (Total polyphenol content)	105
<b>Table 17.</b>	Quantitative determination of total flavonoid content present in lyophilized extracts of different herb and spice samples	106
<b>Table 17A.</b>	ANOVA of data presented in Table 17 (Total flavonoid content)	106
<b>Table 18.</b>	Quantitative determination of total flavonol content present in lyophilized extracts of different herb and spice samples	107
<b>Table 18A.</b>	ANOVA of data presented in Table 18 (Total flavonol content)	107
<b>Table 19.</b>	Quantification of total chlorophyll and total carotenoid contents in different herbs	108
<b>Table 19A.</b>	ANOVA of data presented in Table 19 (Total chlorophyll content)	108
<b>Table 19B.</b>	ANOVA of data presented in Table 19 (Total carotenoids content)	108
<b>Table 20.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of DPPH scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different herbs	114
<b>Table 20A.</b>	ANOVA of data presented in Table 20 (IC <sub>50</sub> values of DPPH scavenging activity)	114
<b>Table 21.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of DPPH scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different spices	115
<b>Table 21A.</b>	ANOVA of data presented in Table 21 (IC <sub>50</sub> values of DPPH scavenging activity)	115
<b>Table 22.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of superoxide radical scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different herbs	120
<b>Table 22A.</b>	ANOVA of data presented in Table 22 (IC <sub>50</sub> values of superoxide radical scavenging activity)	120
<b>Table 23.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of superoxide radical scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different spices	121
<b>Table 23A.</b>	ANOVA of data presented in Table 23 (IC <sub>50</sub> values of superoxide radical scavenging activity)	121
<b>Table 24.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of hydrogen peroxide scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different herbs	126
<b>Table 24A.</b>	ANOVA of data presented in Table 24 (IC <sub>50</sub> values of hydrogen peroxide scavenging activity)	126
<b>Table 25.</b>	IC <sub>50</sub> values ( $\mu\text{g mL}^{-1}$ ) of hydrogen peroxide scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different spices	127

<b>Serial no.</b>	<b>LIST OF TABLES</b>	<b>Page no.</b>
<b>Table 25A.</b>	ANOVA of data presented in Table 25 (IC <sub>50</sub> values of hydrogen peroxide scavenging activity)	127
<b>Table 26.</b>	IC <sub>50</sub> values (µg mL <sup>-1</sup> ) of nitric oxide radical scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different herbs	132
<b>Table 26A</b>	ANOVA of data presented in Table 26 (IC <sub>50</sub> values of nitric oxide radical scavenging activity)	132
<b>Table 27.</b>	IC <sub>50</sub> values (µg mL <sup>-1</sup> ) of nitric oxide radical scavenging activity exhibited by lyophilized hot aqueous and methanolic extracts of different spices	133
<b>Table 27A.</b>	ANOVA of data presented in Table 27 (IC <sub>50</sub> values of nitric oxide radical scavenging activity)	133
<b>Table 28.</b>	Comparative antibacterial activities of different herbs against Gram positive and Gram negative bacterial strains at different doses of lyophilized extracts	135
<b>Table 29.</b>	Minimum inhibitory dose (MID) determination of herb extracts against target bacterial strains using DAD assay	136
<b>Table 30.</b>	Comparative antibacterial activities of different spices against Gram positive and Gram negative bacterial strains at different doses of extracts	141
<b>Table 31.</b>	Minimum inhibitory dose (MID) determination of spice extracts against target bacterial strains using DAD assay	143
<b>Table 32.</b>	Preliminary screening for anti-QS activity by intact herb and spice sample against <i>C. violaceum</i> 2656	144
<b>Table 33.</b>	Comparative anti-QS activities of selected herb and spice extracts on <i>C. violaceum</i> 2656 by DAD assay	145
<b>Table 33A.</b>	ANOVA of data presented in Table 33 (Antimicrobial activity)	146
<b>Table 33B.</b>	ANOVA of data presented in Table 33 (Anti-QS activity)	146
<b>Table 34.</b>	Inhibitory effect of <i>I. verum</i> methanolic extract on protease production by <i>Ps. aeruginosa</i> .	149
<b>Table 35.</b>	Effect of spice extracts on swarming motility of <i>Ps. aeruginosa</i>	150
<b>Table 36.</b>	Effect of <i>I. verum</i> and <i>G. oppositifolius</i> methanolic extracts on body weight in diabetic rats	153
<b>Table 36A.</b>	ANOVA of data presented in Table 36 (body weight change in experimental rats)	153
<b>Table 37.</b>	Effect of <i>I. verum</i> and <i>G. oppositifolius</i> methanolic extracts on blood glucose level in diabetic rats	154

<b>Serial no.</b>	<b>LIST OF TABLES</b>	<b>Page no.</b>
<b>Table 37A.</b>	ANOVA of data presented in Table 37 (blood glucose level change in experimental rats)	155
<b>Table 38.</b>	Effect of <i>I. verum</i> and <i>G. oppositifolius</i> methanolic extracts on serum lipid profile in normal and Streptozotocin-induced diabetic rats	155
<b>Table 38A.</b>	ANOVA of data presented in Table 38 (Total cholesterol)	156
<b>Table 38B.</b>	ANOVA of data presented in Table 38 (Total triglycerides)	156
<b>Table 38C.</b>	ANOVA of data presented in Table 38 (HDL cholesterol)	156
<b>Table 39.</b>	Effect of <i>I. verum</i> and <i>G. oppositifolius</i> extracts on liver enzymes (SGPT and SGOT level) in normal and streptozotocin induced diabetic rats	157
<b>Table 39A.</b>	ANOVA of data presented in Table 39 (SGPT level)	157
<b>Table 39B.</b>	ANOVA of data presented in Table 39 (SGOT level)	158
<b>Table 40.</b>	Effect of <i>I. verum</i> and <i>G. oppositifolius</i> extracts on serum urea and creatinine level of normal and Streptozotocin-induced diabetic rats	158
<b>Table 40A.</b>	ANOVA of data presented in Table 40 (Serum urea)	158
<b>Table 40B.</b>	ANOVA of data presented in Table 40 (Serum creatinine)	159
<b>Table 41.</b>	Compounds identified in the methanolic fraction of <i>I. verum</i> ( <i>IvME</i> )	162
<b>Table 42.</b>	Compounds identified in the hexane fraction of <i>I. verum</i> ( <i>IvHE</i> )	169
<b>Table 43.</b>	Compounds identified in the ethyl acetate fraction of <i>I. verum</i> ( <i>IvEA</i> )	176
<b>Table 44.</b>	Compounds identified in the methanolic fraction of <i>G. oppositifolius</i> ( <i>GoME</i> )	186
<b>Table 45.</b>	Compounds identified in the hexane fraction of <i>G. oppositifolius</i> ( <i>GoHE</i> )	193
<b>Table 46.</b>	Compounds identified in the ethyl acetate fraction of <i>G. oppositifolius</i> ( <i>GoEA</i> )	199