

Serial no.	LIST OF FIGURES	Page no.
Figure 1.	Diseases and disorders caused by reactive oxygen species, ROS	14
Figure 2.	Mechanism of action of plant-derived antimicrobials on possible targets	28
Figure 3.	Herbs and spices selected for the bioassay	55
Figure 4.	Illustration of maintenance and handling of rats for antidiabetic test	77
Figure 5.	Water soluble extractive (WSE) and methanol soluble extractive (MSE) values of selected herbs and spices	94
Figure 6.	DPPH scavenging activity of lyophilized hot aqueous extracts of different herbs	110
Figure 7.	DPPH scavenging activity of lyophilized methanolic extracts of different herbs	111
Figure 8.	DPPH scavenging activity of lyophilized hot aqueous extracts of different spices.	112
Figure 9.	DPPH scavenging activity of lyophilized methanolic extracts of different spices.	113
Figure 10.	Superoxide radical scavenging activity of lyophilized hot aqueous extracts of different herbs	116
Figure 11.	Superoxide radical scavenging activity of lyophilized methanolic extracts of different herbs.	117
Figure 12.	Superoxide radical scavenging activity of lyophilized hot aqueous extracts of different spices.	118
Figure 13.	Superoxide radical scavenging activity of lyophilized methanolic extracts of different spices	119
Figure 14.	Hydrogen peroxide (H ₂ O ₂) scavenging activity of lyophilized hot aqueous extracts of different herbs	122
Figure 15.	Hydrogen peroxide (H ₂ O ₂) scavenging activity of lyophilized methanolic extracts of different herbs	123
Figure 16.	Hydrogen peroxide (H ₂ O ₂) scavenging activity of lyophilized hot aqueous extracts of different spices	124
Figure 17.	Hydrogen peroxide (H ₂ O ₂) scavenging activity of lyophilized methanolic extracts of different spices	125
Figure 18.	Nitric oxide (NO) scavenging activity of lyophilized hot aqueous extracts of different herbs	128
Figure 19.	Nitric oxide (NO) scavenging activity of lyophilized methanolic extracts of different herbs	129

Serial no.	LIST OF FIGURES	Page no.
Figure 20.	Nitric oxide (NO) scavenging activity of lyophilized hot aqueous extracts of different spices	130
Figure 21.	Nitric oxide (NO) scavenging activity of lyophilized methanolic extracts of different spices	131
Figure 22.	Antibacterial activity of <i>M. piperita</i> methanolic extract	136
Figure 23.	Antibacterial activity of <i>T. foenum-graecum</i> methanolic extract	137
Figure 24.	Antibacterial activity of <i>C. sativum</i> methanolic extract	137
Figure 25.	Antibacterial activity of <i>M. koenigii</i> methanolic extract	138
Figure 26.	Antibacterial activity of <i>G. oppositifolius</i> methanolic extract	138
Figure 27.	Antibacterial activity of <i>I. verum</i> methanolic extract	139
Figure 28.	Antibacterial activity of <i>M. fragrans</i> methanolic extract	139
Figure 29.	Antibacterial activity of <i>C. annuum</i> methanolic extract	140
Figure 30.	Antibacterial activity of <i>P. perlata</i> methanolic extract	140
Figure 31.	Antibacterial activity of <i>D. volubilis</i> methanolic extract	141
Figure 32.	Effect of different spice extracts on violacein synthesis in <i>C. violaceum</i> at different doses (3-10 mg disc ⁻¹) evaluated by disc-agar diffusion assay	146
Figure 33.	Graphical representation showing the effect of <i>I. verum</i> on violacein production by wild-type <i>C. violaceum</i>	147
Figure 34.	Effect of <i>I. verum</i> on violacein production by wild-type <i>C. violaceum</i>	147
Figure 35.	Graphical representation of the effect of <i>I. verum</i> on pyocyanin production by <i>Ps. aeruginosa</i> at different concentrations (100-1000 µg mL ⁻¹)	148
Figure 36.	Effect of <i>I. verum</i> methanolic extract on synthesis of pyocyanin by <i>Ps. aeruginosa</i> at C (control): 0; 1: 100; 2: 250; 3: 500; 4: 750 and 5: 1000 µg mL ⁻¹ concentrations	148
Figure 37.	Effect of <i>I. verum</i> methanolic extract on protease production by <i>Ps. aeruginosa</i> at different concentrations	149
Figure 38.	Effect of <i>I. verum</i> methanolic extract on swarming behaviour of <i>Ps. aeruginosa</i> . (a) Untreated control, (b-e) treated with <i>I. verum</i> extract(150, 200, 250 and 500 µg mL ⁻¹)respectively	150
Figure 39.	Effect of <i>I. verum</i> lyophilized methanolic extract on <i>in vitro</i> biofilm formation by <i>Ps. aeruginosa</i> at A: 1000; B: 750; C: 500; D: 250 and E: 100 µg mL ⁻¹ . F: Control	151

Serial no.	LIST OF FIGURES	Page no.
Figure 40.	Graphical representation of biofilm inhibition by methanolic extract of <i>I. verum</i> fruit at different concentrations	151
Figure 41.	Partially purified different fractions of <i>I. verum</i> and <i>G. oppositifolius</i> . A. Methanolic fraction, B. Ethyl acetate fraction, C. Hexane fraction of <i>I. verum</i> ; D. Methanolic fraction, E. Ethyl acetate fraction and E. Hexane fraction of <i>G. oppositifolius</i>	160
Figure 42.	GC-MS Total Ion Chromatogram (TIC) of methanolic fraction of <i>I. verum</i> (IvME)	161
Figure 43.	Mass spectra of the compounds in the GC peaks of retention time of 10.167, 13.009, 15.377 and 16.052 min, which were identified as A. Linalool; B. Estragole, C. Benzaldehyde, 4-methoxy- and D. Benzene, 1-methoxy-4-(1-propenyl)- or cis-Anothole respectively by the MS library	166
Figure 44.	Mass spectra of the compounds in the GC peaks of retention time of 18.538, 22.467, 25.235 and 31.292 min, which were identified as A. 2-Propanone, 1-(4-methoxyphenyl)-, B. Benzhydrazide, 4-methoxy-N2-(2-trifluoroacetyl cyclohepten-1-yl)-, C. 1-(3-Methyl-2-butenoxy)-4-(1-propenyl) benzene and D. n-Hexadecanoic acid respectively by the MS library	167
Figure 45.	Mass spectra of the compounds in the GC peaks of retention time of 34.571, 34.890, 41.093 and 46.215 min, which were identified as A. cis-Vaccenic acid, B. Octadecanoic acid, C. (2R,3S,5S,6R)-2,5-bis(4-Methoxyphenyl)-3,6-dimethyl-1,4-dioxane-rel-, and D. Carinol respectively by the MS library	168
Figure 46.	GC-MS Total Ion Chromatogram (TIC) of hexane fraction of <i>I. verum</i> (IvHE)	169
Figure 47.	Mass spectra of the compounds in the GC peaks of retention time of 14.617, 15.592, 17.491 and 17.987 min, which were identified as A. Benzaldehyde, 4-methoxy-; B. Anethole, C. Anisaldehyde dimethyl acetal, and D. 2-Propanone, 1-(4-methoxyphenyl)- respectively by the MS library	173
Figure 48.	Mass spectra of the compounds in the GC peaks of retention time of 22.920, 23.105, 25.184 and 31.121 min, which were identified as A. 4-(p-Methoxyphenyl)-1-butanol, B. 1-(4-Methoxyphenyl) propane-1,2-diol; C. 1-(3-Methyl-2-butenoxy)-4-(1-propenyl) benzene and D. n-Hexadecanoic acid respectively by the MS library	174

Serial no.	LIST OF FIGURES	Page no.
Figure 49.	Mass spectra of the compounds in the GC peaks of retention time of 34.425, 39.135, 41.829 and 42.154 min, which were identified as A. cis-Vaccenic acid, B. (2R,4R,5S)-2,4-bis (4-Methoxyphenyl)-5-methyl-1,3-dioxolane-rel-; C. 4-Methoxybenzoic acid N'-[2-(4-methoxy-phenyl)-acetyl]-hydrazide and D. Ethanone, 2-hydroxy-1,2-bis (4-methoxyphenyl)- respectively by the MS library	175
Figure 50.	GC-MS Total Ion Chromatogram (TIC) of ethyl acetate fraction of <i>I. verum</i> (IvEA)	176
Figure 51.	Mass spectra of the compounds in the GC peaks of retention time of 10.162, 13.038, 16.249 and 18.541 min, which were identified as A. Linalool, B. Estragole, C. Benzene, 1-methoxy-4-(1-propenyl)- and D. 2-Propanone, 1-(4-methoxyphenyl)- respectively by the MS library	182
Figure 52.	Mass spectra of the compounds in the GC peaks of retention time of 19.397, 22.448, 23.430 and 25.334 min, which were identified as A. Trans-.alpha.-bergamotene, B. S-(p-Methoxybenzoyl) thiohydroxylamine, C. 1-(4-Methoxyphenyl)propane-1,2-diol and D. 1-(3-Methyl-2-butenoxy)-4-(1-propenyl)benzene respectively by the MS library	183
Figure 53.	Mass spectra of the compounds in the GC peaks of retention time of 31.366, 34.700, 39.212 and 39.670 min, which were identified as A. n-Hexadecanoic acid, B. cis-Vaccenic acid, C. (2R,4R,5S)-2,4-bis(4-Methoxyphenyl)-5-methyl-1,3-dioxolane-rel- and D. Benzene, 1-methoxy-4-(1-methylpropyl)- respectively by the MS library	184
Figure 54.	Mass spectra of the compounds in the GC peaks of retention time of 41.140, 41.926 and 51.659 min, which were identified as A. (2R,3S,5S,6R)-2,5-bis (4-Methoxyphenyl)-3,6-dimethyl-1,4-dioxane-rel-, B. (2R,3S,5S,6R)-2,5-bis(4-Methoxyphenyl)-3,6-dimethyl-1,4-dioxane-rel- and C. Stigmast-5-en-3-ol, (3.beta.)- respectively by the MS library	185
Figure 55.	GC-MS Total Ion Chromatogram (TIC) of methanolic fraction of <i>G. oppositifolius</i> (GoME)	186
Figure 56.	Mass spectra of the compounds in the GC peaks of retention time of 10.278, 28.108 31.082 and 33.779 min, which were identified as A. 1H-Pyrrole, 2,5-dihydro-; B. 1-Deutero-2,2,5,5-tetramethylcyclopentanol, C. n-Hexadecanoic acid and D. Phytol respectively by the MS library	191

Serial no.	LIST OF FIGURES	Page no.
Figure 57.	Mass spectra of the compounds in the GC peaks of retention time of 34.335, 52.984, 54.854 and 55.407 min, which were identified as A. 8,11,14-Eicosatrienoic acid, (Z,Z,Z)-; B. Lanost-8-en-3-ol, (3.beta.)-; C. (4aS, 8S, 8aR)-8-Isopropyl-5-methyl-3,4,4a,7,8,8a-hexahydronaphthalen-2-yl; and D. Retinol, acetate respectively by the MS library	192
Figure 58.	GC-MS Total Ion Chromatogram (TIC) of hexane fraction of <i>G. oppositifolius</i> (GoHE)	193
Figure 59.	Mass spectra of the compounds in the GC peaks of retention time of 30.261, 31.048, 33.542 and 33.772 min, which were identified as A. Hexadecanoic acid, methyl ester; B. n-Hexadecanoic acid; C. 9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-; and D. Phytol 34 respectively by the MS library	197
Figure 60.	Mass spectra of the compounds in the GC peaks of retention time of 34.151, 34.714, 44.943 and 51.491 min, which were identified as A. Linoelaidic acid, B. Octadecanoic acid, C. Squalene and D. Chondrillasterol respectively by the MS library	198
Figure 61.	GC-MS Total Ion Chromatogram (TIC) of ethyl acetate fraction of <i>G. oppositifolius</i> (GoEA)	199
Figure 62.	Mass spectra of the compounds in the GC peaks of retention time of 10.257, 17.908, 31.021 and 33.759 min, which were identified as A. 1H-Pyrrole, 2,5-dihydro-, B. DL-Proline, 5-oxo-, methyl ester, C. DL-Proline, 5-oxo-, methyl ester and D. Phytol respectively by the MS library	203
Figure 63.	Mass spectra of the compounds in the GC peaks of retention time of 50.432, 52.731, 54.804 and 55.338 min, which were identified as A. Lanosterol, B. Lup-20(29)-en-28-ol, C. Beta.-copaen-4 .alpha.-ol and D. Retinol, acetate respectively by the MS library	204