

## **LIST OF FIGURES**

<b>1.1</b> Schematic representation of the molecular arrangements in the nematic phase .....	4
<b>1.2</b> Schematic representation of molecular arrangements in the Cholesteric (N*) phase.....	5
<b>1.3</b> Schematic representation of molecular arrangements in the smectic A phase .	7
<b>1.4</b> Schematic representation of molecular arrangements in the smectic C phase..	8
<b>1.5</b> Schematic representation of molecular arrangements in the smectic B, smectic I and smectic F phase .....	9
<b>1.6</b> Schematic representation of molecular arrangements in the crystal B, crystal J and crystal G phase .....	9
<b>1.7</b> Schematic representation of molecular arrangements in the ferroelectric (SmC*) and antiferroelectric (SmC <sub>A</sub> *) phase .....	11
<b>1.8</b> Schematic representation of molecular arrangements in the discotic phase ...	12
<b>1.9</b> Schematic representation of blue phase structure showing blue phase I (BPI) and blue phase II (BPPII) .....	13
<b>1.10</b> Schematic representation of molecular arrangements in the twist grain boundary phase.....	14
<b>1.11</b> Schematic representation of bent core liquid crystals.....	15
<b>2.1</b> Schematic diagram of the experimental set-up for refractive index measurement from the thin prism technique.....	40
<b>2.2</b> Schematic diagram of the experimental set-up for high resolution optical birefringence ( $\Delta n$ ) measurement .....	41
<b>2.3</b> Schematic representation of the director configuration is shown (a) below threshold voltage ( $V_{th}$ ) and (b) above the threshold voltage ( $V_{th}$ ) respectively .....	45
<b>2.4</b> Voltage dependent capacitance change due to Freedericksz transition in a liquid crystal. The arrow indicates the Freedericksz threshold voltage ( $V_{th}$ ) .....	46

## List of Figures

<b>2.5</b> Difference in the free energy density ( $\Delta f$ ) as a function of scalar order parameter S.....	60
<b>2.6</b> The nematic order parameter (S) vs. temperature. $T^*$ and $T^{**}$ are the temperatures slightly lower and higher than $T_{NI}$ respectively. $T^*$ and $T^{**}$ are also known as the spinodal temperatures. ....	60
<b>3.1</b> Phase diagram of the binary system 9OCB+7CB. $x_{9OCB}$ represents the mole fraction of 9OCB. I- isotropic phase, N- nematic phase and SmA- smectic A phase. o - Nematic to isotropic phase transition temperature ( $T_{NI}$ ); □- smectic A to nematic phase transition temperature ( $T_{SN}$ ) .....	76
<b>3.2</b> Phase diagram of the binary system 10OCB+7CB. $x_{10OCB}$ represents the mole fraction of 10OCB. I- isotropic phase, N- nematic phase and SmA- smectic A phase. o – nematic to isotropic phase transition temperature ( $T_{NI}$ ); □ – smectic A to nematic phase transition temperature ( $T_{SN}$ ) .....	77
<b>3.3</b> Phase diagram of the binary system 8OCB + 8CB. $x_{8OCB}$ represents the mole fraction of 8OCB. I- isotropic phase, N- nematic phase and SmA- smectic A phase. o- nematic to isotropic phase transition temperature ( $T_{NI}$ ); □- smectic A to nematic phase transition temperature ( $T_{SN}$ ). Solid, dashed and dotted line represent the nematic-isotropic, smectic A-nematic phase transition and melting temperatures respectively throughout the entire composition as obtained in ref. [25,26] .....	78
<b>3.4(a-b)</b> Temperature variation of birefringence ( $\Delta n$ ) for (a) $x_{9OCB} = 0.200$ and (b) $x_{9OCB} = 0.270$ of the binary system 9OCB+7CB. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines represent a fit to Equation (3.2). In the inset variation of $\Delta n$ in the vicinity of $T_{SN}$ is shown.....	80
<b>3.4(c-d)</b> Temperature variation of birefringence ( $\Delta n$ ) for (c) $x_{9OCB} = 0.300$ and (d) $x_{9OCB} = 0.370$ of the binary system 9OCB+7CB. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition	

## List of Figures

---

temperatures respectively. The solid lines represent a fit to Equation (3.2). In the inset variation of $\Delta n$ in the vicinity of $T_{SN}$ is shown.....	81
<b>3.4(e-f)</b> Temperature variation of birefringence ( $\Delta n$ ) for (e) $x_{9OCB} = 0.420$ and (f) $x_{9OCB} = 0.500$ of the binary system 9OCB+7CB. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines represent a fit to Equation (3.2). In the inset variation of $\Delta n$ in the vicinity of $T_{SN}$ is shown.....	82
<b>3.4(g-h)</b> Temperature variation of birefringence ( $\Delta n$ ) for (g) $x_{9OCB} = 0.620$ and (h) $x_{9OCB} = 0.700$ of the binary system 9OCB+7CB. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines represent a fit to Equation (3.2). In the inset variation of $\Delta n$ in the vicinity of $T_{SN}$ is shown.....	83
<b>3.4(i-j)</b> Temperature variation of birefringence ( $\Delta n$ ) for (i) $x_{9OCB} = 0.810$ and (j) $x_{9OCB} = 0.890$ of the binary system 9OCB+7CB. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines represent a fit to Equation (3.2). In the inset variation of $\Delta n$ in the vicinity of $T_{SN}$ is shown.....	84
<b>3.5(a-b)</b> Birefringence ( $\Delta n$ ) as a function of temperature for the different mixtures of the binary system (a) 10OCB+7CB and (b) 8OCB+8CB. Different mole fractions are indicated in the figure. Solid upper head arrows indicate nematic–isotropic phase transition temperature ( $T_{NI}$ ) and dashed lower head arrows indicate smectic A-nematic phase transition ( $T_{SN}$ ) temperature.....	85
<b>3.6</b> Change in the birefringence value $\delta(\Delta n)$ near the smectic A-nematic (SmA-N) phase transition as a function of mixture concentrations for the binary system 9OCB+7CB.....	86
<b>3.7</b> Temperature variation of $Q(T)$ near the SmA-N phase transition at different concentrations of 9OCB. Solid lines are fit to Equation (3.4) for $Q(T)$ for the different mixtures. Different concentrations are mentioned in the figures.....	90

## List of Figures

- 3.8(a-b)** Temperature variation of  $Q(T)$  near the SmA-N phase transition at different concentrations for the binary system (a) 10OCB+7CB and (b) 8OCB+8CB. Different concentrations are mentioned in the figures. Solid lines are fit to Equation (3.4) for  $Q(T)$  for the different mixtures ..... 91
- 3.9** Variation of the critical exponent ( $\alpha$ ) with McMillan ratio ( $T_{SN}/T_{NI}$ ) for the system 9OCB+7CB. (●) represents the  $\alpha$  values averaged over the two values obtained from fitting above and below the transition temperature  $T_{SN}$ . The vertical bars indicate the error associated with  $\alpha$  value. The  $\alpha$  value ( $\square$ ) for  $x_{9OCB}=1.0$  is obtained from Cusmin *et al.* [36]. Dashed line represents a polynomial fit to the  $\alpha$  values. Upper and lower head arrows denote the tricritical point (TCP) and the 3D-XY limit of SmA-N phase transition respectively ..... 95
- 3.10** Variation of the critical exponent ( $\alpha$ ) with McMillan ratio ( $T_{SN}/T_{NI}$ ) for the system 10OCB+7CB. (●) represents the  $\alpha$  values averaged over the two values obtained from fitting above and below the transition temperature  $T_{SN}$ . The vertical bars indicate the error associated with  $\alpha$  value. Dashed line represents a polynomial fit to the  $\alpha$  values. Upper and lower head arrows denote the tricritical point (TCP) and the 3D-XY limit of the SmA-N phase transition respectively..... 96
- 3.11** Variation of critical exponent ( $\alpha$ ) with McMillan ratio for the system 8OCB+8CB. (●)-  $\alpha$  values averaged over the two values obtained from fitting above and below the transition temperature  $T_{SN}$ . ( $\square$ ) -  $\alpha$  value obtained from the Ref. [40] and [41] for 8OCB and 8CB respectively. (★) -  $\alpha$  value obtained by Sied *et al.* [26] for the present system. The vertical bars indicate the error associated with  $\alpha$  value. Dashed line represents a polynomial fit to the  $\alpha$  values. Upper and lower head arrows denote the TCP and the 3D-XY limit of SmA-N transition respectively.. ..... 97
- 3.12** The critical exponent ( $\alpha$ ) vs. McMillan ratio ( $T_{SN}/T_{NI}$ ). The  $\alpha$  values ( $\square$ ) for the three pure compounds *i.e.* 8CB, 8OCB and 9OCB are obtained from the different literatures [37,40,41]. Dashed line represents a polynomial fit to the  $\alpha$

## List of Figures

values. The vertical bar indicates the error associated with $\alpha$ value. Upper and lower head arrows denote the tricritical point (TCP) and the 3D-XY limit of the SmA-N phase transition respectively .....	98
<b>3.13</b> Temperature variation of the transmitted intensity ( $I_t$ ) in the vicinity of the nematic-isotropic phase transition for the mixture $x_{9OCB} = 0.890$ . The horizontal dashed line represents a linear fit to the transmitted intensity ( $I_t$ ) data far away from the N-I transition temperature ( $T_{NI}$ ).....	100
<b>3.14</b> Temperature variation of the parameter $Q(T)$ near the N-I phase transition at different concentrations of 9OCB for the binary system 9OCB+7CB. Solid lines are fit to Equation (3.4) for different mixtures near the N-I phase transition .....	101
<b>4.1</b> Phase diagram of the binary system 5DBT + 10OCB. $x_{5DBT}$ is the mole fraction of 5DBT. I- isotropic phase, N- nematic phase and SmA- smectic A phase. o- nematic to isotropic (or smectic A to isotropic) phase transition temperature; ■- smectic A to nematic phase transition temperature and □-melting temperature.....	113
<b>4.2</b> Microphotographs (400x magnification) of the textures observed under the polarizing optical microscope for the concentration $x_{5DBT} = 0.741$ . (a) Marble texture of the nematic phase at 62°C during heating cycle, (b) Four brush schlieren texture of the nematic phase at 64°C during cooling, (c) Striped pattern in the smectic A phase at 40°C in the heating cycle. (d) Striped pattern in the smectic A phase at 35°C in the cooling cycle .....	114
<b>4.3(a)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for $x_{5DBT} = 0.100$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic phase transition ( $T_{SN}$ ) temperatures respectively.....	115
<b>4.3(b-c)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (b) $x_{5DBT} = 0.161$ and (c) $x_{5DBT} = 0.203$ . Solid arrow and	

## List of Figures

---

dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic phase transition ( $T_{SN}$ ) temperatures respectively .....	116
<b>4.3(d-e)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (d) $x_{5DBT} = 0.252$ and (e) $x_{5DBT} = 0.301$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic phase transition ( $T_{SN}$ ) temperatures respectively .....	117
<b>4.3(f-g)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (f) $x_{5DBT} = 0.447$ and (g) $x_{5DBT} = 0.503$ . Solid arrow denotes the nematic-isotropic ( $T_{NI}$ ) phase transition temperature.....	118
<b>4.3(h-i)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (h) $x_{5DBT} = 0.604$ and (i) $x_{5DBT} = 0.696$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic phase transition ( $T_{SN}$ ) temperatures respectively .....	119
<b>4.3(j-k)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (j) $x_{5DBT} = 0.741$ and (k) $x_{5DBT} = 0.804$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic phase transition ( $T_{SN}$ ) temperatures respectively .....	120
<b>4.3(l-m)</b> Temperature variation of the extraordinary ( $n_e$ ) and ordinary ( $n_o$ ) refractive indices for (l) $x_{5DBT} = 0.887$ and (m) $x_{5DBT} = 0.952$ . Solid arrow, dashed arrow and dotted arrow denote the nematic-isotropic ( $T_{NI}$ ), smectic A-nematic ( $T_{SN}$ ) and smectic A-isotropic ( $T_{SI}$ ) phase transition temperatures respectively..	121
<b>4.4(a)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (a) $x_{5DBT} = 0.100$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) phase transition temperatures respectively .....	122
<b>4.4(b-c)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (b) $x_{5DBT} = 0.161$ and (c) $x_{5DBT} = 0.203$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow and dashed arrow denote the	

## List of Figures

nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) phase transition temperatures respectively.....	123
<b>4.4(d-e)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (d) $x_{5DBT} = 0.252$ and (e) $x_{5DBT} = 0.301$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) phase transition temperatures respectively.....	124
<b>4.4(f-g)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (f) $x_{5DBT} = 0.447$ and (g) $x_{5DBT} = 0.503$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow denotes the nematic-isotropic ( $T_{NI}$ ) phase transition temperature.....	125
<b>4.4(h-i)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (h) $x_{5DBT} = 0.604$ and (i) $x_{5DBT} = 0.696$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) phase transition temperatures respectively.....	126
<b>4.4(j-k)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (j) $x_{5DBT} = 0.741$ and (k) $x_{5DBT} = 0.804$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) phase transition temperatures respectively.....	127
<b>4.4(l-m)</b> Birefringence ( $\Delta n$ ) as a function of temperature for (l) $x_{5DBT} = 0.887$ and (m) $x_{5DBT} = 0.952$ . ( $\Delta$ )–OT data for $\lambda = 632.8\text{nm}$ ; ( $\square$ )–OT data for $\lambda = 532.0\text{nm}$ ; ( $\bullet$ )–Thin prism data for $\lambda = 632.8\text{nm}$ . Solid arrow, dashed arrow and dotted arrow denote the nematic–isotropic ( $T_{NI}$ ), smectic A-nematic ( $T_{SN}$ ) and smectic A-isotropic ( $T_{SI}$ ) phase transition temperatures respectively .....	128
<b>4.5</b> Birefringence ( $\Delta n$ ) as a function of mixture concentration at $T = 330\text{K}$ obtained from two different methods and at different wavelengths .....	129

## List of Figures

<b>4.6(a-b)</b> Temperature variation of the parameter $Q(T)$ in the vicinity of smectic A-nematic (SmA-N) phase transition. Open circles (o) represent the calculated $Q(T)$ values (obtained from $\Delta n$ data for $\lambda = 532\text{nm}$ ) and the solid lines are fit to Equation (4.2). Different concentrations are indicated in the figures.....	131
<b>4.7</b> Concentration dependence of the critical exponent ( $\alpha$ ) obtained by fitting the $Q(T)$ data to Equation (4.2). The plotted data are taken by averaging the two $\alpha$ values obtained from the fitting for $T < T_{\text{SN}}$ and $T > T_{\text{SN}}$ . Vertical bar represents the error associated with the critical exponent ( $\alpha$ ). The vertical dashed line corresponds to the tricritical points (TCP). The solid lines are 2 <sup>nd</sup> order polynomial fit to the $\alpha$ values.....	133
<b>4.8</b> Variation of the critical exponent ( $\alpha$ ) with the McMillan ratio ( $T_{\text{SN}}/T_{\text{NI}}$ ). The solid line is polynomial fit to the $\alpha$ values. Vertical bar represents the error associated with the critical exponent ( $\alpha$ ) .....	134
<b>5.1(a-b)</b> The dipole moments of (a) 5DBT and (b) 10OCB in <i>p</i> -xylene solution as a function of weight percentage .....	144
<b>5.2(a-b)</b> Temperature variation of the dielectric parameters for (a) $x_{\text{5DBT}} = 0.0$ (10OCB) and (b) $x_{\text{5DBT}} = 0.100$ . $x_{\text{5DBT}}$ is the mole fraction of 5DBT. Solid arrow, dashed arrow and dotted arrow indicate nematic–isotropic ( $T_{\text{NI}}$ ), smectic A–nematic ( $T_{\text{SN}}$ ) and smectic A–isotropic ( $T_{\text{SI}}$ ) phase transition temperatures respectively.....	146
<b>5.2(c-d)</b> Temperature variation of the dielectric parameters for (c) $x_{\text{5DBT}} = 0.161$ and (d) $x_{\text{5DBT}} = 0.203$ . $x_{\text{5DBT}}$ is the mole fraction of 5DBT. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{\text{NI}}$ ) and smectic A–nematic ( $T_{\text{SN}}$ ) phase transition temperatures respectively.....	147
<b>5.2(e-f)</b> Temperature variation of the dielectric parameters for (e) $x_{\text{5DBT}} = 0.252$ and (f) $x_{\text{5DBT}} = 0.301$ . $x_{\text{5DBT}}$ is the mole fraction of 5DBT. Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{\text{NI}}$ ) and smectic A–nematic ( $T_{\text{SN}}$ ) phase transition temperatures respectively.....	148

## List of Figures

<p><b>5.2(g-h)</b> Temperature variation of the dielectric parameters for (g) <math>x_{5DBT} = 0.447</math> and (h) <math>x_{5DBT} = 0.503</math>. <math>x_{5DBT}</math> is the mole fraction of 5DBT. Solid arrow indicates nematic–isotropic (<math>T_{NI}</math>) phase transition temperature.....</p>	149
<p><b>5.2(i-j)</b> Temperature variation of the dielectric parameters for (i) <math>x_{5DBT} = 0.604</math> and (j) <math>x_{5DBT} = 0.696</math>. <math>x_{5DBT}</math> is the mole fraction of 5DBT. Solid arrow and dashed arrow indicate nematic–isotropic (<math>T_{NI}</math>) and smectic A–nematic (<math>T_{SN}</math>) phase transition temperatures respectively.....</p>	150
<p><b>5.2(k-l)</b> Temperature variation of the dielectric parameters for (k) <math>x_{5DBT} = 0.741</math> and (l) <math>x_{5DBT} = 0.803</math>. <math>x_{5DBT}</math> is the mole fraction of 5DBT. Solid arrow and dashed arrow indicate nematic–isotropic (<math>T_{NI}</math>) and smectic A–nematic (<math>T_{SN}</math>) phase transition temperatures respectively.....</p>	151
<p><b>5.2(m-n)</b> Temperature variation of the dielectric parameters for (m) <math>x_{5DBT} = 0.887</math> and (n) <math>x_{5DBT} = 0.952</math>. <math>x_{5DBT}</math> is the mole fraction of 5DBT. Solid arrow, dashed arrow and dotted arrow indicate nematic–isotropic (<math>T_{NI}</math>), smectic A–nematic (<math>T_{SN}</math>) and smectic A-isotropic (<math>T_{SI}</math>) phase transition temperatures respectively.....</p>	152
<p><b>5.2(o)</b> Temperature variation of the dielectric parameters for <math>x_{5DBT} = 1.0</math> (5DBT). <math>x_{5DBT}</math> is the mole fraction of 5DBT. Dotted arrow indicates smectic A-isotropic (<math>T_{SI}</math>) phase transition temperature.....</p>	153
<p><b>5.3(a)</b> Variation of <math>(\epsilon_{iso} - \epsilon_{av})</math> with the mole fraction of 5DBT.....</p>	153
<p><b>5.3(b)</b> Variation of <math>\epsilon_{  }</math>, <math>\epsilon_{\perp}</math>, <math>\epsilon_{av}</math>, <math>\epsilon_{iso}</math> and <math>\Delta\epsilon</math> with the mole fraction of 5DBT at 50°C. (■) -<math>\epsilon_{  }</math>, (o) -<math>\epsilon_{av}</math>, (□) -<math>\Delta\epsilon</math>, (●) - <math>\epsilon_{iso}</math> and (◇) - <math>\epsilon_{\perp}</math>.....</p>	154
<p><b>5.4</b> Temperature dependence of the effective dipole moment (<math>\mu_{eff}</math>) and the angle of inclination (<math>\omega</math>) for the mixture <math>x_{5DBT} = 0.301</math>. (o) represents <math>\mu_{eff}</math> and (□) represents <math>\omega</math>.....</p>	156
<p><b>5.5</b> Dielectric permittivity as a function of temperature for the concentration <math>x_{5DBT} = 0.301</math>. Solid lines are fit to Equations (5.5) and (5.6). The fitting parameters in both nematic and isotropic phases are mentioned in the figure. The temperatures</p>	

## List of Figures

---

<p><math>T^*</math> and <math>T^{**}</math> are indicated with a star symbol (<math>\star</math>) and the dotted line represents the nematic-isotropic phase transition temperature (<math>T_{NI}</math>).....</p>	158
<p><b>5.6(a)</b> Temperature variation of the dielectric permittivity for the different mixtures measured in the isotropic phase. The concentrations are indicated in the figures. Solid lines are the fit to Equation (5.5) and the star symbol represents the critical temperature (<math>T^*</math>).....</p>	159
<p><b>5.6(b)</b> Temperature variation of the dielectric permittivity for the different mixtures measured in the isotropic phase. The concentrations are indicated in the figures. Solid lines are the fit to Equation (5.5) and the star symbol represents the critical temperature (<math>T^*</math>).....</p>	160
<p><b>5.7(a-b)</b> Concentration dependence of (a) the dielectric constant (<math>\epsilon^*</math>) at the critical temperature <math>T^*</math> and discontinuity (<math>\Delta T^*</math>) of the nematic-isotropic phase transition. (<math>\blacksquare</math>) - <math>\epsilon^*</math> and (<math>\bullet</math>) - <math>\Delta T^*</math> and (b) the critical amplitudes <math>A_T^+</math> and <math>a_T^+</math>. (<math>\blacksquare</math>) - <math>A_T^+</math> and (<math>\bullet</math>) - <math>a_T^+</math>.....</p>	161
<p><b>5.8</b> <math>E(T)</math> vs. temperature near SmA-N transition. Different concentrations are indicated in the figure. Solid lines are fit to Equation (5.8). Dotted arrow indicate <math>T_{SN}</math>. In the inset variation of <math>\alpha</math> with McMillan ratio is shown. TCP indicates the tricritical composition. The solid line is linear fit to the <math>\alpha</math> (<math>\bullet</math>) values.....</p>	164
<p><b>5.9(a)</b> Temperature dependence of density for the concentration <math>x_{5DBT} = 0.301</math>. The dashed arrow denotes the nematic-isotropic (N-I) phase transition temperature (<math>T_{NI}</math>).....</p>	165
<p><b>5.9(b)</b> Molar volume as a function of temperature for <math>x_{5DBT} = 0.301</math>. Solid lines are fit to Equations (5.9) and (5.10). The fitting parameters in both the nematic and isotropic phases are mentioned in the figure. The temperatures <math>T^*</math> and <math>T^{**}</math> are indicated with a star symbol.....</p>	166
<p><b>6.1</b> Phase diagram of the binary system 7CB+ME6O.5. (<math>\square</math>)- nematic-isotropic transition temperature (<math>T_{NI}</math>), (<math>\boxplus</math>)- smectic A-isotropic transition temperature (<math>T_{SI}</math>),</p>	

## List of Figures

and (o)- smectic A-nematic transition temperature ( $T_{SN}$ ). TCP indicates the tricritical points. ....	175
<b>6.2(a-b)</b> Microphotographs (400x magnification) of the textures observed under the polarizing optical microscope for the two mixtures during cooling. (a) Schlieren texture of the nematic phase at 57°C for $x_{7CB} = 0.852$ and (b) Marble type texture of the nematic phase at 58.3°C for $x_{7CB} = 0.127$ .....	176
<b>6.2(c)</b> Microphotograph of the fan shaped texture observed under the polarizing optical microscope during cooling in the SmA phase at 61.8°C for $x_{7CB}=0.424$ ..	177
<b>6.3(a-b)</b> Temperature dependence of density for (a) $x_{7CB} = 0.095$ and (b) $x_{7CB} = 0.127$ in the isotropic, nematic and smectic A phases. Solid arrow and dashed arrow denote the nematic-isotropic (N-I) and smectic A-nematic (SmA-N) transition temperatures respectively.....	178
<b>6.3(c-d)</b> Temperature dependence of density for (c) $x_{7CB} = 0.142$ and (d) $x_{7CB} = 0.164$ in the isotropic, nematic and smectic A phases. Solid arrow and dashed arrow denote the nematic-isotropic (N-I) and smectic A-nematic (SmA-N) transition temperatures respectively.....	179
<b>6.3(e-f)</b> Temperature dependence of density for (e) $x_{7CB} = 0.235$ and (f) $x_{7CB} = 0.352$ in the isotropic, nematic and smectic A phases. Solid arrow, dashed arrow and dotted arrow (for $x_{7CB} = 0.352$ ) denote the nematic-isotropic (N-I), smectic A-nematic (SmA-N) and smectic A-isotropic (SmA-I) transition temperatures respectively.....	180
<b>6.3(g-h)</b> Temperature dependence of density for (g) $x_{7CB} = 0.424$ and (h) $x_{7CB} = 0.477$ in the isotropic and smectic A phases. Dotted arrow denotes the smectic A-isotropic (SmA-I) transition temperature .....	181
<b>6.3(i-j)</b> Temperature dependence of density for (i) $x_{7CB} = 0.535$ and (j) $x_{7CB} = 0.600$ in the isotropic and smectic A phases. Dotted arrow denotes the smectic A-isotropic (SmA-I) transition temperature .....	182
<b>6.3(k-l)</b> Temperature dependence of density for (k) $x_{7CB} = 0.700$ and (l) $x_{7CB} = 0.800$ in the isotropic, nematic and smectic A phases. Solid arrow and dashed	

## List of Figures

---

arrow denote the nematic-isotropic (N-I) and smectic A-nematic (SmA-N) transition temperatures respectively. In the inset the variation of $\rho$ in the vicinity of $T_{NI}$ is shown. $\rho_1$ and $\rho_2$ are density values obtained by extrapolation on either side of $T_{NI}$ .....	183
<b>6.4(a-b)</b> Thermal expansion coefficient for (a) $x_{7CB} = 0.424$ near the smectic A-isotropic (SmA-I) transition, (b) $x_{7CB} = 0.800$ near the nematic-isotropic (N-I) and smectic A-nematic (SmA-N) transitions. Results have been calculated from the pairs of neighbouring density data points .....	185
<b>6.5(a)</b> Logarithm of molar volume as a function of temperature for $x_{7CB} = 0.095$ in the vicinity of nematic-isotropic (N-I) phase transition. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures $T^*$ and $T^{**}$ are indicated with a star symbol ( $\star$ ). Vertical arrow denotes the N-I transition temperature ( $T_{NI}$ ).....	187
<b>6.5(b-c)</b> Logarithm of molar volume as a function of temperature for (b) $x_{7CB} = 0.127$ and (c) $x_{7CB} = 0.142$ in the vicinity of nematic-isotropic (N-I) phase transition. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures $T^*$ and $T^{**}$ are indicated with a star symbol ( $\star$ ). Vertical arrow denotes the N-I transition temperature ( $T_{NI}$ ).....	188
<b>6.5(d-e)</b> Logarithm of molar volume as a function of temperature for (d) $x_{7CB} = 0.164$ and (e) $x_{7CB} = 0.235$ in the vicinity of nematic-isotropic (N-I) transition. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures $T^*$ and $T^{**}$ are indicated with a star symbol ( $\star$ ). Vertical arrow denotes the N-I transition temperature ( $T_{NI}$ ).....	189
<b>6.5(f-g)</b> Logarithm of molar volume as a function of temperature for (f) $x_{7CB} = 0.352$ and (g) $x_{7CB} = 0.424$ in the vicinity SmA-I transition. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures $T^*$ and $T^{**}$ are indicated with a star symbol ( $\star$ ). Dotted arrow denotes the SmA-I transition temperature ( $T_{SI}$ ) .....	190

## List of Figures

- 6.5(h-i)** Logarithm of molar volume as a function of temperature for (h)  $x_{7CB} = 0.477$  and (i)  $x_{7CB} = 0.535$  in the vicinity of smectic A-isotropic (SmA-I) transition. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures  $T^*$  and  $T^{**}$  are indicated with a star symbol ( $\star$ ). Dotted arrow denotes the SmA-I transition temperature ( $T_{SI}$ ) ..... 191
- 6.5(j-k)** Logarithm of molar volume as a function of temperature for (j)  $x_{7CB} = 0.600$  and (k)  $x_{7CB} = 0.700$  in the vicinity of SmA-I and N-I transitions respectively. Solid lines represent the fit according to Equation (6.1) and (6.2). The temperatures  $T^*$  and  $T^{**}$  are indicated with a star symbol ( $\star$ ). Solid and dotted arrows denote the N-I transition and SmA-I transition temperatures respectively..... 192
- 6.5(l)** Logarithm of molar volume as a function of temperature for  $x_{7CB} = 0.800$  in the vicinity of nematic-isotropic (N-I) transition along with the fittings according to Equation (6.1) and (6.2). The temperatures  $T^*$  and  $T^{**}$  are indicated with a star symbol ( $\star$ ). Solid arrow denotes the N-I transition temperature ( $T_{NI}$ ) ..... 193
- 6.6**  $R(T)$  as a function of temperature for some mixtures in the vicinity of SmA-N transition. Solid lines represent the fit according to Equation (6.4). For better representation  $R(T)$  data for the following three concentrations  $x_{7CB} = 0.095, 0.127$  and  $0.700$  has been shifted by  $-8K, -5K$  and  $+3K$  respectively. In the inset variation of thermal expansion coefficient for  $x_{7CB} = 0.800$  is illustrated. .... 198
- 6.7 (a)** Birefringence as a function of temperature for three concentrations. Dashed lower head arrow denotes  $T_{SN}$ . **(b)**  $Q(T)$  as a function of temperature near SmA-N transition. Solid lines represent the fit according to Equation (3.4). For better representation  $Q(T)$  data for the following three concentrations  $x_{7CB} = 0.095, 0.127$  and  $0.700$  has been shifted by  $-8K, -5K$  and  $+3K$  respectively..... 200
- 6.8** Variation of critical exponent with McMillan ratio for some concentrations of the binary system 7CB+ME6O.5 obtained from two sets of measurements *i.e.* density and birefringence data. ( $\bullet$  or  $\bullet$ ) represents the  $\alpha$  values averaged over the

## List of Figures

two values obtained from fitting above and below the transition temperature $T_{SN}$ . Solid line represents a polynomial fit to the $\alpha$ values.....	201
<b>7.1</b> Phase diagram of the binary system of 4DBT+11OCB. $x_{4DBT}$ is the mole fraction of 4DBT. (o)-nematic to isotropic (or smectic A to isotropic) phase transition temperature; (■)-smectic A to nematic phase transition temperature.....	216
<b>7.2(a)</b> Refractive indices ( $n_e$ , $n_o$ ) and birefringence ( $\Delta n = n_e - n_o$ ) as a function of temperature for $x_{4DBT} = 0.208$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines are fit to Equation (7.3).....	217
<b>7.2(b-c)</b> Refractive indices ( $n_e$ , $n_o$ ) and birefringence ( $\Delta n = n_e - n_o$ ) as a function of temperature for (b) $x_{4DBT} = 0.291$ and (c) $x_{4DBT} = 0.393$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines are fit to Equation (7.3).....	218
<b>7.2(d-e)</b> Refractive indices ( $n_e$ , $n_o$ ) and birefringence ( $\Delta n = n_e - n_o$ ) as a function of temperature for (d) $x_{4DBT} = 0.491$ and (e) $x_{4DBT} = 0.592$ . Solid arrow denotes nematic-isotropic transition ( $T_{NI}$ ) temperature. The solid lines are fit to Equation (7.3) .....	219
<b>7.2(f-g)</b> Refractive indices ( $n_e$ , $n_o$ ) and birefringence ( $\Delta n = n_e - n_o$ ) as a function of temperature for (f) $x_{4DBT} = 0.665$ and (g) $x_{4DBT} = 0.707$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines are fit to Equation (7.3).....	220
<b>7.2(h-i)</b> Refractive indices ( $n_e$ , $n_o$ ) and birefringence ( $\Delta n = n_e - n_o$ ) as a function of temperature for (h) $x_{4DBT} = 0.787$ and (i) $x_{4DBT} = 0.884$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid lines are fit to Equation (7.3).....	221
<b>7.3(a-b)</b> Temperature dependence of order parameter ( $\langle P_2 \rangle$ ) for (a) $x_{4DBT} = 0.208$ and (b) $x_{4DBT} = 0.291$ . Solid arrow and dashed arrow denote the nematic-isotropic	

## List of Figures

( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid line represents $\langle P_2 \rangle$ from the McMillan's theory.....	223
<b>7.3(c-d)</b> Temperature dependence of order parameter ( $\langle P_2 \rangle$ ) for (c) $x_{4DBT} = 0.393$ and (d) $x_{4DBT} = 0.491$ . Solid arrow denotes the nematic-isotropic ( $T_{NI}$ ) transition temperature. The solid line represents $\langle P_2 \rangle$ from the Maier-Saupe theory.....	224
<b>7.3(e-f)</b> Temperature dependence of order parameter ( $\langle P_2 \rangle$ ) for (e) $x_{4DBT} = 0.592$ and (f) $x_{4DBT} = 0.665$ . Solid arrow denotes the nematic-isotropic ( $T_{NI}$ ) transition temperature. The solid line represents $\langle P_2 \rangle$ from the Maier-Saupe theory.....	225
<b>7.3(g-h)</b> Temperature dependence of order parameter ( $\langle P_2 \rangle$ ) for (g) $x_{4DBT} = 0.707$ and (h) $x_{4DBT} = 0.787$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid line represents $\langle P_2 \rangle$ from the McMillan's theory.....	226
<b>7.3(i)</b> Temperature dependence of order parameter ( $\langle P_2 \rangle$ ) for $x_{4DBT} = 0.884$ . Solid arrow and dashed arrow denote the nematic-isotropic ( $T_{NI}$ ) and smectic A-nematic ( $T_{SN}$ ) transition temperatures respectively. The solid line represents $\langle P_2 \rangle$ from the McMillan's theory.....	227
<b>7.4(a)</b> Temperature variation of the dielectric parameters for $x_{4DBT} = 0.208$ . Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A–nematic ( $T_{SN}$ ) phase transition temperatures respectively.....	230
<b>7.4(b-c)</b> Temperature variation of the dielectric parameters for (b) $x_{4DBT} = 0.291$ and (c) $x_{4DBT} = 0.393$ . Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A–nematic ( $T_{SN}$ ) phase transition temperatures respectively..	231
<b>7.4(d-e)</b> Temperature variation of the dielectric parameters for (d) $x_{4DBT} = 0.491$ and (e) $x_{4DBT} = 0.592$ . Solid arrow indicates nematic–isotropic phase transition temperature ( $T_{NI}$ ).....	232
<b>7.4(f-g)</b> Temperature variation of the dielectric parameters for (f) $x_{4DBT} = 0.665$ and (g) $x_{4DBT} = 0.707$ . Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A–nematic ( $T_{SN}$ ) phase transition temperatures respectively..	233

## List of Figures

<b>7.4(h-i)</b> Temperature variation of the dielectric parameters for (h) $x_{4DBT} = 0.787$ and (i) $x_{4DBT} = 0.884$ . Solid arrow and dashed arrow indicate nematic–isotropic ( $T_{NI}$ ) and smectic A–nematic ( $T_{SN}$ ) phase transition temperatures respectively..	234
<b>7.5</b> Measured dielectric anisotropy ( $\Delta\epsilon$ ) as a function of temperature for the mixture $x_{4DBT} = 0.665$ . Dotted lower head arrow indicates the temperature $T^{**}$ . The inset shows a double logarithmic plot of $(\Delta\epsilon - \Delta\epsilon^{**})$ vs $(T^{**} - T)$ . The dashed lines are fit to the data points according to the Equation (7.4) .....	236
<b>8.1</b> Voltage dependent capacitance of the planar aligned LC cell at different temperatures for $x_{4DBT} = 0.393$ . The enhancement of the threshold voltage with lowering the temperature is clearly observed. Different temperatures in °C are mentioned in the figure .....	246
<b>8.2</b> Variation of the splay elastic constant ( $K_{11}$ ) of different mixtures as a function of shifted temperature. Dashed upper head arrow denotes SmA-N transition temperature ( $T_{SN}$ ) for the respective mixtures. Different mole fractions of 4DBT are indicated in the figure.....	247
<b>8.3(a-b)</b> Variation of $K_{11}$ as a function of $(\langle P_2 \rangle)^2$ at the different concentrations of 4DBT. Solid lines are linear fit to the data points. Different mole fractions of 4DBT are indicated in the figure.....	248
<b>8.4</b> Relaxation time ( $\tau_0$ ) as a function of shifted temperature for the studied mixtures. Dashed upper head arrow denotes SmA-N transition temperature ( $T_{SN}$ ) for the respective mixtures. Different mole fractions of 4DBT are indicated in the figure .....	250
<b>8.5</b> Rotational viscosity ( $\gamma_1$ ) as a function of shifted temperature for the different mixtures. Dashed upper head arrow denotes smectic A-nematic transition ( $T_{SN}$ ) temperature for the respective mixtures. Different mole fractions of 4DBT are indicated in the figure. Dashed lines are fit to Equation 8.4 .....	251
<b>8.6</b> Variation of $\ln(\gamma_1 / \langle P_2 \rangle)$ with $1/T$ for the concentration $x_{4DBT} = 0.665$ . The solid line is linear fit to the data points .....	253

## **List of Figures**

---

<b>8.7</b>	Activation energy as a function of concentration of 4DBT .....	254
------------	--	-----