

LIST OF SCHEMES

<u>CHAPTERS</u>	<u>SCHEMES</u>	<u>PAGE NO.</u>
<i>Chapter-IV</i>	Scheme 1: The molecular structures of Tyrosine, Tryptophan and Ascorbic acid.	342-343
	Scheme 2. Diagrammatic representation of the Probable Geometrical Configurations involving tyrosine molecules surrounded by ascorbic acid molecules involving three H-bonding.	
	Scheme3. Diagrammatic representation of the Probable Geometrical Configurations involving tryptophan molecules surrounded by ascorbic acid molecules involving two H-bonding.	
<i>Chapter-V</i>	Scheme 1. Three Dimensional Representation of Solvent Cage	368-370
	Scheme 2. Array of Limiting Apparent Molar Volumes at a definite concentration.	
	Scheme 3. Array of Molecular Interactions.	
	Scheme 4. Different possible ionic group interactions in aqueous solutions of ascorbic acid in presence of LiX [X = Cl, Br, I].	
	Scheme 5. Ion-solvent interaction	
<i>Chapter-VI</i>	Scheme 1 Structure of Cyclodextrin and Crown molecules.	397-400
	Scheme 2 Molecular structure of [EMPyrr] Br.	
	Scheme 3 (a) Schematic representation of Inclusion complexation of [EMPyrr] Br with β - Cyclodextrin.	

	(b) Incorporation of [EMPyrr] Br into the cavity of truncated conical structure of β -cyclodextrin showing the stereospecific complexation.	
	Scheme 4 Schematic representation of encapsulation interaction of [EMPyrr] Br with 18 C-6.	
	Scheme 5 Schematic representation of solvation of ions screening Solute – Solvent Interactions.	
	(a) Incorporation of IL into the hollow space of β -CD.	
	(b) Encapsulation of [EMPyrr] Br with 18 C-6.	
<i>Chapter-VII</i>	Scheme 1: Molecular structure of ascorbic acid	418, 419, 423, 425
	Scheme 2: Pictorial representation of Oxalate ion .	
	Scheme 3: Pictorial representation of the four oxalate salts.	
	Scheme 4. Keto-enol tautomerism of ascorbic acid.	
	Scheme 5. The stability of the conjugate base of ascorbate ion.	
	Scheme 6: p ^H ranges of interactions of aqueous ascorbic acid solutions	
	Scheme 7: Probable interaction pattern of the studied system.	
	Scheme 8 : Probable mechanism of the studied system.	

	Scheme 9. The schematic representation of solute–solvent interaction, for the studied OXALATES in aqueous vitamins binary mixtures.	
<i>Chapter-VIII</i>	Scheme 1. Optimised geometry would involve a central Li ⁺ ion tetrahedrally surrounded by four benzonitrile molecules.	464-465
	Scheme 2. Diagrammatic representation of the Probable Geometrical Configurations of the complex I(PhcN) ₂ ⁻ .	
<i>Chapter-IX</i>	Scheme 1. Schematic representation of mechanism for the formation of 1: 1 inclusion complex of Tertiary-Leucine with both α and β -cyclodextrin.	466
	Scheme2. (a) Proposed schemes of different possibilities of host-guest ratio for inclusion complexation. (b) Proposed schemes of feasible and restricted inclusion of the guest	

	molecule into the host molecule.	
<i>Chapter-X</i>	Scheme 1. Schematic representation of mechanism for the formation of 1: 1 inclusion complex of nicotinamide with β -cyclodextrin.	512
	Scheme 2. Schematic representation of mechanism for the formation of 1: 1 inclusion complex of pyridoxine with β -cyclodextrin.	