

LIST OF TABLES

CHAPTERS	TABLES	PAGE NO.
<i>Chapter IV</i>	<p>Table 1. Values of density (ρ), viscosity (η), at 298.15, 303.15, 308.15 K and refractive index (n_D) at 298.15 K in different mass fraction (w_1) of aqueous Ascorbic Acid.</p> <p>Table 2. Experimental values of density (ρ) and viscosity (η) at 298.15 K, 303.15 K, 308.15 K and Refractive index (n_D) at 298.15 K of amino acids in different mass fraction of aqueous AA (w_1).</p> <p>Table 3. Molality (m), apparent molar volume (ϕ_V) and $(\eta_r - 1)/\sqrt{m}$ at 298.15 K, 303.15 K, 308.15 K and molar refraction (R) at 298.15 K of amino acids in different mass fraction of aqueous AA (w_1).</p> <p>Table 4. Partial molar volumes (ϕ_V^0), experimental slopes (S_V^*), viscosity A- and B-coefficients of amino acids in different mass fraction of aqueous AA (w_1) at 298.15 K, 303.15 K and 308.15 K respectively.</p> <p>Table 5. Values of empirical coefficients (a_0, a_1, and a_2) of Equation 4 for amino acids in different mass fraction of aqueous vitamin (w_1) at 298.15K to 308.15K respectively.</p> <p>Table 6. Limiting apparent molal expansibilities (ϕ_E^0) for amino acids in different mass fraction of aqueous vitamin (w_1) at 298.15K to 308.15K respectively.</p> <p>Table 7. Values of dB/dT for the amino acids in different mass fraction of aqueous vitamin C (w_1) at 298.15 to 308.15K respectively.</p> <p>Table 8: Properties of Amino acids –VitaminC Complexes calculated by the UB3LYP/6-31+g(d) method.</p>	326-336
<i>Chapter V</i>	<p>TABLE 1: Values of empirical coefficients (a_0, a_1, and a_2) of Equation 8 for Lithium Halides in co-solute (ascorbic acid, water) at 298.15 K, 303.15K, 308.15 K.</p> <p>TABLE 2. Limiting apparent molal expansibilities (ϕ_E) for Lithium salts in co-solute (ascorbic acid and water) at 298.15 K to 308.15 K respectively.</p> <p>TABLE 3: Refractive index (n_D), Molar refraction (R_M), Limiting molar refraction (R_M^0) of LiCl, LiBr, LiI in aqueous solution of ascorbic acid .</p>	360-362

<p><i>Chapter VI</i></p>	<p>Table 1: Experimental values of density (ρ), viscosity (η), refractive index (n_D), surface tension (γ) in deferent mass fraction of aqueous 18 C-6 and β-cyclodextrin mixtures at 298.15 K^a.</p> <p>Table 2: Experimental values of density (ρ), viscosity (η), refractive index (n_D), of ionic liquid in different mass fraction of aqueous 18 C-6 and β-CD mixtures at 298.15 K^a.</p> <p>Table 3: Apparent molar volume (ϕ_V), $(\eta_r-1)/\sqrt{c}$, and molar refraction (R_M) of selected ionic liquid in different mass fraction of aqueous 18 C-6 and β-CD mixtures at 298.15 K^a.</p> <p>Table 4: Limiting Apparent molar volume (ϕ_V^0), (S_V^*), Viscosity B- and A- co-efficients and Limiting Molar Refraction (R_M^0) of selected ionic liquid in different mass fractions of aqueous 18 C-6 and β-cyclodextrin mixtures at 298.15 K^a.</p> <p>Table 5: Surface tension (γ) and the corresponding concentration (c) of ionic liquid in 0.005(m) mass fraction of aqueous 18 C-6 and β- cyclodextrin mixtures at 298.15 K^a.</p> <p>Table 6(a): Molar conductance (Λ_m) and the corresponding concentration (c) of ionic liquid in different mass fractions of aqueous 18 C-6 and β- cyclodextrin mixtures at 298.15 K^a.</p> <p>Table 6(b): Binding constants [(K_b) and (K_b')] $\times 10^{-3} \text{ M}^{-1}$ of IL: β-CD and IL: 18 C-6 complexes.</p> <p>Table 7: Estimated vibrational frequencies for [β - CD : IL] and [18 C-6 : IL] Complex formation.</p> <p>Table 8: Analysis of 1H NMR data of [β-CD + EMPyrr] in D₂O environment.</p> <p>Table 9: Analysis of 1H NMR data of [18 C-6 + EMPyrr] in D₂O environment.</p> <p>Table 10: The observed experimental peaks at different m/ z with corresponding ions for the solid Inclusion and encapsulated complexes.</p>	<p>401-407</p>
<p><i>Chapter VII</i></p>	<p>Table 1. Sample description and calculating mass-fractions of the samples.</p> <p>Table 2. Solubility test.</p> <p>Table 3. Experimental values of densities (ρ), refractive Index (n_D) of Oxalates in different mass fraction of aqueous ascorbic acid at temperatures, T = (298.15K) and at</p>	<p>420, 424, 426, 427, 428, 429, 430</p>

	<p>ambient pressure.</p> <p>Table 4: Measurement of viscosity (η) in centipoises at 298.15 K, 303.15K, 308.15 K respectively.</p> <p>Table 5: Examining viscosity A, B coefficients of Lithium, Sodium, Potassium, Ammonium Oxalates at 298.15 K to 308.15 K respectively.</p> <p>Table 6: Data of $(\eta / \eta^{0-1})/m^{1/2}$ obtained from the Viscosity calculations of Oxalates at different temperatures:</p> <p>Table 7: Molality (m), apparent molar volume (ϕ_v), molar refraction (R) of Lithium Oxalate, Sodium Oxalate, Potassium Oxalate, Ammonium Oxalate in aqueous vitamin -C Solution in different mass fraction at room temperature.</p> <p>Table 8. Limiting apparent molar volumes (ϕ_v^0), experimental slopes (Sv^*) of Lithium Oxalate, Sodium Oxalate, Potassium Oxalate, Ammonium Oxalate in aqueous Ascorbic acid at 298.15 K.</p>	
<i>Chapter VIII</i>	<p>Table 1: Properties of Li+...PhcN Complexes calculated by the UB3LYP/6-31+g(d) method</p> <p>Table 2: Estimated vibrational frequencies for Li+...PhcN complexes</p> <p>Table 3: Properties of I-...PhcN Complexes calculated by the UB3LYP/midix method</p> <p>Table 4. Estimated vibrational frequencies for I-...PhcN Complexes</p> <p>Table 5. Molar conductance (Λ) and molar concentration of lithium iodide salts in benzonitrile solution at various temperatures</p> <p>Table 6. Equivalent conductance at infinite dilution at various temperatures of Lithium iodide in benzonitrile solution</p>	451-453
<i>Chapter IX</i>	<p>Table 1: Values of surface tension (γ) at the break point with corresponding concentrations of cyclodextrins and amino acid and values of conductivity (k) at the break point with corresponding concentrations of cyclodextrins and amino acids at 298.15 K^a.</p> <p>Table 2: Contributions of zwitter ionic group (NH_3^+), (COO^-); (CH) and side group (R) to the limiting apparent molar volume (ϕ_v^0) and viscosity B-coefficient for the amino acid in different mass fraction of aqueous α and β-cyclodextrin respectively at 298.15K^a, 303.15K^a, 308.15K^a.</p>	483-498

Table 3: Hydration number (n_H), solvation number (S_n) of the amino acid at different mass fraction of aqueous α - and β -cyclodextrin respectively and binding constant (K_b) of amino acid-cyclodextrin inclusion complexes at 298.15 K^a, 303.15K^a, 308.15K^a.

Table S1. Experimental values of density (ρ), viscosity (η), refractive index (n_D) and pH of different mass fractions of aqueous α , β -cyclodextrin mixtures in tertiary leucine at 298.15 K^a.

Table S2. Experimental values of density (ρ), viscosity (η), refractive index (n_D) and pH of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 298.15 K^a.

Table S3. Experimental values of density (ρ), viscosity (η), refractive index (n_D) and pH of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 303.15 K^a.

Table S4. Experimental values of density (ρ), viscosity (η), refractive index (n_D) and pH of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 308.15 K^a.

Table S5. Data for surface tension and conductivity study of aqueous Tert-Leucine – α -CD system at 298.15K^a.

Table S6. Data for surface tension and conductivity study of aqueous Tert-Leucine – β -CD system at 298.15K^a.

Table S7. Apparent molar volume (ϕ_V), $(\eta_r-1)/\sqrt{m}$, molar refraction (R_M) of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 298.15 K^a.

Table S8. Apparent molar volume (ϕ_V), $(\eta_r-1)/\sqrt{m}$, molar refraction (R_M) of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 303.15 K^a.

Table S9. Apparent molar volume (ϕ_V), $(\eta_r-1)/\sqrt{m}$, molar refraction (R_M) of selected amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 308.15 K^a.

Table S10. Limiting apparent molar volume (ϕ_V^0), experimental slope (S_V^*), viscosity A and B -coefficient and limiting molar refraction (R_M^0) of amino acid in different mass fractions of aqueous α and β -cyclodextrin mixtures at 298.15 K^a.

Table S11. Limiting apparent molar volume (ϕ_V^0),

	<p>experimental slope (S_V^*), viscosity A and B-coefficient and limiting molar refraction (R_M^0) of amino acid in different mass fractions of aqueous α and β-cyclodextrin mixtures at 303.15 K^a.</p> <p>Table S12. Limiting apparent molar volume (ϕ_V^0), experimental slope (S_V^*), viscosity A and B-coefficient and limiting molar refraction (R_M^0) of amino acid in different mass fractions of aqueous α and β-cyclodextrin mixtures at 308.15 K^a.</p> <p>Table S13: Data for the Benesi-Hildebrand double reciprocal plot performed by ¹H NMR spectroscopy for aqueous Tert -Leucine-α-CD system at 298 K (concentration of Tert-Leucine = 500 μM).</p> <p>Table S14: Data for the Benesi-Hildebrand double reciprocal plot performed by ¹H NMR spectroscopy for aqueous Tert -Leucine-β-CD system at 298 K (concentration of Tert-Leucine = 500 μM).</p>	
<p><i>Chapter X</i></p>	<p>TABLE 1</p> <p>Provenance and mass fraction purity of chemical samples</p> <p>TABLE 2</p> <p>Density (ρ), ^a Limiting apparent molar volume of solute (ϕ_V^0), experimental slopes (S_V), absolute viscosity (η) and relative viscosity(η_r) of an aqueous solutions of pyridoxine at different temperatures T = (298.15, 303.15 and 308.15) K at 10⁵ N. m⁻².</p> <p>TABLE 3</p> <p>Density (ρ), ^a Limiting apparent molar volume of solute (ϕ_V^0), experimental slopes (S_V), absolute viscosity (η) and relative viscosity(η_r) of an aqueous solutions of nicotinamide at different temperatures T = (298.15, 303.15 and 308.15) K at 10⁵ N. m⁻².</p> <p>TABLE 4</p> <p>Standard deviation (u) in (ϕ_V^0), absolute average deviation (AAD), in (ϕ_V^0), limiting apparent molal expansivity of solute (E_ϕ^0) at T/K = (298.15 to 308.15), $\delta^2\phi_V^0/\delta T^2$ and coefficient of thermal expansion (α^*) at T = 298.15 K of both aqueous β-CD binary systems with mass fraction ($w=0.005$).</p> <p>TABLE 5</p> <p>Experimental values of Falkenhagen coefficient A, Jones-Dole coefficient B and D viscosity coefficients, temperature coefficient of B, i.e. (dB/dT), Gibbs free energies of</p>	<p>538-542</p>

	<p>activation per mole of solvent ($\Delta\mu_1^{0\#}$) and per mole of solute ($\Delta\mu_2^{0\#}$), entropy of activation per mole of solute ($\Delta S_2^{0\#}$), enthalpy of activation per mole of solute ($\Delta H_2^{0\#}$) at and hydration number (n_H), for both aqueous binary systems at $T/K = (298.15, 303.15 \text{ and } 308.15)$, ($w = 0.005$).</p> <p>Table 6</p> <p>Estimated vibrational frequencies for [β - CD: nicotinamide] Complex formation</p> <p>Table 7</p> <p>Estimated vibrational frequencies for [β - CD: pyridoxine] Complex formation.</p>	
--	--	--