

## LIST OF FIGURES

FIGURES	PAGE NO.
<b>CHAPTER IV</b>	
<b>Figure 1.</b> Job plot of DEPM- $\beta$ -Cyd systems at 298.15K in neutral (pink, ●) and alkaline (blue, ■) medium.	94
<b>Figure 2.</b> Adsorption spectra of DEPM ( $2 \times 10^{-4}$ M) in (a) pH=7.0 $\pm$ 0.5 and (b) pH=9.2 $\pm$ 0.5 at different concentration of $\beta$ -Cyd (1) absence of $\beta$ -Cyd, (2) 0.002M, (3) 0.004M, (4) 0.006M, (5) 0.008M, (6) 0.010M, (7)0.012M.	96
<b>Figure 3.</b> Benesi-Hildebrand plot of $1/A-A_0$ vs. $1/[\beta\text{-CD}]$ for DEPM in pH=7.0 $\pm$ 0.5 and pH=7.0 $\pm$ 0.5 at 298.15K.	98
<b>Figure4.</b> $^1\text{H}$ NMR spectra of (a) $\beta$ -Cyd, (b) 1:1 molar ratio of $\beta$ -CD:DEPM IC in $\text{D}_2\text{O}$ and $^1\text{H}$ NMR spectra of (c) DEPM in $\text{DMSO-D}_6$ at 298.15 K.	101
<b>Figure 5.</b> FT-IR spectra of (a) $\beta$ -Cyd (in KBr), (b) DEPM and (c) DEPM: $\beta$ -Cyd (1:1 molar ratio) solid complex (in KBr).	104
<b>Figure 6.</b> Powder X-ray diffraction pattern of (a) $\beta$ -Cyd and (b) DEPM: $\beta$ -Cyd (1:1 molar ratio) inclusion complex.	105
<b>Figure7.</b> Scanning electron microscope (SEM) morphology of (a) $\beta$ -Cyd and (b) DEPM: $\beta$ -Cyd (1:1 molar ratio) inclusion complex.	106
<b>CHAPTER V</b>	
<b>Figure 1.</b> Molar conductance vs mole ratio of $[\text{H}]/[\text{EMPyrr}]^+$ of 18C6+IL at 293.15K (pink), 298.15K (blue) and 303.15K (orange).	114
<b>Figure 2.</b> Molar conductance Vs mole ratio of $[\text{H}]/[\text{EMPyrr}]^+$ of DB18C6+IL at 293.15K (green), 298.15K (red) and 303.15K (violet).	114
<b>Figure 3.</b> $\log K_f$ Vs $1000/T$ for 18C6:IL (■) and DB18C6:IL (●) complexation.	117

<b>Figure 4.</b> $^1\text{H}$ NMR spectra of (a) 1-ethyl-1 methylpyrrolidinium hexafluorophosphate and (b) 1:1 molar ratio of 18C6:[EMPyrr] $^+$ in $\text{D}_2\text{O}$ at 298.15K.	120
<b>Figure 5.</b> $^1\text{H}$ NMR spectra of (a) 1-ethyl-1 methylpyrrolidinium hexafluorophosphate and (b) 1:1 molar ratio of DB18C6:[EMPyrr] $^+$ in $\text{D}_2\text{O}$ at 298.15K.	120
<b>Figure 6.</b> Electrospray mass spectra of equimolecular mixture of IL with (a) 18-crown-6 and (b) dibenzo 18-crown-6.	123
<b>Figure 7.</b> IR spectra of 18-crown-6 (blue) and the resulting complex I (18C6:[EMPyrr] $^+$ ) (red).	124
<b>Figure 8.</b> IR spectra of Dibenzo-18-crown-6 (blue) and the resulting complex II (DB18C6:[EMPyrr] $^+$ ) (violet).	125
<b>CHAPTER VI</b>	
<b>Figure 1.</b> Molar conductivity of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] in aqueous solution with L-tyr and (b) [MOIM]Cl in aqueous solution with L-tyr at three different temperatures.	133
<b>Figure 2.</b> Molar conductivity of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] in aqueous solution with L-phe and (b) [MOIM]Cl in aqueous solution with L-phe at three different temperatures.	133
<b>Figure 3.</b> Adsorption spectra of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b) [MOIM]Cl at different concentration of L-tyrosine (1) absence of L-tyrosine, (2) 0.0004 M, (3) 0.0008 M, (4) 0.0012 M, (5) 0.0016 M, (6) 0.0018 M, (7)0.002 M respectively.	136
<b>Figure 4.</b> Adsorption spectra of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b)[MOIM]Cl at different concentration of L-phenylalanine (1) absence of L-phenylalanine, (2) 0.0004 M, (3) 0.0008 M, (4) 0.0012 M, (5) 0.0016 M, (6) 0.0018 M, (7)0.002 M respectively.	136
<b>Figure 5.</b> Fluorescence spectra of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b) [MOIM]Cl at different concentration of L-tyrosine (1) absence of L-tyrosine, (2)	137

0.0003 M, (3) 0.0006 M, (4) 0.0009 M, (5) 0.0012 M, (6) 0.0015 M, (7)0.0018 M respectively.	
<b>Figure 6.</b> Fluorescence spectra of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b) [MOIM]Cl at different concentration of L-phenylalanine (1) absence of L-phenylalanine, (2) 0.0003 M, (3) 0.0006 M, (4) 0.0009 M, (5) 0.0012 M, (6) 0.0015 M, (7)0.0018 M respectively.	137
<b>Figure 7.</b> Benesi-Hildebrand plot of $1/\Delta A$ vs. $1/[AA]$ in UV-vis spectroscopy for (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b) MOIM]Cl in L-tyr and for (c)[BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (d) MOIM]Cl in L-phe at 298.15K <sup>a</sup> .	142
<b>Figure 8.</b> Benesi-Hildebrand plot of $1/\Delta A$ vs. $1/[AA]$ in fluorescence spectroscopy for (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (b) MOIM]Cl in L-tyr and for (c)[BMIM][C <sub>8</sub> SO <sub>4</sub> ] and (d) MOIM]Cl in L-phe at 298.15K <sup>a</sup> .	142
<b>Figure 9.</b> <sup>1</sup> H NMR spectra of (a) [BMIM][C <sub>8</sub> SO <sub>4</sub> ], (b) ([BMIM][C <sub>8</sub> SO <sub>4</sub> ]+ L-tyr) system and (c) ([BMIM][C <sub>8</sub> SO <sub>4</sub> ] + L-phe) system in D <sub>2</sub> O at 298.15K.	144
<b>Figure 10.</b> <sup>1</sup> H NMR spectra of (a) [MOIM]Cl, (b) ([MOIM]Cl+ L-tyr) system and (c) ([MOIM]Cl + L-phe) system in D <sub>2</sub> O at 298.15K.	144
<b>CHAPTER VII</b>	
<b>Figure 1.</b> Job plot of NTHCL/ $\beta$ -Cyd systems at 298.15K.	153
<b>Figure 2.</b> Fluorescence spectra of NTHCL in different $\beta$ -Cyd concentrations ( $\mu$ M): (1) 0, (2) 20, (3)30, (4) 40, (5) 50, (6) 60, (7) 70 and (8) 80.	155
<b>Figure 3.</b> Benesi-Hildebrand double reciprocal plot of $1/I-I_0$ vs. $1/[\beta$ -Cyd] for NTHCL in (a) UV-absorption and (b) fluorescence at 298.15K.	158
<b>Figure 4.</b> <sup>1</sup> H NMR spectra of (a) $\beta$ -Cyd, (b) 1:1 molar ratio of NTHCL/ $\beta$ -Cyd inclusion complex and (c) NTHCL in D <sub>2</sub> O at 298.15 K.	159
<b>Figure 5.</b> 2D ROESY spectra of solid inclusion complex of NTHCL and $\beta$ -CD in D <sub>2</sub> O (correlation signals are marked by green circles).	161

<b>Figure 6.</b> . <sup>13</sup> CNMR spectra of (a) β- Cyd, (b) 1:1 molar ratio of solid NTHCL/β-Cyd inclusion complex and (c) NTHCL in D <sub>2</sub> O at 298.15 K.	162
<b>Figure 7.</b> FT-IR spectra of (a) β- Cyd (in KBr), (b) NTHCL and (c) NTHCL/β- Cyd (1:1 molar ratio) solid complex (in KBr).	164
<b>Figure 8.</b> Powder X-ray diffraction pattern of (a) β-Cyd, (b) NTHCL and (C) NTHCL/β-Cyd (1:1 molar ratio) inclusion complex.	166
<b>CHAPTER VIII</b>	
<b>Figure 1.</b> Job plot of (a) PMO/α- CD and (b) PMO/β-CD systems at 298.15K.	173
<b>Figure 2:</b> Benesi-Hildebrand double reciprocal plot for the effect of α- Cyd on the absorbance of PMO (308 nm) at different temperatures.	177
<b>Figure 3:</b> Benesi-Hildebrand double reciprocal plot for the effect of β- Cyd on the absorbance of PMO (308 nm) at different temperatures.	177
<b>Figure 4.</b> Plot of ln <i>K</i> <sub>a</sub> vs. 1/ <i>T</i> for the interaction of CDP with α-CD (■) and β-CD (■).	178
<b>Figure 5.</b> FTIR spectra of (a) PMO, (b) α-cyd and (c) PMO/α-cyd (1:1 molar ratio) solid inclusion complex in KBr.	179
<b>Figure 6.</b> FTIR spectra of (a) PMO, (b) β-cyd and (c) PMO/β-cyd (1:1 molar ratio) solid inclusion complex in KBr.	181
<b>Figure 7.</b> ESI mass spectra of (a) PMO/α-CD inclusion complex and (b) PMO/β-CD inclusion complex.	182
<b>Figure 8.</b> Powder X-ray diffraction pattern of (a) α-CD, (b) PMO/α-CD (1:1 molar ratio), (c) β-CD and (d) PMO/ β-CD (1:1 molar ratio) inclusion complex.	183
<b>Figure 9.</b> TGA profiles of (a) α-cyd ; PMO/α-cyd and (b) β-cyd; PMO/β-cyd inclusion complex systems.	184
<b>CHAPTER IX</b>	
<b>Figure 1.</b> Variation of surface tension of aqueous (A) L-Methionine solution, (B) L-Proline solution, and (C) L-Glutamine solution	193

respectively with increasing concentration of $\beta$ -CD.	
<b>Figure 2.</b> Plot of limiting molar volume ( $\phi_{v^{\circ}}$ ) against mass fraction ( $w$ ) of $\beta$ -CD for L-Methionine (blue), L-Proline (brown) and L-Glutamine (green) respectively at 298.15 K <sup>a</sup> ( <sup>a</sup> Standard uncertainties $u$ is: $u(T)=0.01$ K).	196
<b>Figure 3.</b> <sup>1</sup> H NMR spectra of (a) $\beta$ -CD, (b) L-Methionine and (c) 1:1 molar ratio of $\beta$ -CD & L-Methionine in D <sub>2</sub> O at 298.15 K <sup>a</sup> ( <sup>a</sup> Standard uncertainties $u$ is: $u(T)=0.01$ K).	205
<b>Figure 4.</b> <sup>1</sup> H NMR spectra of (a) $\beta$ -CD, (b) L-Proline and (c) 1:1 M ratio of $\beta$ -CD & L-Proline in D <sub>2</sub> O at 298.15 K <sup>a</sup> ( <sup>a</sup> Standard uncertainties $u$ is: $u(T) = 0.01$ K).	206
<b>Figure 5.</b> <sup>1</sup> H NMR spectra of (a) $\beta$ -CD, (b) L-Glutamine and (c) 1:1 molar ratio of $\beta$ -CD & L-Glutamine in D <sub>2</sub> O at 298.15 K.	207
<b>Figure 6.</b> Powder X-ray diffraction pattern of (a) $\beta$ -CD, (b) methionine (Met), (c) proline (Pro), (d) glutamine (Glu), (e) $\beta$ -Cyd: Met, (f) $\beta$ -CD: Pro and (g) $\beta$ -CD: Glu (1:1 molar ratio) inclusion complex.	209