

List of Figures

- Fig. 1.1:** Map of north-east India showing the north-east Indian plains.
- Fig. 1.2:** Virus infected bottle gourd (a), tomato (b) and cucumber (c) fields of north-east Indian plains. Red circles in the photographs indicate the infected plants in the fields.
- Fig. 2.1:** Genome organization of *Potyvirus* showing different proteins coding regions: N-terminal protein (P1 protein), helper component protein (HC-Pro), P3 protein, 6 kDa protein 1 (6K1), cytoplasmic inclusion protein (CI), 6 kDa protein2 (6K2), nuclear inclusion A linked VPg protein (NIa-VPg), nuclear inclusion protein A (NI-a), nuclear inclusion protein B (NI-b), coat protein (CP).
- Fig. 2.2:** Genome organization of *Potexvirus* showing different proteins coding regions: RNA dependent RNA polymerase (RdRp), Triple gene block 1 (TGB1), Triple gene block 2 (TGB2), Triple gene block 3 (TGB3), coat protein (CP).
- Fig. 2.3:** Genome organization of *Emaravirus* showing different protein coding RNA particles: RNA dependent RNA polymerase (RdRp), glycoprotein precursor, nucleocapsid, movement protein and three RNA particles (p5, p6 and p7) of unknown function.
- Fig. 3.1:** (a) Map of northern West Bengal and (b-g) regions of collection of infected plant samples from different districts of North Bengal: (b) Siliguri, (c) Jalpaiguri, (d) Islampur, (e) Malda, (f) Cooch behar and (g) Balurghat. The red pins showing the collection spots.
- Fig. 3.2:** (a) Map of Tripura and (b) region of collection of infected plant samples from Agartala, Tripura. The red pin is showing the collection spot.

- Fig. 3.3:** (a) Map of northern Assam and (b-o) regions of collection of infected plant samples from different districts of Assam: (b) Silchar, (c) Sapotgram, (d) North Salmara, (e) Nayagaon, (f) Bilasipara (g) Talguri, (h) Jagdoi, (i) Anthaibari, (j) Dhaligaon, (k) Dibrugarh, (l) Silchar, (m) Goalpara, (n) Guwahati and (o) Sonitpur. The red pins showing the collection spots.
- Fig. 3.4:** Leaves of the plants used for the management of *Papaya ringspot virus* disease in bottle gourd: (a) *Azadirachta indica*, (b) *Bougainvillea spectabilis*, (c) *Clerodendrum infortunatum*, (d) *Lantana camara* and (e) *Camellia sinensis*.
- Fig. 4.1.A:** Naturally infected papaya (*Carica papaya*) leaf samples collected from different places of West Bengal (WB) and Assam [Pa=Papaya; Isl= Islampur, WB; Blg= Balurghat, WB; Guw= Guwahati, Assam; Ran= Siliguri, WB; Fsd= Siliguri, WB; Lnp= Siliguri, WB; Jpg= Jalpaiguri, WB; An= Anthaibari, Assam and Cob= Cooch behar, WB].
- Fig. 4.1.B:** Naturally infected papaya (*Carica papaya*) leaf samples and papaya fruit with ring spot symptom collected from different places of West Bengal and Assam [Pa= Papaya; PaF= Papaya fruit; NS= North Salmara, Assam; Sa= Sapotgram, Assam; Ng= Nasyagaon, Assam; Gp= Goalpara, Assam and Mld= Malda, WB]
- Fig. 4.2:** Naturally infected common bean (*Phaseolus vulgaris*) leaf samples collected from different places of West Bengal (WB) and Assam [Be= common bean; Dbg= Dibrugarh, Assam; Slg= Siliguri, WB; Cob= Cooch behar, WB; Dg= Dhaligaon; Bs= Bilasipara, Assam; Sr= Srirampur, Assam and Ng= Nayagaon].
- Fig. 4.3:** Naturally infected field bean (*Dolichos lablab*) leaf samples collected from different places of Assam [Fb= Field bean; Guw= Guwahati, Assam and Dg= Dhaligaon, Assam].

- Fig. 4.4:** Naturally infected potato (*Solanum tuberosum*) leaf samples collected from different places of West Bengal (WB) and Assam [Po= Potato; Slg= Siliguri; Dg= Dhaligaon, Assam and An= Anthaibari, Assam].
- Fig. 4.5:** Naturally infected tomato (*Lycopersicon esculentum*) leaf samples collected from different places of West Bengal (WB) and Assam [To= Tomato; An= Anthaibari, Assam; Slg= Siliguri, WB and Guw= Guwahati, Assam].
- Fig. 4.6:** Naturally infected chili (*Capsicum annum*) leaf samples collected from different places of West Bengal (WB) and Assam [Ch= Chili; Snt= Sonitpur, Assam; Blg= Balurghat, WB and Sa= Sapatgram].
- Fig. 4.7:** Naturally infected ash gourd (*Benincasa hispida*) leaf samples collected from different places of West Bengal (WB) and Assam [Ag= Ash gourd; Guw= Guwahati, Assam; Mld= Malda, WB and Slg= Siliguri, WB].
- Fig. 4.8:** Naturally infected coriander (*Coriandrum sativum*) leaf samples collected from different places of West Bengal and Assam [Co= Coriander; Ng= Nayagaon, Assam; Gp= Goalpara, Assam and Cob= Cooch behar, WB].
- Fig. 4.9:** Naturally infected bottle gourd (*Lagenaria siceraria*) leaf samples collected from different places of West Bengal (WB), Assam and Tripura [L= Bottle gourd; Phn= Siliguri, WB; Agt= Agartala, Tripura; Cob= Cooch behar, WB; Ng= Nayagaon, Assam; Tg= Talguri, Assam; Jd= Jagdoi, Assam; An= Anthaibari, Assam; Slg= Siliguri, WB and Slc= Silchar, Assam].
- Fig. 4.10:** Naturally infected cucumber (*Cucumis sativus*) leaf samples and cucumber fruit with curling and mosaic symptom collected from different places of West Bengal (WB), Assam and Tripura [Cu= Cucumber; CuF= Cucumber fruit; Slc= Silchar, Assam; Slg= Siliguri, WB and Agt= Agartala, Tripura].

- Fig. 4.11:** Naturally infected garden rose (*Rosa* sp.) leaf samples collected from West Bengal [Ro= Garden rose and Slg= Siliguri, WB].
- Fig. 4.12:** Total RNA isolated from plant samples on 1% agarose gel under UV-transilluminator (a) From healthy plant samples and (b) From infected plant samples.
- Fig. 4.13:** Amplified PCR products from infected plant samples on 1% agarose gel under UV-transilluminator: (a), (b) and (c) Amplified with primer set 'CPuP/P9502'; (d) Amplified with primer set 'Potex5/Potex2RC' and (e) Amplified with primer set 'RRV For/RRV Rev'.
- Fig. 4.14:** (a) Healthy bottle gourd plant before mechanical sap inoculation; (b) Infected bottle gourd plant after 7 days of mechanical sap inoculation; (c) Healthy papaya plant before mechanical sap inoculation and (d) Infected papaya plant after 7 days of mechanical sap inoculation.
- Fig. 4.15:** Transmission electron microscopic photograph of (a) infected papaya leaf showing the presence of ~700 nm long filamentous rods of *Potyvirus*; (b) and (c) infected bottle gourd leaves showing presence of 500-700 nm long filamentous rods of *Potexvirus*; (d) and (e) infected rose leaves showing the presence of 120-150 nm double membrane bodies (DMBs) of *Rose rosette virus*. The bar given at the bottom right corner of each photograph represents 200 nm in cases of (a), (b) and (c); and 100 nm in cases of (d) and (e).
- Fig. 4.16:** (a) Vector map of pGEM-T vector (<https://www.promega.com/-/media/files/resources/protocols/technical-manuals/0/pgem-t-and-pgem-t-easy-vector-systems-protocol.pdf>, Website reference 4); (b and c) Blue-white screening of the cloned insert into pGEM-T vector on IXA (LB agar supplemented with IPTG, X-gal and ampicillin) plate.

- Fig. 4.17:** Nucleotide sequence identity matrix of CP gene of PRSV isolates of the present study and other PRSV isolates worldwide infecting different hosts. Identity percentages are indicated on the right side corner of the matrix.
- Fig. 4.18:** Phylogenetic tree generated by Neighbour-Joining method consisting of CP gene of PRSV isolates of the present study and other PRSV isolates worldwide infecting different hosts. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with their hosts have been indicated at the end of each branch.
- Fig. 4.19:** Nucleotide sequence identity matrix of CP gene of PVY isolates of the present study and other PVY isolates infecting different hosts. Identity percentages are indicated on the right side corner of the matrix.
- Fig. 4.20:** Phylogenetic tree generated by Neighbour-Joining method consisting of CP gene of PVY isolates of the present study and other PVY isolates infecting different hosts. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the hosts of the viruses have been indicated at the end of each branch.
- Fig. 4.21:** Nucleotide sequence identity matrix of CP gene of SMV isolates of the present study and other SMV isolates infecting different hosts (*G. max*, *V. unguiculata*, *P. edulis* and *L. siceraria*). Identity percentages are indicated on the right side corner of the matrix.
- Fig. 4.22:** Phylogenetic tree generated by Neighbour-Joining method consisting of CP gene of SMV isolates of the present study and other SMV isolates infecting different hosts (*G. max*, *V. unguiculata*, *P. edulis* and *L. siceraria*). Values at the nodes

indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the respective hosts have been indicated at the end of each branch.

Fig. 4.23: Nucleotide sequence identity matrix of RdRP gene of LaMMoV isolates of the present study and other *Potexvirus* isolates. Identity percentages are indicated on the right side corner of the matrix.

Fig. 4.24: Phylogenetic tree generated by Neighbour-Joining method consisting of RdRP gene of LaMMoV isolates of the present study and other *Potexvirus* isolates. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. GenBank accession numbers along with the respective viruses have been indicated at the end of each branch.

Fig. 4.25: Nucleotide sequence identity matrix of RdRP gene of RRV isolates of the present study and other *Emaravirus* isolates. Identity percentages are indicated on the right side corner of the matrix.

Fig. 4.26: Phylogenetic tree generated by Neighbour-Joining method consisting of RdRP gene of RRV isolates of the present study and other *Emaravirus* isolates. Values at the nodes indicate percentage of bootstrap support (out of 1000 bootstrap replicates) and are indicated if greater than 50. Acronym of the viruses along with their collection spots (acronym) have been indicated at the end of each branch. US/USA=United States of America; SLG=Siliguri, India; IND= India; SRV= Serbia; ITL=Italy; JPN=Japan; CAN= Canada.

Fig. 4.27: (a) Relationship between effective number of codons (ENC) and GC3 content of the coat protein genes of the four virus isolates. The continuous curve represents the expected curve between

GC3 content and ENC under random codon usage; (b) Relationship between the GC1/2 content and GC3 content of four virus isolates; (c) Contributions of 40 axes based on the correspondence analysis; (d) Correspondence analysis of the CP and RdRP genes. x and y axes correspond to Axis 1 and Axis 2 respectively. The coloured circles represent the values of different viruses as indicated at the top right corner of the graphs.

Fig. 4.28: Correlation between effective number of codons (ENC) of CP genes of PRSV with (a) adenine and thymine at 2nd codon position (AT2); (b) guanine and cytosine at 2nd codon position (GC2); (c) number of synonymous codons (L_sym); (d) frequency of aromatic amino acids (Aromo) and (e) Axis 4.

Fig. 4.29: Correlation between effective number of codons (ENC) of CP genes of PVY with (a) thymine at 3rd codon position(T3); (b) adenine at 3rd codon position(A3); (c) guanine at 3rd codon position(G3); (d) codon bias index (CBI); (e) frequency of optimal codon (Fop); (f) adenine and thymine at 1st codon position (AT1); (g) total percentage of guanine and cytosine (%GC); (h) adenine and thymine at 3rd codon position (AT3); (i) number of synonymous codons (L_sym) and (j) Axis 1.

Fig. 4.30: Correlation between effective number of codons (ENC) of RdRP genes of LaMMoV with (a) adenine and thymine at 2nd codon position (AT2); (b) guanine and cytosine at 2nd codon position (GC2); (c) total percentage of guanine and cytosine (%GC); (d) number of synonymous codons (L_sym); (e) frequency of aromatic amino acids (Aromo) and (f) Axis 1.

Fig. 4.31: Correlation between effective number of codons (ENC) of RdRP genes of RRV with (a) adenine and thymine at 2nd codon position (AT2); (b) guanine and cytosine at 2nd codon position(GC2) and (c) Axis 1.

- Fig. 4.32:** (a-e) *Myzus persicae* on the undersurface of infected leaves; (f) Mature *M. persicae* under simple microscope (10X) and (g-i) Different parts of mature *M. persicae* under compound microscope (100X).
- Fig. 4.33:** (a) Amplified PCR products from *M. persicae* on 1% agarose gel under UV-transilluminator with primer set 'CPuP/P9502'; (b) Healthy bottle gourd plantlet before vector inoculation and (c) Infected bottle gourd plant after 17 days of vector inoculation.
- Fig. 4.34:** Efficacy of BTH in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (BTH treated-uninoculated); (d) Healthy bottle gourd plant after 9 days post-treatment and inoculation (BTH treated-PRSV inoculated).
- Fig. 4.35:** Efficacy of AABA in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (AABA treated-uninoculated); (d) Infected bottle gourd plant after 9 days post-treatment and inoculation (AABA treated-PRSV inoculated).
- Fig. 4.36:** Efficacy of GABA in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (GABA treated-uninoculated); (d)

Healthy bottle gourd plant after 9 days post-treatment and inoculation (GABA treated-PRSV inoculated).

Fig. 4.37: Reduction of disease in bottle gourd plants (of Gadda-1 variety) caused by *Papaya ringspot virus* (PRSV) after application of chemical inducers: (a) BTH; (b) AABA and (c) GABA.

Fig. 4.38: Efficacy of *Azadirachta indica* ethanolic extract in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (*A. indica* treated-uninoculated); (d) Healthy bottle gourd plant after 9 days post-treatment and inoculation (*A. indica* treated-PRSV inoculated).

Fig. 4.39: Efficacy of *Bougainvillea spectabilis* ethanolic extract in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (*B. spectabilis* treated-uninoculated); (d) Healthy bottle gourd plant after 9 days post-treatment and inoculation (*B. spectabilis* treated-PRSV inoculated).

Fig. 4.40: Efficacy of *Camellia sinensis* ethanolic extract in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (*C. sinensis* treated-uninoculated); (d) Healthy bottle gourd plant after 9 days post-treatment and inoculation (*C. sinensis* treated-PRSV inoculated).

Fig. 4.41: Efficacy of *Clerodendrum infortunatum* ethanolic extract in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (*C. infortunatum* treated-uninoculated); (d) Infected bottle gourd plant after 9 days post-treatment and inoculation (*C. infortunatum* treated-PRSV inoculated).

Fig. 4.42: Efficacy of *Lantana camara* ethanolic extract in bottle gourd plants for reducing disease severity following challenge inoculation with PRSV. (a) Healthy bottle gourd plants (untreated-uninoculated control); (b) Infected bottle gourd plant after 9 days post PRSV inoculation (untreated-PRSV inoculated); (c) Healthy bottle gourd plant after 9 days post-treatment (*L. camara* treated-uninoculated); (d) Healthy bottle gourd plant after 9 days post-treatment and inoculation (*L. camara* treated-PRSV inoculated).

Fig. 4.43: Reduction of disease in bottle gourd plants (of Gadda-1 variety) caused by *Papaya ringspot virus* (PRSV) after application of botanicals: (a) *A. indica*; (b) *B. spectabilis*; (c) *C. sinensis*; (d) *C. infortunatum* and (e) *L. camara*.