

7. REFERENCES

- Agrell, L.P.S., & Lindquist, A.M. (1973). Physiological and Biochemical changes during insect development. In M. Rockstein (Eds.), *The Physiol. Insecta* (pp. 159 – 247).
- Akitamo, S., Egi, Y., Nakamura, Y., Suetsugu, Y., Oishi, K. & Sakamoto, K., (2017). Genome wide micro array screening for *Bombyx mori* genes related to transmitting the determination outcome of whether to produce diapause or non diapause eggs. *Insect. Sci*, 24 (2), 187 – 193.
- Amy, R.L. (1961). The embryology of *Habrobracon juglandis* (Ashmead). *J. Morph.*, 109, 199-217.
- Anderson, D.T. (1962). The embryology of *Dacus tryoni* (Frogg.) [Diptera]. *J. Embryol. Exp. Morph.*, 10, 248-292.
- Anderson, D.T. (1972). The development of hemimetabolous insects. In S.J. Counce, & C.H. Waddington, *Developmental Systems- Insects* (pp 95-163, 165-242).
- Anderson, D.T. (1973). *Embryology and Phylogeny in Annelids & Arthropods*. London: Pergamon Press.
- Anderson, D.T., & Wood, E.C. (1968). The morphological basis of embryogenic movements in the light brown apple moth, *Epiphyas postvittana* (Walk.) (Lepidoptera: Tortricidae). *Aust. J. Zool.*, 16, 763-793.
- Angelini, D.R., & Kaufman, T.C. (2005). Functional analysis in the milkweed bug *Oncopeltus fasciatus* (Hemiptera) support the role for Wnt signalling in body segmentation but not appendage development. *Dev. Biol.*, 283, 409 – 423.
- Audrewarthe, H.G., & Birchi, L.C. (1954). *The distribution of animals*. USA: The University of Chicago Press.
- Baba, H., Kuwabara, N., & Iwashita, Y. (1997). Scanning electron microscopic observation on the embryogenesis in the wild silk moth *Antheraea yamani*. *J. Seri. Sci. Japan*, 66 (2), 99-106.

- Bale, J.S., & Hayward, S.A. (2010). Insect over wintering in a changing climate. *J. Expt. Biol.*, 213, 980 – 994.
- Balinsky, B.I. (1981). *An Introduction to Embryology*. Philadelphia: Saunders College.
- Bardoli, S., & Hazarika, L.K. (1994). Body temperature and thermoregulation of *Antheraea assama* larvae. *Entomologia Experimentalis et Applicata*, 72, 207 – 217.
- Becker, A., Schlöder, P., Steele, J.E., & Wegener, G. (1996). The regulation of trehalase metabolism in insects. *Experientia*, 52, 433-439.
- Beenackers, A.M. Th., Van-der -Horst, D.J., & Marrewijk, W.J.A. (1981). Role of lipids in energy metabolism. *Energy Meta. Insects*, 53 – 100.
- Berdu Jr, J.G., & Pereira, M.F. (2013). Silk Production in Latin American Regions – Constrains & the way forward. *Sericologia*, 53(1), 15-23.
- Bergmeyer, H.U., Gawehn, K., & Grassl, M. (1974). In H.U. Bergmeyer (Eds.), *Methods of Enzymatic Analysis* (Volume I, pp. 521-522). New York, NY : Academic Press Inc.
- Biswas, I., and Ray, N. (2007). Studies on short term cold preservation for supply of seed during main commercial crop rearing of muga silkworm. *J.Exp., India.*, 10(2), 271 -276.
- Bloter, C.J., & Chefurka, W., (1990). The effect of the phosphate treatment on superoxide dismutase, catalase and peroxidase in the granary weevil, *Sitophilus granaries*. *Pestic. Biochem. Physiol.*, 36, 52-60.
- Bora, A., Borah, B., & Sengupta, A.K. (1990) Studies on presentation of muga cocoons at high altitude and low temperature. *Ann Report (1988-90), R.M.R.S. Boko*, 30 – 32.
- Bora, D.S. (2006). Sensitivity of *Antheraea assama* Westwood to interaction of photo period and thermoperiod. In Muga silkworm : Bio chemistry, Bio technology, and Molecular Biology (pp. 67–74).
- Bosewell, R.E., and Mahowald, A.P. (1985). Cytoplasmic determinants in embryogenesis. In G.A Kerkut & L.I. Gilbert (Eds.), *Comprehensive Insect Physiology, Biochemistry and Pharmacology* (Vol. I, pp. 387-405). Oxford: Pergamon Press.

- Boyer, R.F. (2006). *Modern experimental Biochemistry* (pp.389-405). New Delhi: Pearson.
- Brachet, J. (1957). *Biochemical Cytology*. New York: Academic Press.
- Brachet, J. (1962). Nucleic acid in development. *J. Cell. Comp. Physiol.*, 60, Supple.I, 1-18.
- Casperson, T. (1950). *Cell growth and cell function*. New York: W.W. Norton Company Inc.
- Chandrasekhar, P.M., & Bali, G. (1987). Glycogen level and glycogen phosphorylase activity in eggs of silk worm *Bombyx mori* L. *Proc. Indian Acad. Sci. (Anim. Sci.)*, 96(1), 49-54.
- Chaudhury, A. (1998). Biochemical studies of certain enzymes of carbohydrate metabolism at some stages of development of silk worm *Antheraea assama*, Westwood and *Philosamia ricini*, Boisduval. (Ph.D. Thesis), Guwahati University.
- Chen, P.S., & Brigel, H. (1965). Studies on the protein metabolism of *Culex pipiens*, L.V. changes in free amino acids and peptides during embryonic development. *Comp. Biochem. Physiol.*, 14, 463 – 473.
- Chen, Y.R., Jiang, T., Zhu, J., Xie, Y.C., Tang, J.C., Chen, Y. H., Tang, S.M., Hao, B.F., Wang, S.R. Huang, J. S., Shen, X.J. (2017). Transcript sequencing reveals potential mechanism of diapause preparation in bivoltine silk worm, *Bombyx*. (Lepidoptera: Bombycidae) *Comp. Biochem. Physiol. Part D. Genomics proteomics*, 24, 68 – 78.
- Chino, H. (1957a). Carbohydrate metabolism in the diapause egg of the silkworm *Bombyx mori*. I Diapause and change in glycogen content. *Embryologia*, 3(4), 295-316.
- Chino, H. (1958). Carbohydrate metabolism in the diapause egg of the silkworm *Bombyx mori*. II Conversion of glycogen into sorbitol and glycerol during diapauses. *J. Insect Physiol*, 2, 1-12.

- Chippendale, G.M. (1970). Metamorphic changes in the fat body proteins of the south western corn borer, *Diateraea grandiosella*. *J. Insect Physiol.*, 16, 1057-1068.
- Chippendale, G.M. (1978). The function of carbohydrate in insect life process. In M. Rockstein (Eds.), *Biochemistry of Insects* (pp1-55). Academic Press.
- Choudhury, B., & Sahu, A.K. (2000, June 13-14). Muga Seed Production by private grainers. Paper presented at the workshop on *Development of Muga & Eri Silk industries through improved technologies*, N. Lakhimpur, Assam.
- Choudhury, B., Bhattacharya, A., Singh, B. K., Siddiqui, A.A., Das, P., & Jayprakash, P. (2012). Disease incidence dynamics of muga silk worm (*Antheraea assamensis*, Helfer) and their correlation with meteorological factors. *J. Appl. Biosci.*, 38(1), 101-104.
- Choudhury, P., Baruah, S., & Devi, D. (2013). Changes in carbohydrate and protein in the developing healthy and pebrine infected embryos of *Antheraea assamensis*, Helfer. *IJBPAS*, 2(8), 1627-1632.
- Choudhury, S.N. (1981). *Muga Silk Industry*. Assam: Directorate of Sericulture.
- Choudhury, S.N. (2006, Feb. 17-18). Problems and prospects of muga silkworm seed production. Paper presented in the workshop on *Problem and prospect of muga silkworm seed production* (pp. 1-10). Muga silkworm seed organization, Guwahati.
- Choudhury, B., Bhattacharya, A., Singh, B.K., & Provakar, C.J. (2010) Effect of global warming in the rearing performance of muga silkworm *Antheraea assamensis*, Helfer. *J. Assam Sc. Soc.*, 51(1), 134-140.
- Christensen, P. J. H. (1943), Embryologische und zytologische studien über die erste und frühe Eientwicklung bei *Orgyia antique* Linne (Fam. Lymantriidae, Lepidoptera). *Vidensk. Meddr dansk naturh. Foren.*, 106, 1-223.
- Colombo, G., Benassi, C.A., Allegri, G. & Longo, E. (1962). Free amino acids in eggs of *Schistocerca gregaria* F. during development. *Comp. Biochem. Physiol.*, 5, 88 – 93.

- Copf, T., Schroder, R., & Averof, M. (2004). Ancestral role of caudal genes in axes elongation and segmentation. *Proc. Natl. Acad. Sci. USA.*, 101, 17711 – 17715.
- Counce, S.J. (1961). The analysis of insect embryogenesis. *A. Rev. Ent.*, 6, 295-312.
- Counce, S.J. (1973). The causal analysis of insect embryogenesis. In S.J. Counce, & C.H. Waddington, *Developmental Systems- Insects-2* (pp 1-156).
- Damen, W.G. (2007). Evolutionary conservation and divergence of the segmentation process in arthropods. *Dev. Dyn.*, 236, 1379 – 1391.
- Das, P. (2000, June 13-14). Base Paper of Muga Silk Industry in Assam. Paper presented at the workshop on *Development of Muga & Eri Silk industries through improved technologies*, N. Lakhimpur, Assam.
- Dash, A.K., & Nayak, B.K. (1988). Effect of refrigeration on hatching of egg of the tasar silk moth *Antheraea mylitta*, Drury,(Saturindae) *J. Res. Lepid.*, 27 (3-4), 259 -271.
- Datta, R.K., Sengupta, A.K., & Biswas,S.N.(1972). Studies on refrigeration of multi voltine silkworm eggs at low temperature. *Indian J. Seric.*, 11, 20 – 27.
- Davis, G.K., Patel, N.H. (2002). Short, long and beyond molecular and embryological approaches to insect segmentation. *Annu. Rev. Entomol.* 47, 669–699.
- Dearden, P., & Akam, M. (2001). A role for wings in segment morphogenesis but not segment formation in the grasshopper *Schistocerca gregaria*. *Dev. Genes Evol.*, 210, 329-336.
- Denlinger, D.L.,(2002). Regulation of diapause. *Annu. Rev. Entomol.*, 31, 239 –264.
- Denlinger, D.L. (1986). Dormancy in tropical insect. *Anu. Rev. Entomol.*, 47, 93 – 122.
- Denlinger, D.L., & Lee, Jr. R.E. (2010).*Low temperature Biology of Insects*. NewYork: Cambridge University Press.
- Dolin, M.I. (1957). *Journal of Biological Chemistry*, 225, 557-573.
- Dukes, H.H. (1955). The physiology of domestic animals. New York: Camstock Publishing Associates.

- Dutta, R.K., Sengupta, K., & Biswas, S.N.(1972). Studies on the refrigeration of multiroutine silkworm eggs at low temperatures. *Indian Journal of Science*, 11, 20-27.
- Dutta, R.K.,(1988). Embryological studies of non-diapause silkworm *Bombyx mori* L. *Ind. J. Seric.*, 27, 1-6.
- Engelmann, F. (1970). *The Physiology of Insect Reproduction*. Oxford & New York: Pergamon Press.
- Felton, G.W., & Sumners, C.B. (1995). Antioxidant systems in insects. *Arch. Insect Biochem. Physiol.*, 29, 187-197.
- Fujiwara, Y., Tanaka, Y., Iwata, K., Rubio, R.O., Yaginuma, T., Yamashita, O., & Shiomi, K. (2006). ERK/MAPK regulates ecdysteroids and sorbitol metabolism for embryonic diapause termination in the silk worm *Bombyx mori*. *J. Insect Physiol.*, 52(6), 569-575.
- Fullilove, S.L., & Jacobson, A.G. (1971). Nuclear elongation and cytokinesis in *Drosophila montona*. *Dev. Biol.*, 26, 560-577.
- Furusawa, T., Indrasith, L.S., & Shikata, M. (1985). Changes in DNA, RNA content and RNA/DNA ratios during the embryonic diapause of *Bombyx mori*. *J. Seric. Sci. Jpn.*, 54(1),53-62
- Furuswa, T. & Yang, W.J. (1987). Fluctuation of free sugars with embryonic development of non diapause eggs of silk worm *Bombyx mori*. *J. Seric. Sci. Jpn.*, 56, 143-149.
- Gerlach, U.,& Hiby, W. (1974). In H.U. Bergmeyer (Eds.), *Methods of Enzymatic Analysis* (Volume II, pp. 569-573). New York, NY : Academic Press Inc.
- Ghosh, S., & Ray, N. (2005). Effect of low temperature preservation delayed hatching of eggs of of eggs of muga silk worm, *Antheraea assama*, Westwood. In U. Chakraborty & B. Chakraborty (Eds.), *Stress Biology* (pp117-121). New Delhi: Naraosa.

- Ghosh, S., & Ray, N. (2006). Embryological studies of silk worm *Antheraea assama* Ww., Paper presented at the Workshop on problems & prospects of Muga silk worm seed production, MSO, CSB, Guwahati, 17-18th Feb. (pp. 29).
- Goswami, D., Singh, N.I., Ahamed, M., & Giridhar, K. (2013). Embryonic development in Muga silk worm, *Antheraea assamensis* Helfer (Lepidoptera: Saturniidae). *Mu. Entomology & Zoology*, 8 (2): 852-857.
- Govindan, R., Devaiah, C., Rangaswami, H.R., & Thippeswami, C. (1980). Effects of refrigeration of the eggs of Eri Silkworm, *Samia Cynthia Ricini*, Biosduval on hatching. *Indian J Seric.*, 19, 13 -15.
- Gross, R.J. (1952). The early embryology of the book louse, *Liposcelis divergens* Badonnel (Poscoptera: Liposcelidae). *J. Morph.*, 91, 135-167.
- Gross, R.J. (1953). The advanced embryology of the book louse, *Liposcelis divergens* Badonnel (Poscoptera: Liposcelidae). *J. Morph.*, 92, 157-191.
- Gutzeit, H.O., Zissler, D., Grau, V., Liphardt, M., & Heinrich U.R. (1994). Glycogen stores in mature ovarian follicles and young embryos of *Drosophila*. Ultra structural changes in some biochemical correlates. *Eur. J. Cell. Biol.*, 63, 52- 60.
- Görg, R.J. (1952). Untersuchungen am Keim von *Hierodula* (Rhombodera) *crassa* Giglio- Tos, ein Beitrag zur Embryologie der Mauotiden (Mantodea). *Dt. Ent. Z.*, 6, 389-450.
- Hagan, H.R.(1951). *Embryology of the viviparous insects*. New York: Ronald Press Co.
- Hakim, R.S., Boldwin, K., & Snaghe, G. (2010). Regulation of Midgut Growth, Development and Metamorphosis. *Annu. Rev. Entomol.*, 55, 593-608.
- Han, Q., Li, G., & Li, J. (2000a). Chorion peroxidase mediated NADH/O₂ oxidoreduction cooperated by chorion malate dehydrogenase- catalyzed NADH production: a feasible pathway leading to H₂O₂ formation during chorion hardening in *Aedes aegypti* mosquitoes. *Biochim. Biophys. Acta*, 1523 (2-3), 246-253.

- Han, Q., Li, G., & Li, J. (2000b). Purification and characterization of chorion peroxidase from *Aedes aegypti* egg. *Arch. Biochem. Biophys.*, 378 (1), 107-115.
- Hansen, H. (1946). The theoretical aspect of insect metamorphosis. *Biol. Rev.*, I, 21.
- Hasegawa, K., & Yamashita, O. (1965). Studies on the mode of action of the diapause hormone in the silkworm *Bombyx mori*, VI- the target organ of diapause hormone. *J. Exp. Biol.*, 43, 271-277.
- Hayakawa, Y., & Chino, H. (1981). Temperature dependent inter conversion between glycogen and trehalose in diapausing pupae of *Philosamia Cynthia ricini* and *pryeri*. *Insect Biochem.*, 11 (1), 43-47.
- Hayakawa, Y., & Chino, H. (1982). Phosphofructokinase as a possible key enzyme regulating glycerol and trehalose accumulation in diapausing insects. *Insect Biochem.*, 12 (6), 639-642.
- Hiratsuka, K. (1975). *Text Book of Tropical Sericulture*. Tokyo: Japanese Overseas Cooperation Volunteers.
- Hochachka, P. W., Buck, L. T., & Land, S.C. (1996). Unifying theory of hypoxia tolerance: molecular / metabolic defense and rescue mechanisms for surviving oxygen lack. *Proc. Natl. Acad. Sci. USA*, 93, 9493-9498.
- Horie, Y., Kanda, T., & Moehida, Y. (2000). Sorbitol as an arrester of the embryonic development in diapausing eggs of the silkworm *Bombyx mori*. *J. Insect Physiol.*, 46, 1009-1016.
- Howe, R. H. (1967). The temperature effects on embryonic development in insects. *Annu. Rev. Ent.*, 12, 15- 42.
- Huetter, A.F. (1923). The Origin of the germ cells in *Drosophila melanogaster*. *J. Morph.*, 37, 385-423.
- Hutcher, P., & Battery, N. (2011). Sericulture: Silkworms & mulberries. In *Biological Diversity Exploiters & Exploited* (Chapter 5, pp73 – 96). London: Wiley & Blackwell.

- Ibohal Singh, N., Ibotombi Singh, L, Somen Singh, L., & Suryanarayana, N., (2005). An overview of the salient features of *Antheraea frithii*, Moore- a promising wild silk moth of India. In *Proceedings of 20th International Congress of wild silk moth* (Vol II, pp. 74-77), Bangalore: Central Silk Board.
- Ikada,E. (1910). Embryonic development and morphological changes of *Bombyx mori*. *Sangyoh Shinpoh*, 18, 26-42.
- Ikada,E.(1912).Morphological change of silk worm eggs untill 2 days from oviposition. *Sangyoh Shinpoh*, 20,9-11; 43-47.
- Ikekawa, N., Saito, S.M., Kobayashi, M., & Tsuda, K. (1966). Studies on the sterol of *Bombyx mori* L. IV. Conversion of Sterol in the silk worm. *Chem. Pharm. Bull.*, 14,834-836.
- Imanishi, S., Sinbo, H.,& Kiguchi, K.(1996). Cryopreservation of silk worm eggs. *J. Seric. Sci. Jpn.*, 65(4), 235 – 240.
- Indira, T. (1963). Biochemical and cyto chemical studies during development on ovarian growth in *Spheroderma molestum* (Dub) Ph. D. Thesis, Annamalai University, India.
- Jaworski, T., & Hilszezanski (2003). The effect of temperature and humidity change of insect development and their impact on forest ecosystem in the context of expected climate change. *Forest Research Papers*. ,74(4), 345 – 355.
- Jayaprakash, P., Bhattacharya, A., Choudhury, B., Singh, B.K., & Hajarika, H.K.(2012). Muga silk production- its tribulations, prospects and inroads made in the face of unprecedented impugnations. In M. Chutia, S. A. Ahmed, K. Neog, R. Kumar & K. Das (Eds.), *Compendium of Seminar Papers*. Paper presented at the National Seminar on Recent Trends in Research & Development in Muga Culture-Ideas to action. MSO, Guwahati, 3-4May 2012 (pp. 141-150).Guwahati: Central Silk Board.
- Johanusen, D., & Butt, A.H. (1941). *Embryology of insectsand Myriopods*. New York: Mc Graw- Hill.

- Jolly M.S. (1979). *FAO Agricultural Services bulletin (291)*. Rome: Food and Agricultural Organisation of the United Nations.
- Jolly, M.S. (1983). Organization of bivoltine grainage for tropics. In *Sericulture, Project-3*. Mysore :Central Sericultural Research and Training Institute.
- Jolly,M.S., Sen, S.K., & Ahsan, M, (1974). *Tasar Culture*. Bombay: Ambika Publishers.
- Jura, C. (1972). Development of a pterygote insects. In S.J. Counce & C. H. Waddington (Eds.), *Development Systems –Insects (Vol.-I)*. London &New York: Academic Press.
- Jura, C. (1972). Development of Apterygote insects. In S.J. Counce, & C.H. Waddington, *Developmental Systems- Insects-1* (pp 49-94).
- Karlson, P. & Hofmeister (1963). Zur bigenese des ecdysons. I. Unwandlung von cholesterin in ecdyson hoppe. *Seyler's Z. Physiol. Chem.*, 331, 298.
- Karlson, P. (1963). Chemie and biochemie der insekt euhormone. *Angew. Chem.*,75, 257-265.
- Karnavar,, G.K., & Nair, (1969) Changes in body weight, fat, glycogen and protein during diapauses of *Trogoderma granarium*. *J. Insect Physiol.*, 15, 95-102.
- Katsuki, M., Murakami, A., & Watanabe, I. (1980). Fate mapping of some tussues in the genetic mosaics of the silk worm *Bombyx mori*. *Zool. Mas.* 89, 269-276.
- Kawaguchi,Y., Kawakami, K., Kusakabe, T., & Koga, K. (2004). Histological Observation of eggs of the embryonic lethal mutation set in *Bombyx mori*. *J. Insect Biotech.and Sericology*, 73,101 –106.
- Keino, H., & Takesue, S. (1982). Scanning electron microscopic study on the early Development of silkworm Eggs (*Bombyx mori* L.). *Develop. Growth & Differ.* , 24 (3), 287-294.
- Khanikar, D.P., & Choudhury, S.N.(1997, Nov 7-8). Effects of preservation of muga seed cocoons at low temperature. Paper presented at the Seminar on *Environmental potential and conservation of nature and natural resources* (pp.22) Sibsagar College, Sibsagar, Assam.

- Khanikar, D.P. ,& Dutta, S.K. (1998b). Effect of induced pupal diapause on rearing performances and cocoon parameters of muga silkworms, *Antheraea assama* Westwood. *J. Agric. Sci. Soc., North East India*, 11 (20), 140 – 143.
- Khanikar, D.P., & Dutta, S.K. (1998a). Low temperature preservation of seed cocoons for delaying moth emergency in muga silkworm, *Antheraea assama* ,Westwood (Lepidoptera: Saturniidae). *Int. J. Wildsilkmoth and silk*, 5, 216 -220.
- Khanikar ,D.P. and Dutta, S.K. (1998, Nov 11 – 14). Low temperature to seed cocoons for delaying moth emergence in muga silk worms (*Antheraea assama* Westwood) Lepidoptera : Saturnidae. Paper presented at the 3rd *International Conference on Wild Silk moths*. Bhubaneswar, Orissa.
- Kihara,F., Itoh, K., Iwasaka, M., Niimi, T., Yamashita, O., & Yaginuma, T. (2009). Glycerol kinase activity and glycerol kinase-3 gene are up regulated by acclimation to 5°C in diapauses eggs of *Bombyx mori*. *Insect. Biochem. Mol. Biol*, 39 (11),763-769.
- King, R.C. (1970). *Ovarian Development inDrosophila melanogaster*. New York: Academic Press.
- Kodri, K. D., Bednaraova, A., Zemavova, M. & Krishnan, N. (2015). Hormonal regulation of response to oxidative stress in insects- an update. *Int. J. Mol. Sci.*, 16, 25788-25816.
- Krause, G. (1957). Neue Beitrage zur Entwicklungsphysiologie des Insekten Keimes. *Verh. dt. zool. Ges.(Graz)*, 396-424.
- Krishnappa, B. L., Reddy, D.N.R., and Narayanaswamy, K.C. (2001) Changes in carbohydrate, lipid and moisture contents during the embryonic development of *Samia cynthia ricini*, Biosdual. *Indian J. Seric.*, 2,180-181.
- Kühn, A. (1971). *Lectures on Developmental physiology*. Berlin & New York: Springer.
- Kumaresan, P., & Quadri, S.M.H.(2012). Growth & instability of mulberry silk production in India. *Indian J. Seric.*, 51(1),64-71.

- Kumereshan, P., Thangavelu, K., & Sinha, R.K.(2004). Studies on long term preservation of eggs of Indian tropical multi-voltine silk worm (*Bombyx mori*, L.) genetic resources. *Int. J. Indust. Entomol.* 9(1). 79-87.
- Kurata, S., Yaginuma, T., Kobayashi, M., Koga, K., & Sakaguchi, B. (1980). DNA content and cell number during embryogenesis of *Bombyx mori*. *J. Sericult. Sci. Jpn.*, 49 (2), 107-110.
- Kuwana, J., & Takami, T. (1957). *Insecta. An Embryology of Invertebrates*. In M.Kume & K. Dan, (pp287-343). Tokyo: Baifukan .
- Lalouette, L., Williams, C.M., Hervant, F., Sinclair, B.J., & Renault, D. (2011). Metabolic and oxidative stresses in insects exposed to low temperature thermal fluctuation. *Comp. Biochem. Physiol.*, 158, 229-234.
- Larinink, O. (1997). Apomorphic & plesiomorphic characteristics in Archeognatha, Monura and Zygentoma. *Pedobiologia*, 41, 3-8.
- Lee, Jr. R.E. (2010). A primer on insect cold tolerance. In D. L. Denlinger & Jr. R.E. Lee(Eds.), *Low temperature Biology of Insects* (Chapter 1). NewYork: Cambridge University Press.
- Legay, J.M. (1974). Sur la croissance sequentielle de l'ovocyte dans le tube ovarique an *Bombyx mori* pendant le stade nymphal. *Annls. Embryol. Morph.* 7, 277-282.
- Leopold, R.A. (2000). Short term cold storage of housefly (Diptera: Muscidae) Embryos : survival and quality of subsequent stages. *Annals of Entomological Society of America* Vol 93 (4), 884 – 889.
- Levinson, Z.H. (1960). The evolution of sterol requirements.the function of dietary syterol in phytophagus insects. Paper presented at *the 11th Int.Congr. Ent.*, 3, (pp.145-156).
- Lim, S.H., Kim, Y.T., Lee, S.P., Rhee. I.J., Lin, J.S., & Lin, B.H. (1990). Sericulture training Manual. *FAO Agricultural Services Bulletin* 80. Rome: Food & agricultural training organization of the United Nations.
- Liu, P. Z., & Kaufman, L. C. (2005). Short and long germ segmentation: unanswered questions in the evolution of a developmental mode. *Evol. Dev.*, 7(6). 629 – 646.

- Liu, W. (2012). Functional analyses in the silk worm, *Bombyx mori*, support a role for Notch signalling in appendage development but not the groucho-dependent pair rule process. *J. Exp. Zool. (Mol. Dev. Evol.)* 9999B, 1 – 12.
- Lonvet, J.P. (1964). La Segregation du mesoderme chez l'embryon du Phasme *Carausius morosus* Br. *Bull. Soc. Zool. Fr.*, 89, 688-701.
- Lowry, O.H., & Passonneau, T.V. (1972). *A flexible system of enzymatic Analysis* (pp. 71-77, 179-182). New York: Academic Press.
- Lynnech, J. A., Brent, A. E., Leaf, D.S., & Plutz, M.A . Desplan.C.(2006). Localized maternal orthodenticle patterns anterior and posterior in the long germ wasp *Nasonia*. *Nature* ,439, 728 – 732.
- Mahowald, A.P. (1963). Electron microscopy of the formation of the cellular blastoderm in *Drosophila melanogaster*. *Exp. Cell Res.*, 32, 457-468.
- Makka, T., Seino, A., Tomita, S., Fujiwara, H., Sonobe, H. (2002) A possible role of 20-hydroxyecdysone in embryonic development of the silk worm *Bombyx mori*. *Arch. Insect Biochem. Physiol.* 51, 111-120.
- Makka, T., & Sonobe, H. (2000). Ecdysone metabolism in diapauses and non diapause eggs of silk worm *Bombyx mori*. *Zool. Sci.*, 17, 89-95.
- Martini, A., Baldassari, N., & Baronis, p. (2006). Embryonic development in *Neodiprion sertifer*. *Bulletin of Insectology* ,59 (1), 45-52.
- Mc Gregor, D.A., & Loughton, B.G. (1979) Yolk protein degradation during embryogenesis of the African migratory Locust. *Can. J. Zool.*, 7, 907 – 917.
- Mc Gregor, A. and Loughton, G. (1974). Yolk- protein degradation during embryogenesis of the African migratory locust, *Can. J. Zool.*, 52, 907 – 917.
- Miya, K. (1950). *Trans. Sapporo Nat. Hist. Soc.*, 19, 36-39.
- Miya, K. (1953). The presumptive genital region at the blastoderm stage of the silk worm egg. *J. Fac. Arg. Iwate. Univ.*, I, 223-227.

- Miya, K. (1958). Studies on the embryonic development of gonad in the silk worm *Bombyx mori* L. Part I: Differentiation of germ cells. *J. Fac. Arg. Iwate. Univ.*, 3, 436-467.
- Miya, K. (1960). Electron microscopic studies on the embryonic development in the silkworm *Bombyx mori*, II, Fine structure of the serosa. *J. Seric. Sci. Jpn.*, 29,273.
- Miya, K. (2003). *The early embryonic development of Bombyx mori: An ultra structure point of view*. Tokyo: GENDAITOSHO.
- Miyawaki, K., Mito, T. Sarashina, I. Zhang, H., Kinmyo, Y., Ohuchi, H.,& Noji, S. (2004). Involvement of Wingles / Armadilo signalling in the posterior sequential segmentation in the cricket. *Gryllus bimaculatus* (orthoptera) as revealed by RNAi analysis. *Mech. Dev.*,121, 119 – 130.
- Mizuno, T., Watanbe, K., and Ohushi, E.(1981). Developmental changes of Ecdysteroids in the silk worm *Bombyx mori*. *Develop. Growth and Differ.*, 23(5),543-552.
- Moorthy, S.M., Krishnan, N., Bhattacharya, T., & Choudhuri, A. (2007). Changes in the titre of protein and cholesterol content in non diapause, artificially diapause terminated and diapauses eggs of silk worm, *Bombyx mori*. *Int. J. Indust. Entomol.*, 15(2), 165-169.
- Myer, B.L.B. and Burggrem, W. W. (2010) Metabolic Allometry during development and metamorphosis of the silk worm *Bombyx mori*. *Analysis Patterns and Mechanisms. Physiological and Biochemical Zoology*, 83(2), 215 – 231.
- Nace, G.W. (1970). Specific macromolecular changes during development. In O.A. Schjeide & J. Devellis (Eds.), *Cell Differentiation* (pp. 322-350). New York : Norstrand Reinhold Company.
- Nagina, N., & Nagesh Chandra, B.K. (1988). On the effect of low temperature on egg hatch and subsequent development of *Samia Cynthia Ricini* Biosduval (Lep:Sat). *Sericologia*, 28 (3), 337 – 341.

- Nagy, L., Riddiford, L., & Kiguchi, K. (1994), Morphogenesis in the early embryo of the Lepidopteran *Bombyx mori*. *Dev. Biol.*,165(1) , 137-151.
- Nagy, L.M. & Carrol, S. (1994). Conservation of wingless patterning functions in the short germ embryo of *Tribolium castaneum*. *Nature*, 367, 460–463.
- Nakada, T. (1932). The resistance of Silkworm embryo to abnormal high temperature and dryness on the external morphology of the embryo. *Bull. Fukuoka Sericult. Expt. Stn.*, 1, 1 -26
- Nakao, H. (2010). Characterization of Bombyx embryo segmentation process : expression profile of engrailed , even-skipped, Caudal and wnt 1/ Wingless homologue. *J. Exp. Zool. (Mol. Dev. Evol.)*, 314B, 224 – 231.
- Nakao,H. (2012). Anterior and posterior Centres jointly regulate *Bombyx* embryo body segmentation. *Dev. Biol.* [http:// dx.doi.org/10.1016/j.ydbio.2012.08.029](http://dx.doi.org/10.1016/j.ydbio.2012.08.029).
- Narayanaswamy, T.K.,& Govindan, R. (1987). Effect of Refrigeration of eggs of pure Mysore silkworm, *Bombyx mori*, at blue stage. *Entomon*, 12, 105-107.
- Nation, J.L. (2008). *Insect Physiology and Biochemistry*. Boca Raton, USA: CRC Press.
- Nayak, B.K. & Dash, A.K.(1989). Effect of refrigeration egg incubation period of the Tasar silk *Antheraea mylitta* Drury (Saturnidae) *J. Lepidopteran Society.*, 43(2), 152 -153.
- Newman, S. A. (2011). Animal egg as Evolutionary Innovation : A solution to the “Embryonic Hourglass” Puzzle. *J. Exp. Zool. (Mol. Dev. Evol.)* 316, 467 – 483.
- Niimi, T., & Yaginuma, T. (1992). Biosynthesis of NAD- Sorbitol dehydrogenase induced by acclimation at 5°C in diapauses eggs of the silk worm *Bombyx mori*. *Comp. Biochem. Physiol.*, 102,169-173.
- Niimi,T., Yamashita,O., & Yaginuma, T. (1993). Developmental profile of the gene expression of the Bombyx homolog of mammalian sorbitol dehydrogenase during embryogenesis of non diapauses eggs. *Comp. Biochem. Physiol. B*, 106 (2), 437-442.

- Ohnishi, E., Mizuno, T., Chatani, F., & Sakurai, S. (1977). 2- Deoxy-d-ecdysone from ovaries and eggs of the silk worm, *Bombyx mori*. *Dev. Growth Differ.*, 19, 67-70.
- Ohtsuki, Y., Mori, S., Kanda, T., & Kitazawa, T. (1976). Morphological observation on embryonic moult in the silk worm, *Bombyx mori*. *J. Sericult. Sci. Japan* ,45(3), 225-231.
- Ohtsuki, Y. (1979). Silk worm eggs. In The Japanese Society of Sericultural Science (Eds.), *A general text book of Sericulture* (pp.156-173). Tokyo: Nihon Sanshi Sinbun Sha.
- Okazaki, T. & Yamashita, O. (1981). Changes in glucose and fructose contents during embryonic development of silk worm, *Bombyx mori*. *J. Seric. Sci. Jpn.*50(3),190-196.
- Ohtsuki, R., Sato, S., Nagai, O., Horigone, I., Yoshitake, S., Goto, S., Sato, S., Totani, C., Nagashima, E., & Yoshitake, S. (1997). Silk worm egg production. New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Overgaard, J., & Mac Millan, H.A. (2017). The integrative physiology of insect chill tolerance. *Annu. Rev. Physiol.*, 79,187-208.
- Pandey, R.K., Luoklam. R., Das, P.K., & Noamani, M.K.R.(1992). Effect of cold storage on the hatchability of oak tasar silkworm. *Indian Silk*, 31(2), 155 -157.
- Panfilio, K.A. (2008). Extraembryonic development in insects and the acrobatics of blastokinesis. *Dev. Biol.* 313, 471 – 491.
- Pant, R. & Nautiyal, G.C. (1974). Change in protein, glycogen, free sugar content and active phosphorylase activity during embryogenesis of *Philosamia ricini*. *Proc. Indian Acad. Sci. (Sec. B)*, 80, 121-126.
- Pant, R., & Sharma, K.K. (1976). Variation in glycogen, total carbohydrate, free sugars, proteins, total free amino acid and some enzymes (active phosphorylase , acid and alkaline phosphatases) during embryonic development of *Antheraea mylitta*. *Curr. Sci.*, 45, 125-128.

- Park, K.E., & Yoshitake, N. (1969). Effect of incubation temperature on nucleic acid and protein synthesis during embryonic development of the silk worm *Bombyx mori* L.(Lepidoptera: Bombycidae). *Appl. Ent. Zool.*, 4(4), 171-176.
- Peel, A.D., Chapman, A.D., & Akam, M. (2005). Arthropod segmentation : beyond the *Drosophila* paradigm. *Nature Rev. Genet.*, 6, 905-916.
- Pflugfelder, O. (1958). *Entwicklungsphysiologie der Insekten*. Leipzig: Akademische Verlagsgesellschaft.
- Plummer, D.T. (1998). *An Introduction to Practical Biochemistry* (pp. 216-219).New Delhi: Tata Mc Graw-Hill.
- Polten, K.A, Davidowitz, G., & Woods, H.A. (2011) Gross stage consequences of egg temperature in the insect *Manduca sexta*. *Functional Ecology*, 25, 548 – 556.
- Ponnuvel, et. al. 2010. Ponnuvel, K.M., Murthy, G.N., Awasthi, A.K., Rao, G., & Vijayprakash, N.B. (2010). Differential gene expression during early embryonic development in diapause and non diapause eggs of multivoltine silk worm *Bombyx mori*. *Indian J. Exp. Biol.*, 48(11), 1143 – 1151.
- Sarkar, B.N., Sarmah, M.C., Dutta, P., & Dutta, K.(2012). Embryo Isolation and egg preservation technology of Eri silkworm *Samia ricini* (Donovan) (Lepidoptera: Saturniidae). *Munis Entomology & Zoology*, 7 (2), 792-797.
- Rabecea, L.F., & Palfey, B.A. (2010).Flavin independent enzymes. In *Reference module in Chemistry, Molecular Science and Chemical engineering: Comprehensive natural products II*, Vol. 7, Cofactors, (pp.37-43).
- Raikhel, A.S., & Dhadialla, T.S.(1992). Accumulation of yolk proteins in insect oocytes. *Annu. Rev. Entomol.*, 37, 217 – 251.
- Ranjan, R.K., & Hazarika, U. (2012). Constrains in muga culture: strategies & research programmes undertaken at CMERTI Lahdoigarh. In M. Chutia, S. A. Ahmed, K. Neog, R. Kumar, & K. Das (Eds.), *Compendium of Seminar Papers*. Paper presented at the National Seminar on Recent Trends in Research & Development in Muga Culture-Ideas to action. MSO, Guwahati, 3-4May 2012 (pp. 3-11).Guwahati: Central Silk Board.

- Ranna, B., Luikham,R., Chalapatty, M.V., Debaraj,Y., Singh, N .I., Sinha, R.K., & Thangavelu, K. (2002). Effect of refrigeration on the hatchability of different ages of Oak tasar silkworm eggs. *Sericologica*, 42(4), 581 -584.
- Read, E. M., & Day, M.R. (1966). Embryonic movements during development light Brown Apple Mouth. *Aust. J. Zool.*, 14, 253-263.
- Richards, O.W., & Davies, R.G. (1977). *Imms general text book of entomology*. London: Chapman & Hall.
- Rockstein ,M. (1974). The insect and external environment. In *The physiology insects* (Vol –II). NewYork: Academic Press.
- Roonwal, M.L. (1936). Studies on the embryology of the African Migratory Locust, *Locusta migratoria migratorioides*, I. *Phil. Trans. R. Soc. B.* 226, 391-421.
- Roonwal, M.L. (1937). Studies on the embryology of the African Migratory Locust, *Locusta migratoria migratorioides*, II. *Phil. Trans. R. Soc. B.* 227, 175-244.
- Rubio, R.O., Suzuki,A., Mitsumasu,K., Homma,T., Niimi,T., Yamashita, O. & Yaginuma, T. (2011).Cloning of cDNAs encoding sorbitol dehydrogenase-2a and b enzyme characterization and up regulated expression of the gene in *Bombyx mori* diapause eggs exposed to 5°C. *Insect. Biochem.Mol. Biol.*, 12 (4),361-366.
- Ruuge, E.K., Ledenev, A.N.; Lakomkin, V.L.; Konstantinov, A.A., & Ksenzenko, M.Y.(1991). Free radical metabolites in myocardium during ischemia and reperfusion. *Am. J. Physiol., Suppl.*, 261, 81-86.
- Sahu, M., Mukherjee, P.K., & Mondal, T. (1998, Nov. 11-15) Constrains during pre seed crop rearing of *Antheraea assama*, Ww. at seed research level in lower Assam. Paper presented at the 3rd *International Conference on wild silk moths*, Bhubaneswar, Orissa.
- Saito, M., Yamazaki, M., & Kobayashi, M. (1963). Steroid biosynthesis in the silk worm. *Nature*, 198, 1324.
- Sakano, D., Furusawa, T., Sugimura, Y., Storey, J.M., & Storey, K.B. (2004). Metabolic shifts in carbohydrate metabolism during embryonic development of non diapause eggs of silk worm *Bombyx mori*, *J. Insect Biotech. & Sericol.*, 73, 15-22.

- Salt, R.W. (1961). Principles of cold hardiness. *Annu. Rev. Ent.*, 6, 55- 74.
- Sander, K. (1976), Specification of basic body pattern in Insect embryogenesis. *Adv. Insect Physiol.* 12, 125-238.
- Sanders,E.J. (1975). Aspects of furrow membrane formation in the cleaving *Drosophila* embryo. *Cell. Tiss. Res.* 156, 463-474.
- Satio, S. (1937). *J. Fac. Agric. Hokkaido Imp Univ.* , 40, 35-109.
- Saunders , D. S. (1982). *Insect clocks*. Oxford: Pergamon Press.
- Seidiel, F. (1961). Entwicklungsphysiologisch Zentren im Eisystem der Iusekten. *Verh. dt. zool. Ges.*, 121-142.
- Sengupta, K. & Singh, K.(1974). Some studies on the effect of refrigeration of muga *Antheraea assama* Ww. cocoons on moth emergence, egg laying and hatching of eggs. Paper presented at the *First International Seminar on Non- mulberry silks* (pp.174). CTR & TI, Ranchi.
- Shingleton, A.W., Geoffroy, C., Sisk & David L Stern . Diapause in the pea aphid phia (*Acysthosphonpisum*) is a slowing but not a cessation of development. *BMC Developmental Biology*,
- Shinmyo, Y., Mito, T., Matsushita, T., Sarashina, I.,Miyasaki, K., Ohuchi, H.,& Noji, S. (2005). Caudal is required for gnathal and thoracic patterning and for posterior elongation in the intermediate germband in cricket *Gryllus bimaemlatus*. *Mech. Dev.*, 122, 236 – 239.
- Singh, S.P., & Singh, J. (1980). Carbohydrate turn over in relation to trehalase activity in eri silk moth *Philosamia ricini*, during embryogenesis. *Acta Physiol. Acad. Sci. Hung.*, 55,149-152.
- Singha, B.B., Sahu, A.K., & Das, P.K. (1998, Nov. 11-14). *Identification of low temperature resistant embryonic stages in muga silkworm, Antheraea assama*. Ww. Paper presented at the 3rd International conference on wild silk moth, Bhubaneswar, Orissa.

- Sinha, A.K., Sinha, U.S.P., & Sengupta, K. (1989). Changes in the free amino acids during embryonic development of *Antheraea proylei*. *Indian J. Sericulture*, 28(1), 118-120.
- Sinha, A.K., Sinha, U.S.P., Sengupta, A.K., & Sengupta, K. (1987). Changes in free amino acid in the larval and pupal haemolymph of *Antheraea proylei*, J. Paper presented at the XV International Sericultural Congress, Thailand
- Sinha, U.S.P., Sinha, A.K. & Sinha, S.S.(1991). Changes in concentration of proteins and carbohydrates in the developing healthy and pebrine infected embryos of tropical tasar silk worm, *Antheraea mylitta*, Drury. *Indian J. Seric.*, 30(2), 155-156.
- Snobbe, H., Masumoto, T., Tokushige, H., & Makka, T.(1997). Developmental changes in accumulation and metabolism of ecdysteroids in diapause eggs and non diapause eggs of silk worm *Bombyx mori*. In S. Kawashima, & S. Kikuyama (Eds.), *Advances in Comparative Endocrinology* (Vol. I, pp 185-189). Bologna: Monduzzi Editorem.
- Sohal,R.S., Aggarwal, S., Dubey, A., & Orr, W.C. (1993). Protein oxidative damage associated with life expectancy in house flies. *Proc. Natl. Acad. Sci. USA*, 90, 7255-7529.
- Sonobe, H., Maotaini, K., & Nakajima, H. (1986). Studies on embryonic diapause in the PND mutant of the silkworm, *Bombyx mori*, Genetic control of embryogenesis. *J Insect Physiol.*, 32, 215-220.
- Sonobe, H., & Yamada, R. (2004). Ecdysteroids during early embryonic development in silk worm *Bombyx mori*: metabolism and functions. *Zool. Sci.*, 21, 503-516.
- Sonobe, H., Tokushige, Makka, T., Hara, N.,& Fujimoto, Y. (1999).Comparative studies of ecdysteroid metabolism between diapause eggs and non diapauses eggs of silk worm *Bombyx mori*. *Zool.Sci.*,16, 935-943.
- Srivastava, U.S. & Bahadur, I. (1961). The development of the Malpighian in *Dysdercuskoemigi* (Hemiptera). *Quarterly Journal of Microscopical Science*, 102 (3), 347 – 60.

- Steele, J. E. (1982). Glycogen Phosphorylase in insects. *Insect Biochem.*, 12, 131 – 147.
- Storey, K. B., & Storey, J. M. (1988). Freeze tolerance in animals. *Physiol. Rev.*, 68, 27-84.
- Storey, K., & Storey, J.M. (2012). Insect cold hardiness: metabolic, gene, and protein adaptation. *Can. J. Zool.*, 90, 456-475.
- Storey, K.B. (1982). Phosphofructokinase from the over wintering gall fly larva, *Eurosta solidaginis* : control of cryoprotectant polyol synthesis. *Insect Biochem.*, 12, 501-505.
- Storey, K.B., & Storey, J.M. (1991). Carbon balance and energetic of cryoprotectant synthesis in a freeze tolerant insect: responses to perturbation by anoxia. *J. Comp. Physiol.B*, 160, 77-84.
- Storey, K.B., & Storey, J.M.(1990). Biochemistry of cryoprotectants. In D. Denlinger and R.E. Lee (Eds.), *Insects at low temperature* (pp.64-93). New York: Chapman Hall.
- Stribel, H. (1960). Zur Embryoualentwicklung der Teumiten. *Acta trop.*, 17, 193-260.
- Stuart, J.A., & Brown, M.F. (2006). Energy quiescence and the cellular basis of animal life spans. *Comp. Biochem. Physiol.*, A, 143, 12-23.
- Sujuki, K., Fujita, M., & Miya, K. (1983). Changes in super cooling point of the silk worm eggs . *J. Seric. Sci. Jpn.*, 52, 185-190.
- Takami, (1969). *A general text book of the silk worm eggs*. Tokyo, Japan : Zenkoku Sanshu Kyokai.
- Takami, T. and Kitazawa, T. (1960). Eternal observation of embryonic development in the silkworm. *Sanshi Shikenjo, Kokoku, Sericult. Expt. Stn. Tech. Bull.*, 75, 1-31.
- Takei,R., & Nagashima,E. (1975). Electron microscopic investigation on the early developmental stages of diapauses and non diapauses eggs in the silk worm *Bombyx mori* L. *J. Seric. Sci. Jpn.*, 44, 118-124.
- Takesue, S., Keino, H., & Onitake, K. (1980). Blastoderm formation in the silkworm egg (*Bombyx mori* L.). *J. Embryol. Exp. Morph.*, 60, 117-124.

- Takesue, S., Onitake, K., & Keino, H. (1977). Studies on the blastoderm & germband formation during early embryogenesis in the silk worm, *Bombyx mori* L. *Zool. Magazine*, 86, 345.
- Takimi, T. (1946). *Seibutsu*, 1, 208-211.
- Tang, S. & Li, Y. (1994). primary studies on changes in quantity of phospholipids and one day hatchability of Bluish silk worm egg by different Cold -storage Treatments. *B.B.A.*, 1218(3), 366 – 374.
- Tao, H., Liu, H.J., Cheng, Y.Q., Sima, Y., Yin, W.M., & Xu, S.Q. (2015). Parental environmental exposure leads to glycometabolic disturbances that affect fertilization of eggs in Silkworm, *Bombyx mori*: parental transcript legacy. *J. Comp. Physiol.*, 185(1), 47-55.
- Tayadae, D.S. (1987). Heterosis effect on economic traits of new hybrids of silkworm, *Bombyx mori* L. *Sericologia*, 27, 301-307.
- Tayede, D.S., Jawale, M.D., & Uncheganixar, P.K. (1987). Effect of refrigeration on hatchability of eggs of multivoltine race *Bombyx mori* L. *Sericologia*, 27(2), 297.
- Tazima, Y. (1964). *The Genetics of the silkworm*. London: Academic Press.
- Tazima, Y. (1978). *The silkworm: an Important Laboratory Tool*. Tokyo: Kodausha.
- Tazima, Y. (1984). Silkworm Moths. In I.L. Mason (Eds.), *Evolution of Domesticated Animals* (pp.416 -424). London: Longman.
- Thangavelu, K., Bhagowati, A.K. & Chakraborty, A.K. (1985a). Preservation of muga cocoon for seed purposes. *Indian Silk*, 23(12), 25- 26.
- Thangavelu, K., Bhagowati, A.K., & Chakraborty, A.K. (1985). Introduction of quiescence in muga silkworm *Antheraea assama* Westwood. *Current Science*, 54(19), 1011 -1013.
- Thangavelu, K., Bhagowati, A.K. and Chakraborty, A.K (1987). Diapause potential in muga silkworm *Antheraea assama*. *Insect Sci. Let.* 8(1), 61 -63.

- Thangavelu, K., Bhagowati, A.K. and Chakraborty, A.K. (1985 b). Induction of quiescence in muga silkworm *Antheraea assama* Westwood. *Curr.Sci.*, 54(19) 1011 -1013.
- Thangavelu, K.; Chakraborty, A.K.; Bhagowati, A.K., & Isa, Md. (1988). *Handbook of muga culture*. Bangalore: Central Silk Board
- Thangavelu, K and Sahu, A.K. (1986). Further studies on the indoor rearing of muga silkworm (*Antheraea assama* Westwood) (Lepidoptera : Saturniidae). *Sericologia*, 26 (2), 215 – 224.
- Tomita, S., & Kikuchi, A. (2009). Abd – B suppresses lepidopteran proleg development in posterior abdomen. *Developmental Biology*. 328, 403 – 409.
- Toyama, K. (1896). Structure of the eggs and development of the embryos in the silkworm, *Bombyx mori*, L. *Ngakukai-hoh*, 28, 28-80.
- Toyama, K. (1902). Contribution to the Study of Silk Worm, I. On the embryology of the silk worm. *Bull. Coll. Agric. Tokyo Imp. Univ.*, 5, 73 -118.
- Trevlyan, W.E., & Harrison, J.S. (1952). Yeast metabolism: fractionation and microdetermination of cell carbohydrate. *Biochem. J.*, 50, 298-303.
- Tricohniroff, A. (1879). Uber die entwicklungsgeschichte des Seidenwurm. *S. Zool. Anz.*, 2, 64-67.
- Turner, F.R., & Mahowald, A.P. (1976). Scanning electron microscopy of *Drosophila* embryogenesis. I. The Structure of the egg envelop & formation of the cellular blastoderm. *Dev. Biol.* 50, 95-108.
- Upadhyay, V.B., & Pandey, A.K. (2000). Effect on refrigeration on eggs and pre refrigeration period on the survival of larvae of *Bombyx mori*. *Ind. J. Ent.*, 62(1), 28-33.
- Vemenanda Reddy, G., Veeraiah, T.M., & Samson, M.V. (2004). Silk worm seed production schedules for bivoltines-New dimensions. *Indian J. Seric.* 43(1), 25-34.
- Viswakarma, S.R. (1982 – 83) Effects of Refrigeration of Erik Silkworm, *Philosamia ricini* Hutt. eggs on hatching (Lep. Sat.). *Indian J. Seric.*, 36 – 39.

- Waddington, C.H. (1967). *Principles of Development and Differentiation*. New York: Macmillan.
- Waddington, C.H. (1962). *New patterns in genetics and development*. New York: Columbia University Press.
- Walters, F.L. (1967). Effects of Temperature on Hatching & Development of *Tribolium confusum*. *J. Stored Products*, Referral to in Howe, R.H. *Annu. Rev. Ent.*, 12, 15- 42.
- Wigglesworth, V.B. (1972). *The principles of insect physiology*. London: Methuen & Co.
- Wyatt, G.R. (1967). Biochemistry of sugars and polysaccharides in insects. In J.W. Beament, J.E. Treherne and V.B. Wigglesworth (Eds.), *Advances in Insect Physiology* (pp. 287-360). London & New York : Academic Press.
- Wyatt, G.R., & Kalf, G.F. (1957). The chemistry of insect haemolymph. *J. Gen. Physiol.*, 40, 833.
- Yaginuma, O. & Yamashita, T. (1979). NAD-dependent sorbitol dehydrogenase activity in relation to the termination of diapause in eggs of *Bombyx mori*. *Insect Biochem.*, 9(5), 547- 553.
- Yaginuma, T., & Yamashita, O. (1978). Polyol metabolism related to diapauses in *Bombyx* eggs : different behaviour of sorbitol from glycerol during diapause and post diapause. *J. Insect. Physiol.*, 24, 347-354.
- Yaginuma, T., & Yamashita, O. (1979). NAD- dependent sorbitol dehydrogenase in relation to the determination of diapauses in eggs of *Bombyx mori*. *Insect. Biochem.*, 9(5), 533-547.
- Yaginuma, T., & Yamashita, O. (1999). Oxygen consumption in relation to sorbitol utilization at the termination of diapauses in eggs of the silk worm *Bombyx mori*. *J. Insect Physiol.*, 44, 621-627.
- Yaginuma, T., Kobayashi, M., & Yamashita, O. (1990a). Effect of low temperatures on NAD sorbitol dehydrogenase activity and morphogenesis in non diapauses eggs of the silk worm *Bombyx mori*. *Comp. Biochem. Physiol.*, B. 97 (3), 495-506.

- Yaginuma, T., Kobayashi, M., & Yamashita, O. (1990b). Distinct effects of different low temperatures on the induction of NAD sorbitol dehydrogenase activity in diapause egg of the silk worm of *Bombyx mori*. *J. Comp. Physiol.*, B, 160, 277-285.
- Yamada, R., & Sonobe, H. (2003). Purification, kinetic characterization and molecular cloning of novel enzyme ecdysteroid- phosphatase. *J. Biol.Chem.*, 278, 26365-26373.
- Yamahama, Y., Seno, K., & Hariyama, T. (2008). Changes in lipid droplet localization during embryogenesis of the silk worm *Bombyx mori*. *Zool.Sci.*, 25, 580-586.
- Yamashita, O. (1965). Carbohydrate metabolism during embryonic development of the silk worm, *Bombyx mori*, L. *J. Seric. Sci. Jpn.*, 34, 1-7.
- Yamashita, O. and Hasegawa, K. (1985). Embryonic diapause. In G. A. Kerkut, & L.I. Gilbert (Eds.), *Comprehensive Insect Physiology Biochemistry and Pharmacology* (pp. 407-434). Oxford: Pergamon Press.
- Yamashita, O., & Hasegawa, K. (1976). Diapause hormone action in silkworm ovaries incubated *in vitro* ¹⁴C- trehalose incorporation into glycogen. *J. Insect Physiol.*, 22(3),409-414.
- Yamashita, O., & Yaginuma, T. (1991). Silk worm eggs at low temperatures: implications for sericulture. In R.E. Lee & Denlinger, D.L. (Eds.), *Insects at low temperature* (pp. 424-445). New York : Chapman Hall.
- Yamashita, O., & Yaginuma, T. (1991). Silk worm eggs at low temperatures: implications for sericulture. In R.E. Lee & Denlinger, D.L. (Eds.), *Insects at low temperature* (pp. 424-445). New York : Chapman Hall.
- Yamato, K., Banno, Y., Fujji, H., Maie, F., Kashige, N., & Aso, Y. (2006). Catalase from the silk worm *Bombyx mori*: gene sequence, distribution and over expression. *Insect Biochem. Mol. Biol.*, 35, 277-283.
- Yuyin, C. (2000). Technical standards for commercial egg production of bivoltine silkworm varieties used in China. *Sericologia*, 40,185-201.

- Zhao, L.C., Hou, Y.S., Sima, Y.H.(2014). Changes in glutathione redox cycle during diapause determination and termination in the bivoltine silk worm *Bombyx mori*. *Insect Science*, 21 (1), 39-46.
- Zhao,L.C., & Shi, L.G. (2010). Metabolism of hydrogen peroxide between diapauses and non diapauses eggs of the silk worm *Bombyx mori* during chilling at 5°C. *Arch. of Insect Biochem. Physiol.*, 74 (2), 127-134.
- Zhu, I., Indrasiith, L.S., & Yamashita, O. (1986). Characterization of vitellin, egg specific protein and 30- kDa proteins of *Bombyx* eggs and their fates during oogenesis and embryogenesis. *Biochem. Biophys. Acta*, 882, 427-436.