

## BIBLIOGRAPHY

### **CHAPTER 1:**

- [1] G. Durand and J. D. Lister; Ann. Rev. Mater. Sci., 23, 269 (1973), Recent advances in liquid crystals.
- [2] T. J. Sluckin, D. A. Dunmur and H. Stegemeyer, Crystals That Flow: Classic Papers from the History of Liquid Crystals, History of Physics Group Newsletter, Institute of Physics (2006).
- [3] O. Lehmann; Z. physik. Chem. 4, 462- 472 (1889), Fließende Kristalle.
- [4] F. Reinitzer; Monatshefte für Chemie., 9, 421-441(1888), Beiträge Zur Kenntnis des Cholesterins.
- [5] G. Freidel; Ann. Physique (Paris), 18, 273 (1922), Les états mésomorphes de la matière.
- [6] V. Vill; Liq. Cryst 3.4 Database, LC publisher GmbH, hamburg, May (2000).
- [7] H. Kelker and R. Hatz; Verlag Chemie, Chapter 1 (1980), Hand Book of Liquid Crystals.
- [8] H. Kelker; Mol. Cryst. Liq. Cryst., 21, 1 (1973), History of liquid crystals.
- [9] H. Kelker and P. M. Knoll; Liquid. Cryst. 5(1), 19 (1989), Plenary Lecture: Some pictures of the history of liquid crystals.
- [10] G. W. Gray; Mol. Cryst. Liq. Cryst., 63, 3 (1981), Comments on some recent development in the field of liquid crystals.
- [11] G. W. Gray; Liq. Cryst., 24(1), 5 (1998), Reminiscences from a life with liquid crystals.
- [12] P. J. Coolings and M. Hird, Introduction to Liquid Crystals Chemistry and Physics, Taylor & Francis (1998).
- [13] G. W. Gray and H. Finkelmann in Polymeric Liquid Crystals, ed. A. Ciferri, W. R. Krigbaum and Robert B. Meyer, AP, chap-1 and chap-2 respectively (1982).
- [14] D. Demus; Mol. Cryst. Liq. Cryst., 364, 25 (2001), One century liquid crystal chemistry: from Vorlander's rods to disks, stars and dendrites and G. W. Gray in Handbook of Liquid Crystals. Vol. 1 (Fundamentals), Ed. D. Demus, J. W. Goodby, G. W. Gray, H.-W. Spiess and V. Vill, Introduction and Historical Development, WILEY-VCH, Verlag GmbH, Weinheim, FRG (1998).
- [15] S. Chandrasekhar, B. K. Sadashiva and K. A. Suresh; Pramana, 9, 471- 480 (1977), Liquid crystals of disc-like molecules.
- [16] J. Billard, J. C. Dubois, T. H. Nguyen and A. Zann; Nouveau J. Chim. 2, 535-540 (1978), Une mésophase disquotique.
- [17] Liquid Crystals Application and Uses Vol. 1-3. Ed. B. Bahadur, World Scientific, (1990-1992).
- [18] G. W. Brown, J. W. Doane and V. D. Neff, A Review of the Structure and Physical Properties of Liquid Crystals. CRC Press, Cleveland, Ohio (1971).
- [19] J. W. Goodby; Liq. Cryst. 24, 25-38 (1998), Liquid crystals and life.
- [20] P. G. de Gennes; "The Physics of Liquid Crystals", Clarendon Press, Oxford (1974); P. G. de Gennes and J. Prost, 2nd ed. Clarendon Press, Oxford (1993).
- [21] L. A. Madsen, T. J. Dingemans, M. Nakata, E. T. Samulski; Phys. Rev. Lett. 92 (14) (2004), Thermotropic Biaxial Nematic Liquid Crystals.

- [22] A. de Vries; Mol. Cryst. Liq. Cryst., 10, 31 (1970), Evidence for the Existence of More Than One Type of Nematic Phase.
- [23] A. de Vries; Mol. Cryst. Liq. Cryst., 10, 219 (1970), X-Ray Photographic Studies of Liquid Crystals I. A Cybotactic Nematic Phase and A. de Vries, in Proc. of the International Liquid Crystal Conference, Bangalore, December, Pramana Supplement I, 93 (1973), X-Ray Studies of Liquid Crystals: V. Classification of Thermotropic Liquid Crystals and Discussion of Intermolecular Distances.
- [24] P. E. Cladis; Phys. Rev. Lett., 35, 48-51 (1975), New Liquid-Crystal Phase Diagram.
- [25] G. W. Gray, Molecular Structure and the Properties of Liquid Crystals, Academic Press, London (1962).
- [26] P. S. Pershan, Structure of Liquid Crystals Phases, World Scientific Lecture Note in Physics, Vol 23, World Scientific, Singapore (1988).
- [27] G. W. Gray and J. W. Goodby, Eds., Smectic Liquid Crystals, Textures and Structures, Leonard - Hill, Philadelphia (1984).
- [28] A. J. Leadbetter in Thermotropic Liquid Crystals, Critical Reports on Applied Chemistry, Vol. 22, G. W. Gray(Ed.), Wiley, Chichester, p1 (1987).
- [29] A. de Vries in Liquid Crystals, The Fourth State of Matter (Ed. F. D. Saeva), Marcell Dekker, New York, p1 (1972).
- [30] A. de Vries, Mol. Cryst. Liq. Cryst., 63, 215(1981), A Structural Classification of Smectic Liquid Crystals.
- [31] J. Doucet in The Molecular Physics of Liquid Crystals, Eds. G. R. Luckhurst and G. W. Gray, Academic Press, London, p317 (1979).
- [32] L. V. Azaroff, Mol. Cryst. Liq. Cryst., 60, 73(1980), X-Ray Diffraction by Liquid Crystals.
- [33] D. Demus, J. W. Goodby, G. W. Gray and H. Sackmann, Mol. Cryst. Liq. Cryst., 56, 311(1980), Recommendation for The Use of The Code Letters G And H for Smectic Phases.
- [34] J. W. Goodby and G. W. Gray, Mol. Cryst. Liq. Cryst., 41, 145-150 (1978), Some New Smectic F Materials.
- [35] R. M. Richardson, A. J. Leadbetter and J. C. Frost, Mol. Phys., 45, 1163 (1982), A comparative study of the molecular motions in the three smectic phases of isobutyl 4(4' phenylbenzylideneamino) cinnamate using incoherent neutron scattering.
- [36] A. J. Leadbetter, J. C. Frost, J. P. Gaughan, and M. A. Mazid, J. Phys. (Paris), 40, C3-185-C3-192 (1979), The structure of the crystal, smectic E and smectic B forms of IBPBAC.
- [37] J. W. Goodby in Handbook of Liquid Crystals, Eds. D. Demus, J. Goodby, G.W. Gray, H.-W. Spiess and V. Vill, Vol. 2A, Wiley-VCH, Verlag GmbH, Weinheim, FRG, p7 (1998).
- [38] F. Hardouin, A. M. Levelut, J. Bennattar and G. Sigaud; Solid State Comm., 33, 337-340 (1980), X-rays investigations of the smectic A1 - smectic A2 transition.
- [39] A. J. Leadbetter, J. L. Durrant and M. Rugman; Mol. Cryst. Liq. Cryst. Lett., 34, 231 (1977), The Density Of 4 n-Octyl-4-Cyano-Biphenyl (8CB).
- [40] A. J. Leadbetter, J. C. Frost, J.P. Gaughan, G.W. Gray and A. Mosley; J. Phys. (Paris), 40, 375-380 (1979), The structure of smectic A phases of compounds with cyano end groups.
- [41] F. Hardouin, G. Sigaud, N. H. Tinh, A.F. Achard; J. Phys. (Paris) Lett., 42, 63-66 (1981), A fluid smectic A antiphase in a pure nitro rod-like compound.

- [42] J. Prost; *Advances in Physics*, 33, 1-46 (1984), The smectic state.
- [43] P. S. Pershan, *Structure of Liquid crystals*, World Scientific, Singapore (1988).
- [44] B. R. Ratna, R. Shashidhar and V. N. Raja, *Phys. Rev. Lett.*; 55(14), 1476 (1985), Smectic-A phase with two collinear incommensurate density modulations.
- [45] S. T. Wang, S. L. Wang, X. F. Han, Z. Q. Liu, S. Findeisen, W. Weissflog and C. C. Huang; *Liq. Cryst.*, 32, 5, 609–617 (2005), Optical studies on two tilted smectic phases of meta-substituted three-ring liquid crystal compounds.
- [46] A. L. Tykalo, *Thermophysical properties of liquid crystals*, Gordon and Breach Science Publishers (1991).
- [47] R. Pindak, D. E. Moncton, S. C. Davey and J. W. Goodby; *Phys. Rev. Lett.*, 46, 1135 (1981), X-Ray Observation of a Stacked Hexatic Liquid-Crystal B Phase.
- [48] D. E. Moncton and R. Pindak in *Ordering in Two Dimensions* (Ed. S. K. Sinha), North Holland, New York, p83 (1980).
- [49] A. J. Leadbetter, J. C. Frost, M. A. Mazid; *J. Phys. (Paris) Lett.*, 40, 325-329 (1979), Interlayer correlations in smectic B phases.
- [50] P. S. Pershan, G. Aeppli, J. D. Litster and R. J. Birgeneau; *Mol. Cryst. Liq. Cryst.*, 67, 205(1981), High-Resolution X-ray Study of the Smectic A-Smectic B Phase Transition and the Smectic B Phase in Butyloxybenzylidene Octylaniline.
- [51] D. E. Moncton, and R. Pindak; *Phys. Rev. Lett.*, 43, 701(1979), Long-Range Order in Two- and Three-Dimensional Smectic-B Liquid-Crystal Films.
- [52] P. A. C Gane, A. J. Leadbetter and P.G. Wrighton; *Mol. Cryst. Liq. Cryst.*, 66, 247-266 (1981), Structure and Correlations in Smectic B, F and I Phases.
- [53] S. Diele, D. Demus and H. Sackmann; *Mol. Cryst. Liq. Cryst. Lett.* , 56, 217 (1980), X-Ray Studies in the n-Pentyl 4-[4-n-Dodecyl-Oxybenzylidene-Amino]-Cinnamate in the S<sub>I</sub> Phase.
- [54] J. J. Benattar, F. Moussa, M. Lambert; *J. Phys. (Paris) Lett.*, 42, 67-70 (1981), Two kinds of two-dimensional order : the SmF and SmI phases.
- [55] A. M. Levelut, J. Doucet and M. Lambert; *J. Phys., (Paris)*, 35, 773-779 (1974), X-ray study of the nature of the smectic B mesophases and of the solid-smectic B phase transition.
- [56] J. Doucet, P. Keller, A.M. Levelut and P. Porquet; *J. Phys. (Paris)*, 39, 548-553 (1978), Evidence of two new ordered smectic phases in ferroelectric liquid crystals.
- [57] A. J. Leadbetter, M. A. Mazid, R. M. Richardson, in *Liquid Crystals*, Ed. S. Chandrasekhar, Heyden, London, p65 (1980).
- [58] J. L. Ferguson; *Sci. Am.*, 211, 76-85 (1964), Liquid Crystals.
- [59] D. Demus, J. Goodby, G.W. Gray, H.-W. Spiess and V. Vill, *Handbook of Liquid Crystals*, Ch. IV and V, Vol. 2A, Wiley-VCH, Verlag GmbH, Weinheim, FRG (1998).
- [60] F. D. Saeva; *Mol. Cryst. Liq. Cryst.*, 23, 171-177 (1973), On the Relationship between Cholesteric and Nematic Mesophases.

- [61] T. Nakagiri, H. Kodama and K.K. Kobayashi; Phys. Rev. Lett., 27, 564-567 (1971), Helical Twisting Power in Mixtures of Nematic and Cholesteric Liquid Crystals\*.
- [62] I. G. Chistiyakov; Sov. Phys. Usp., 9, 551-573 (1967), Liquid Crystals.
- [63] A. Saupe; Angew. Chem. Intnl. Edn. 7, 97-112 (1968), Recent Results in the Field of Liquid Crystals.
- [64] A. Hochbaum, Fluorescent Guest-Host in Scattering Liquid crystals and New Physical Phenomena in Thin Liquid Crystal Films (Ph.D. Thesis), Temple University, p.5 (1980).
- [65] J. J. Wysocki, J. Adams and W. Hasse; Phys. Rev. Lett., 20, 1024-1025 (1968), Electric-Field-Induced phase change in cholesteric liquid crystals.
- [66] H. Baessler and M. M. Labes; Phys. Rev. Lett., 21, 1791-1793 (1968), Relationship Between Electric Field Strength and Helix Pitch in Induced Cholesteric-Nematic Phase Transitions.
- [67] O. Lehmann; Z Phys. Chem, 56, 750 (1906).
- [68] Harry J Coles, Pivnenko, N. Mikhail; Nature 436 (7053), 997–1000 (2005), Liquid crystal 'blue phases' with a wide temperature range.
- [69] Jun Yamamoto, Isa Nishiyama, Miyoshi Inoue, Hiroshi Yokoyama; Nature, 437 (7058): 525 (2005), Optical isotropy and iridescence in a smectic blue phase.
- [70] R. M. Hornreich and S. Shtrikman; J. Physique, 41, 335-340 (1980), A body-centered cubic structure for the cholesteric blue phase.
- [71] H. Kikuchi, M. Yokota, Y. Hisakado, H Yang, T. Kajiyama; Nature Materials, 1 (1): 64–8 (2002), Polymer-stabilized liquid crystal blue phases.
- [72] R. B. Meyer, L. Liebert, L. Strzelecki and P. Keller; J. Phys. Lett. 36, L69-L71 (1975), Ferroelectric liquid crystals.
- [73] S. Pirke and M. Glogorova in Ferroelectric – Physical Effects; Ed, M. Lallart, INTECH, Rijeka, Croatia (2011).
- [74] S. Pirkl and M. Glogarova; Intech, Croatia (2011), Ferroelectric liquid crystals with high spontaneous polarization.
- [75] R. B. Meyer; Mol. Cryst. Liq. Cryst., 40, 33-48 (1977), Ferroelectric liquid crystals: A review.
- [76] Yu. P. Panarin, O. Kalinovskaya & J. K. Vij; Liquid Crystals, Vol. 25, No. 2, 241- 252 (1998), The investigation of the relaxation processes in antiferroelectric liquid crystals by broad band dielectric and electro-optic spectroscopy.
- [77] A. D. L. Chandani, E. Gorecka, Y. Ouchi, H. Takezoe, and A. Fukuda; Jpn. J. Appl. Phys. 28(7), L1265 (1989), Antiferroelectric Chiral Smectic Phases Responsible for the Trislable Switching in MHPOBC.
- [78] H. Hiraoke, A. D. L. Chandani, E. Gorecka, Y. Ouchi, H. Takezoe and A. Fukuda; Jpn. J. Appl. Phys. 29(8), L1473 (1990), Electric-Field-Induced Transitions among Antiferroelectric, Ferrielectric and Ferroelectric Phases in a Chiral Smectic MHPOBC.
- [79] J. P. F. Lagerwall, P. Rudquist, S. T. Lagerwall and F. Gießelmann; Liq. Cryst., 30, 399-414 (2003), On the phase sequence of antiferroelectric liquid crystals and its relation to orientational and translational order.
- [80] K. Miyachi and A. Fukuda, in Handbook of Liquid Crystals, Eds. D. Demus, J. Goodby,

- G.W. Gray, H.-W. Spiess and V. Vill, Vol. 2B, Wiley-VCH, Verlag GmbH, Weinheim, FRG, Ch. 3 (1998).
- [81] J. Dijon in Liquid Crystal Application and Uses (Ed. B. Bahadur), Vol. I, World Scientific, Singapore, p 305 (1990).
- [82] N. A. Clark and S. T. Lagerwall; *Appl. Phys. Lett.*, 36, 899-901(1980), Submicrosecond bistable electrooptic switching in liquid crystals.
- [83] D. Armitage, J. I. Thakara and W. D. Eades; *Ferroelectrics*, 85, 291-302 (1988), Ferroelectric liquid-crystal devices and optical processing.
- [84] M. Hird; *Liq. Cryst.*, 38, 1467–1493 (2011), Invited topical review (Ferroelectricity in liquid crystals - materials, properties and applications).
- [85] A. N. Cammidge and R. J. Bushby in *Handbook of Liquid Crystals*, Eds. D. Demus, J. Goodby, G.W.Gray, H.-W. Spiess and V. Vill, Vol. 2B, Wiley-VCH, Verlag GmbH, Weinheim, FRG, Ch.VII (1998).
- [86] S. Kumar; *Liquid Crystals*, 31, 1037-1059 (2004), Recent developments in the chemistry of triphenylene-based discotic liquid crystals.
- [87] M. Schadt and W. Helfrich; *Appl. Phys. Lett.* 18(4), 127 (1971), Voltage-dependent optical activity of a twisted nematic liquid crystal.
- [88] T. J. Scheffer and J. Nehring; *Appl. Phys. Lett.* 45(10), 1021 (1984), A new, highly multiplexable liquid crystal display.
- [89] H. Xu, D. Hartono and K. -L. Yang, *Liquid Crystals*, 37 (10), 1269–1274 (2010), Detecting and differentiating *Escherichia coli* strain TOP10 using optical textures of liquid crystals.
- [90] J. Akaogi, Y. Koizumi, M. Ono and H. Furue; *Mol. Cryst. Liq. Cryst.*, 596, 106–112 (2014), Application of Ferroelectric Liquid Crystals to Optical Devices.
- [91] J. S. Patel and Seong-Woo Suh; *Journal of the Korean Physical Society*, 32, S1048-S1051 (1998), Ferroelectricity in Liquid Crystals and Its Applications.
- [92] C. L. Khetrapal in *Liquid Crystals in the Nineties and Beyond*, Ed. S. Kumar, World Scientific, p 435 (1995).
- [93] B. Bahadur in *Handbook of Liquid Crystals*, Eds. D. Demus, J. Goodby, G.W. Gray, H.-W. Spiess and V. Vill, Vol. 2A, Wiley-VCH, Verlag GmbH, Weinheim, FRG, p257 (1998).

## CHAPTER 2:

- [1] M. Born; Sitzb. kgl. preub. Akad. Wiss., 614 (1916), Theorie der flüssigen Kristalle und des elektrischen Kerr Effects in Flüssigkeiten.
- [2] P. G. de Gennes; The Physics of Liquid Crystals, Clarendon Press, Oxford (1974); P. G. de Gennes and J. Prost, 2nd ed. Clarendon Press, Oxford (1993).
- [3] A. L. Tsikalo; Thermophysical properties of liquid crystals, Gordon and Breach Science Publishers, (1991).
- [4] S. Chandrasekhar, Liquid Crystals, 2nd Edn., Cambridge University Press, Cambridge, (1992).
- [5] I. Musevic, R. Blinc and B. Zeks's, The Phases of Ferroelectric and Antiferroelectric Liquid Crystals, World Scientific, Singapore (2000).
- [6] W. Maier and A. Saupe, Z. Naturforsch 13a, 564 (1958), Eine einfache molekulärstatistische Theorie der nematischen kristallinflüssigen Phase, Dispersionswechselwirkung; W. Maier, and A. Saupe, Z. Naturforsch, 14a, 882 (1959), Eine einfache molekulärstatistische Theorie der nematischen kristallinflüssigen Phase. Teil I; W. Maier, and A. Saupe, Z. Naturforsch, 15a, 287 (1960), Eine einfache molekulärstatistische Theorie der nematischen kristallinflüssigen Phase. Teil II; A. Saupe and W. Maier, Z. Naturforsch, 16a, 816 (1961), Methoden zur Bestimmung des Ordnungsgrades nematischer Schichten; W. Maier, Angew. Chem. 73, 660 (1961), Struktur und Eigenschaften nematischer Phasen.
- [7] A. L. Tsikalo, Thermophysical Properties of Liquid Crystals, Gordon and Breach Science Publishers, p27 (1991).
- [8] K. K. Kobayashi; Phys. Lett. A, 31, 125-126 (1970), On the theory of translational and orientational melting with application to liquid crystals.
- [9] W. L. McMillan; Phys. Rev. A 4, 1238-1246 (1971), Simple molecular model for the smectic A phase of liquid crystals.
- [10] W. L. McMillan; Phys. Rev. A 6, 936 (1972), X-Ray Scattering from Liquid Crystals. I. Cholestryl Nonanoate and Myristate.
- [11] A. L. Tsikalo, Thermophysical Properties of Liquid Crystals, Gordon and Breach Science Publishers, p 164 (1991).
- [12] R. J. Meyer and W. L. McMillan; Phys. Rev. A, 9, 899 (1974), Simple molecular theory of the smectic C, B, and H phases.
- [13] S. Chandrashekhar; Liquid crystals, Cambridge University Press (1977).
- [14] G. Vertogen and W. H. de Jeu; Thermotropic Liquid crystals, Fundamentals. Springer-Verlag, Berlin (1988).
- [15] P. J. Collings and Michael Hird; Introduction to Liquid Crystals Chemistry and Physics. Taylor and Francis (1998).
- [16] G. R. Luckhurst; Molecular Physics of Liquid Crystals, Academic Press (1979).
- [17] R. L. Humphries, P. G. James and G. R. Luckhurst; J. Chem. Soc., Faraday Trans. 2, 68, 1031 – 1044 (1972), Molecular field treatment in nematics.

- [18] K. K. Kobayashi; J. Phys. Soc. (Japan), 29, 101 (1970), Theory of Translational and Orientational Melting with Application to Liquid Crystals I.
- [19] K. K. Kobayashi; Mol. Cryst. Liq. Cryst., 13, 137 (1971), Theory of Translational and Orientational Melting with Application to Liquid Crystals.
- [20] B. K. Vainstein; Diffraction of X-rays by Chain Molecules, Elsevier, Amsterdam (1966).
- [21] H. Kelkar and R. Hatz; Handbook of Liquid Crystals, Verlag Chemie, Ch. 5 (1980).
- [22] P. S. Pershan; Structure of Liquid Crystalline Phases, World Scientific, Singapore (1988).
- [23] G. Ungar in Physical properties of liquid crystals, D. A. Dunmur, A. Fukuda and G. R. Luckhurst (Eds.), INSPEC, London, Ch. 4.1 (2001).
- [24] G. Vertogen and W. H. de Jeu (Eds.), Thermotropic Liquid Crysatsals Fundamentals., Springer-Verlag, p-207 (1988).
- [25] G. R. Luckhurst in Liquid Crystals & Plastic Crystals (Eds. G. W. Gray and P. A. Winsor), Ellis Horwood Limited, Vol 2, p-144 (1974).
- [26] C. L. Khetrapal and A. C. Kunwar in Advances in Liquid Crystals, Vol. 1-6, Ed. Glenn H. Brown, p-173, AP (1983).
- [27] V. D. Neff in Liquid Crystals & Plastic Crystals (Ed. G. W. Gray and P. A. Winsor), Ellis Horwood Limited, Vol 2, p-231 (1974).
- [28] S. J. Gupta, R. A. Gharde and A. R. Tripathi; Mol. Cryst. Liq. Cryst., 364, 461 - 468 (2001), Phase Transition Temperatures of LCs using Fabry-Perot Etalon.
- [29] D. Demus, Textures of liquid crystals, Verlag Chemie Weinheim, New York (1978).
- [30] I. Dierking, Textures of liquid crystals WILEY-VCH, GmbH & Co. KGaA p-16 (2003).
- [31] A. J. Slaney, K. Takatohi and J. W. Goodby in The Optics of Thermotropic Liquid Crystals, Ed. S. Elston and R. Sambles, Taylor & Francis (1998), Defect textures in liquid crystals.
- [32] Y. Bouligand in Handbook of Liquid Crystals. Vol. 1 (Fundamentals), Ed. D. Demus, J. Goodby, G. W. Gray, H.-W. Spiess and V. Vill, WILEY-VCH, Verlag GmbH, Weinheim, FRG (1998), Defects and textures.
- [33] B. Jha and R. Paul; Proc. Nucl. Phys. and Solid State Phys. Symp., India . 19C, 491 (1976) A Design for High Temperature X-ray Camera for the Study of Liquid Crystals in a Magnetic Field.
- [34] B. Jha, S. Paul, R. Paul and P. Mandal; Phase Transitions, 15, 39-48 (1989), Order parameters of some homologue cybotactic nematics from X-ray diffraction measurements.
- [35] S. Diele. P. Brand and H. Sackmann; Mol. Cryst. Liq. Cryst., 16, 105-116 (1972), X-ray Diffraction and Polymorphism of Smectic Liquid Crystals 1. A-, B- and C-modifications.
- [36] A. de Vries; Pramana, Suppl. No. 1, 93 (1975), X-Ray Studies of Liquid Crystals: V, Classification of Thermotropic Liquid Crystals and Discussion of Intermolecular Distances.

- [37] A. de Vries, A. Ekachi and N. Sielberg; *J. de Phys.*, 40, C3-147 (1979), X-Ray Studies of Liquid Crystals VI. The Structure of the Smectic A, C, Bn and Bt Phases of Trans-1, 4-Cyclohexane-DI-NOctyloxybenzoate.
- [38] A. J. Leadbetter, J. Prost, J. P. Gaughan and M. A. Mazid; *J. de Phys.*, 40, C3-185 – C3-193 (1979), The structure of the crystal, SmE and SmB forms of IBPBAC.
- [39] A. J. Leadbetter and P. G. Wrighton; *J. de Phys.*, 40, C3-234 – C3-242 (1979), Order parameters in S<sub>A</sub>, S<sub>C</sub>, and N phases by X-ray diffraction.
- [40] V. M. Sethna, A. de Vries and N. Spielberg; *Mol. Cryst. Liq. Cryst.*, 62, 141-153 (1980), X-Ray Studies of Liquid Crystals VIII: A Study of the Temperature Dependence of the Directly Observed Parameters of the Skewed Cybotactic Nematic Phase of Some Bis-(4'-n-Alkoxybenzal) -2-Chloro-1,4-Phenylenediamines.
- [41] L.V. Azaroff and C. A. Schuman; *Mol. Cryst. Liq. Cryst.*, 122, 309-319 (1985), X-Ray Diffraction by Cybotactic Nematics.
- [42] A. de Vries; *Mol. Cryst. Liq. Cryst.*, 131, 125-145 (1985), The Use of X-Ray Diffraction in the Study of Thermotropic Liquid Crystals with Rod-Like Molecules.
- [43] P. Mandal, M. Mitra, S. Paul and R. Paul; *Liq. Cryst.*, 2, 183-193 (1987), X-Ray Diffraction and Optical Studies of an Oriented Schiff's Base Liquid Crystal.
- [44] A. de Vries; *Mol. Cryst. Liq. Cryst.*, 10, 219-236 (1970), X-ray photographic studies of liquid crystals I. A cybotactic nematic phase.
- [45] A. de Vries; *Mol. Cryst. Liq. Cryst.*, 11, 361-383 (1970), X-Ray Photographic Studies of Liquid Crystals II. Apparent Molecular Length and Thickness in Three Phases of Ethyl-p-Ethoxybenzal-p-Aminobenzoate.
- [46] C. Kittle; *Intro to Solid State Physics*, Wiley Eastern, Chapter 2 (1976).
- [47] G. Pelzl in *Handbook of Liquid Crystals*. Vol. 1 (Fundamentals), Ed. D. Demus, J. Goodby, G. W. Gray, H.-W. Spiess and V. Vill, WILEY-VCH, Verlag GmbH, Weinheim, FRG (1998).
- [48] D. A. Dunmur in *The Optics of the Thermotropic Liquid Crystals*. Ed. S. Elston and R. Sambles, Taylor & Francis (1998).
- [49] S. M. Kelly, M. O'Neill; Chapter 1, *Liquid Crystals For Electro-optic Applications*.
- [50] A. K. Zeminder, S. Paul and R. Paul; *Mol. Cryst. Liq. Cryst.* 61, 191-206 (1980), Refractive Indices and Orientational Order Parameter of Five Liquid Crystals in Nematic Phase.
- [51] M. F. Vuks; *Optics and Spectroscopy*, 20, 361-368, (1966), Determination of optical anisotropy of aromatic molecules from the double refraction of crystals.

- [52] H. E. J. Neugebauer, *J. Canad. Phys.*, 32, 1-8 (1954), Clasius-Mossoti equation for certain types of anisotropic crystals.
- [53] S. Chandrasekhar; *Liquid Crystals*, second edn., Cambridge University Press, Cambridge, p39, (1992).
- [54] W. H. de Jeu, *Physical Properties of Liquid Crystalline Materials*, Ed G. Gray, Gordon and Breach, Vol. 1, p 41 (1980).
- [55] P. G. de Gennes; *Mol. Cryst. Liq. Cryst.* 12, 193-214 (1971), Short Range Order Effects in the Isotropic Phase of Nematics and Cholesterics.
- [56] I. Haller, H. A. Huggins, H. R. Lilenthal and T. R. McGuire; *J. Phys. Chem.*, 77, 950 (1973), Order-Related Properties of Some Nematics; I. Haller; *Jappl. Phys. Lett.*, 24, 349 (1974), Alignment and wetting properties of Nematics.
- [57] K. Toriyama, K. Suzuki, T. Nakagomi, T. Ishibashi and K. Odawara; *J. de Phys.*, 40, C3-317 (1979), A design of liquid crystal material for multiplexed liquid crystal display.
- [58] D. A. Dunmur and W. H. Miller; *Mol. Cryst. Liq. Cryst.*, 60, 281-283 (1980), Dipole-Dipole Correlation in Nematic Liquid Crystals.
- [59] R. E. Michel and G. W. Smith; *J. Appl. Phys.* 45, 3234 (1974), Dependence of birefringence threshold voltage on dielectric anisotropy in a nematic liquid crystals.
- [60] C. J. F. Böttcher, *Theory of Electric Polarization*, 2<sup>nd</sup> edition, Vol I (1973) and C. J. F. Böttcher and P. Bordewijk, 2<sup>nd</sup> edition, Elsevier Scientific Publishing Co., Vol II (1978).
- [61] W. H. de Jeu and T. W. Lathouwers; *Z. Naturforsch* 29a, 905 (1974), Dielectric constants and molecular structure of nematics. I. Terminally substituted azobenzene and azoxybenzenes.
- [62] W. Maier and G. Meier, *Z. Naturforsch*; 16a, 262 (1961), Einfache Theorie der dielektrischen Eigenschaften von homogen orientierten nematischen Phasen.
- [63] L. Onsager; *J. Am. Chem. Soc.*, 58, 1486-1493 (1936), Electric Moments of Molecules in Liquids.
- [64] L. Bata and A. Buka; *Mol. Cryst. Liq. Cryst.*, 63, 307 (1981), Dielectric Permittivity and Relaxation Phenomena in smectic Phases.
- [65] H. Frohlich, *Theory of Dielectrics*. Clarendon Press, Oxford (1949); H. Frohlich, *J. Chem. Phys.*, 22, 1804 (1954).
- [66] P. Bordewijk; *Physica*, 69, 422-432 (1973), On the derivation of the Kirkwood-Fröhlich equation.

- [67] P. Bordewijk; Physica, 75, 146-156 (1974), Extension of the Kirkwood-Fröhlich theory of the static dielectric permittivity to anisotropic liquids.
- [68] P. Bordewijk and W. H. de Jeu; J. Chem. Phys. 68 (1), 116 (1978), Calculation of dipole correlation factors in liquid crystals with use of a semiempirical expression for the internal field.
- [69] W. H. de Jeu and P. Bordewijk; J. Chem. Phys. 68 (1), 109 (1978), Physiical studies of nematic azoxybenzenes. II. Refractive indices and the internal field.
- [70] W. H. de Jeu, T. W. Lathouwers and P. Bordewijk; Phys. Rev. Lett. 32, 40 (1974), Dielectric properties of n and  $S_A$  p,p-di-n-heptylazoxobenzene.
- [71] W. H. de Jeu, J. W. A. Goosens and P. Bordewijk; J. Chem. Phys. 61, 1985(1974), Influence of smectic order on the static dielectric permittivity of liquid crystals.
- [72] P. Debye, Polar molecules, Dover Pub. Inc., NX (1929).
- [73] C. J. F. Böttcher and P. Bordewijk, Ed., Theory of Electric Polarization, Vol II, Dielectric in time-dependent fields, Elsevier, Scientific Publishing Co. (1973).
- [74] S. Havriliak and S. Negami; J. Polime, 8, 161-210 (1967), A complex plane representation of dielectric and mechanical relaxation processes in some polymers.
- [75] A. Wurflinger; Int. Rev. Phys. Chem., 12, 89 (1993), Dielectric studies under pressure on plastic and liquid crystals.
- [76] W. Haase, H. Pranoto, F. J. Bormuth, Ber. Bunsenges. Phys. Chem., 89, 1229-1234 (1985), Dielectric Properties of Some Side Chain Liquid Crystalline Polymers.
- [77] F. J. Bormuth, Ph.D. Thesis, Technische Hochschule, Darmstadt (1998).
- [78] V. I. Minkin, O. A. Osipov, Y. A. Zhdanov, Dipole Moments in Organic Chemistry, New York- London, (1970).
- [79] A. K. Jonscher, Dielectric Relaxation In Solids, Chelsea Dielectric Press, London (1983).
- [80] S. Wrobel, B. T. Gowda, W. Haase, J.Chem. Phys., 106 (10), 5904 (1997), Dielectric study of phase transitions in 3-nitro-4-chloro-aniline.
- [81] V. K. Agarwal and A. H. Price; Mol. Cryst. Liq. Cryst., 98, 193-200 (1983), Dielectric Relaxation and Order Parameters in Mixtures of 4-cyanobiphenyl and MBBA.
- [82] V. V. Daniel, Dielectric Relaxation, Academic Press, London and N.Y. (1967).

- [83] K. S. Cole and R. H. Cole; J. Chem. Phys. 9, 341-351 (1941), Dispersion and Absorption in Dielectrics I. Alternating Current Characteristics.
- [84] J. Hatano, Y. Hakanai, H. Furue, H. Uehara, S. Saito, and K. Murashiro, Jpn. J. Appl. Phys., 33, 5498-5502 (1994), Phase Sequence in Smectic Liquid Crystals Having Fluorophenyl Group in the Core.
- [85] S. Merino, M. R. de la Fuente, Y. Gonzalez, M. A. Perez Jubindo, M. B. Ros, J. A. Puertolas; Phys. Rev., E 54, 5169-5177 (1996), Broadband dielectric measurements on the (R)-1-methylheptyl-6-(4'-decyloxybenzoyloxy)-2-naphthalene carboxylate antiferroelectric liquid crystal.
- [86] A. Fafara, B. Gestblom, S. Wrobel, R. Dabrowski, W. Drzewinski, D. Kilian, W. Haase; Ferroelectrics, 212, 79-90 (1998), Dielectric spectroscopy and electrooptic studies of new MHPOBC analogues.
- [87] M. Marzec, R. Dabrowski, A. Fafara, W. Haase, S. Hiller and S. Wrobel; Ferroelectrics, 180, 127-135 (1996), Gold-stone mode and Domain mode relaxation in ferroelectric phases of 4'-[*S,S*] - 2,3 - epoxyhexyloxy] Phenyl 4 (Decyloxy) Benzoate (EHPDB).
- [88] H. Uehara, Y. Hakanai, J. Hatano, S. Saito and K. Murashiro; Jpn. J. Appl. Phys., 34, 5424-5428 (1995), Dielectric Relaxation Modes in the Phases of Antiferroelectric Liquid Crystals.
- [89] M. Buivydas, S. T. Lagerwall, F. Gouda, R. Dubal, A. Takeichi; Ferroelectrics, 212, 55-65 (1998), The dependence of polarization and dielectric biaxiality on the enantiomeric excess in chiral dopant added to a smectic C host mixture.
- [90] H. Kresse, H. Schmalfuss, W. Weissflog, C. Tschierske, A. Hauser; Mol. Cryst. Liq. Cryst., 366, 505-517 (2001), Dielectric Characterization of Bn Phases.
- [91] W. Kuczynski, Electrooptical studies of relaxation processes in ferroelectric liquid crystals, in: W. Haase, S. Wrobel (Eds), Relaxation phenomena –Springer-Verlag, Berlin-Heidelberg, pp 422-444 (2003), Liquid crystals, magnetic systems, polymers, high-TC superconductors, metallic glasses.
- [92] A. M. Biradar, S. Wrobel and W. Haase; Phys. Rev. A, 39 2693-2702 (1989), Dielectric relaxation in the smectic-A\* and smectic-C\* phases of a ferroelectric liquid crystal.
- [93] J. K. Ahuja and K. K. Raina; Jpn. J. Appl. Phys., 39, 4076-4081 (2000), Polarization Switching and Dielectric Relaxations in Ferroelectric Liquid Crystals.
- [94] Y. P. Panarin, O. Kalinovskaya, J. K. Vij and J. W. Goodby; Phys. Rev. E, 55, 4345-4353 (1997), Observation and investigation of the ferrielectric subphase with high qT parameter.
- [95] I.-S. Baik, S. Y. Jeon, S. H. Lee, K. A. Park, S. H. Jeong, K. H. An, and Y. H. Lee; Appl. Phys. Lett. 87, 263110 (2005), Electrical-field effect on carbon nanotubes in a twisted nematic liquid crystal cell.

- [96] F. Haraguchi, K.-I Inoue, N. Toshima, S. Kobayashi, and K. Takatoh; *Jpn. J. Appl. Phys.* 46, L796 (2007), Reduction of the Threshold Voltages of Nematic Liquid Crystal Electrooptical Devices by Doping Inorganic Nanoparticles.
- [97] F. C. Frank; *Disc. Faraday Soc.*, 25, 19-28 (1958), On the theory of liquid crystals.
- [98] I. W. Stewart, *The Static and Dynamic Continuum Theory of Liquid Crystals*, Taylor and Francis, London (2004).
- [99] B. Kundu, S. K. Pal, S. Kumar, R. Pratibha, N.V. Madhusudana; *Phys. Rev. E: Stat. Nonlinear Soft Matter Phys.* 82, 061703 (2010), Splay and bend elastic constants in the nematic phase of some disulfide bridged dimeric compounds.
- [100] M. Hird; *Liq. Cryst.*, 38, 1467–1493 (2011), Invited topical review (Ferroelectricity in liquid crystals - materials, properties and applications).
- [101] C. B. Sawyer and C. H. Tower; *Phys. Rev.* 35, 269-273 (1930), Rochelle Salt as a Dielectric.
- [102] W. J. Merz; *Phy. Rev.* 95(3), 690-698 (1954), Domain Formation and Domain Wall Motions in Ferroelectric BaTiO<sub>3</sub> Single Crystals; Y. Ouchi, T. Uemura, H. Takezoe and A. Fukuda; *Jpn. J. Appl. Phys.*, 24 (4), L235- L238 (1985), Correspondence between Stroboscopic Micrographs and Spontaneous Polarization Measurements in Surface Stabilized Ferroelectric Liquid Crystal Cells.
- [103] W. J. Merz; *J. Appl. Phys.* 27(8), 938-943 (1956), Switching Time in Ferroelectric BaTiO<sub>3</sub> and Its Dependence on Crystal Thickness.
- [104] H. D. Megaw; *Acta Cryst.*, 7,187-194 (1954), Ferroelectricity and crystal structure II.
- [105] M. E. Lines and A. M. Glan; Oxford, Clarendon 128 (1977), *Principles and Applications of Ferroelectric and Related Materials*.
- [106] K. Miyasato, S. Abe, H. Takezoe, A. Fukuda and E. Kuze; *Jpn. J. Appl. Phys.* 22 (10), L661-L663 (1983), Direct Method with Triangular Waves for Measuring Spontaneous Polarization in Ferroelectric Liquid Crystals.
- [107] S. Kaur, A. K. Thakur, R. Chauhan, S. S. Bawa and A. M. Biradar; *Physica B*, 352, 337-341 (2004), The effect of rotational viscosity on the memory effect in ferroelectric liquid crystal.
- [108] S. S. Bawa, A. M. Biradar and S. Chandra; *Jpn. J. Appl. Phys.* 25(6), L446 (1986), Frequency Dependent Polarization Reversal and the Response Time of Ferroelectric Liquid Crystal by Triangular Wave Method.
- [109] J-Z. Zue, M. A. Handschy and N. A. Clark; *Ferroelectrics*, 73, 305-314 (1987), Electrooptic response during switching of a ferroelectric liquid crystal cell with uniform director orientation.

- [110] K. Skarp, I. Dahl, S. T. Lagerwall and B. Stebler; Mol. Cryst. Liq. Cryst., 114, 283-297 (1984), Polarization and Viscosity Measurements in a Ferroelectric Liquid Crystal by the Field Reversal Method.
- [111] T. Carlsson, B. Zeks, C. Filipic, A. Levstik; Physical Review A, 42, 877–889 (1990), Theoretical model of the frequency and temperature dependence of the complex dielectric constant of ferroelectric liquid crystals near the smectic C<sup>\*</sup> - smectic A phase transition.
- [112] J. D. Bernal and D. Crowfoot; Trans. Farad. Soc., 29, 1032-1049 (1933), Crystalline phases of some substances studied as liquid crystals.
- [113] P. Mandal and S. Paul; Mol. Cryst. Liq. Cryst., 131, 223-235 (1985), X-Ray Studies on the Mesogen 4'-n-Pentyloxy-4-Biphenylcarbonitrile (5OCB) in the Solid Crystalline State.
- [114] P. Mandal, S. Paul, H. Schenk and K. Goubitz; Mol. Cryst. Liq. Cryst., 135, 35-48 (1986), Crystal and Molecular Structure of the Nematogenic Compound 4-Cyanophenyl-4'-n-Heptylbenzoate (CPHB).
- [115] P. Mandal, B. Majumdar, S. Paul, H. Schenk and K. Goubitz; Mol. Cryst. Liq. Cryst., 168, 135-146 (1989), An X-ray Study of Cyanophenyl Pyrimidines, Part I - Crystal Structure of PCCPP.
- [116] P. Mandal, S. Paul, C. H. Stam and H. Schenk; Mol. Cryst. Liq. Cryst., 180, 369-378 (1990), X-Ray Studies of Cyanophenyl Pyrimidines Part II The Crystal and Molecular Structure of 5-(4-Ethylcyclohexyl)-2-(4-Cyanophenyl) Pyrimidine.
- [117] S. Gupta, P. Mandal, S. Paul, M. de Wit, K. Goubitz and H. Schenk; Mol. Cryst. Liq. Cryst., 195, 149-159 (1991), An X-Ray Study of Cyanophenylpyrimidines Part III. Crystal Structure of 5-(trans-4-Heptylcyclohexyl)-2-(4-Cyanophenyl) Pyrimidine.
- [118] P. Mandal, S. Paul, H. Schenk and K. Goubitz; Mol. Cryst. Liq. Cryst., 210, 21-30 (1992), Crystal and Molecular Structure of a Cybotactic Nematic Compound bis-(4'-n-Butoxybenzal)-2-Chloro-1,4-Phenylenediamine.
- [119] P. Mandal, S. Paul, K. Goubitz and H. Schenk; Mol. Cryst. Liq. Cryst., 258, 209-216 (1995), X-ray Structural Analysis of a Mesogenic Compound N,N'-Bis-(p Butoxybenzylidene)- $\alpha$ ,  $\alpha'$ - bi - p-Touidine.
- [120] A. Nath, S. Gupta, P. Mandal, S. Paul and H. Schenk; Liq. Cryst., 20, 765-770 (1996), Structural analysis by X-ray diffraction of a non-polar alkenyl liquid crystalline compound.
- [121] B. R. Jaishi, P. K. Mandal, K. Goubitz, H. Schenk, R. Dabrowski and K. Czuprynski; Liq. Cryst., 30, 1327-1333 (2003), The molecular and crystal structure of a polar mesogen 4-cyanobiphenyl-4'-hexylbiphenyl carboxylate.
- [122] P. A. C. Gane and A. J. Leadbetter; Mol. Cryst. Liq. Cryst., 78, 183-200 (1981), The Crystal and Molecular Structure of N- (4- n octyloxy benzylidene) -4'-butylaniline (80.4) and the Crystal-Smectic G Transition.

- [123] L. Malpezzi, S. Bruckner, D. R. Ferro and G. R. Luckhurst; *Liq. Cryst.*, 28, 357-363 (2001), Crystal structure and packing analysis of the liquid crystal dimer  $\alpha,\omega$ -bis(4-cyanobiphenyl-4'-yloxy)octane.
- [124] M. A. Sridhar, N. K. Lokanath, J. Shashidhara Prasad, C. V. Yelamaggad and Varshney, *Liq. Cryst.*, 28, 45-49 (2001), Crystal structure of a cholesterol-based dimesogen.
- [125] V. K. Gupta, P. Bhandhoria, M. Kalyan, M. Mathews and C. V. Yelamaggad; *Liq. Cryst.*, 32, 741-747 (2005), Crystal structure of a liquid crystal non-symmetric dimer: cholestryl 4-[4-(4-n-butylphenylethynyl)phenoxy]butanoate.
- [126] L. Walz, F. Nepveu and W. Haase; *Mol. Cryst. Liq. Cryst.*, 148, 111-121 (1987), Structural Arrangements of the Mesogenic Compounds 4-Ethyl-4'-(4"-pentylcyclohexyl) biphenyl and 4-Ethyl-2'-fluoro-4'-(4"-pentylcyclohexyl) biphenyl (BCH's) in the Crystalline State.
- [127] P. S. Patil, V. Shettigar, S. M. Dharmaprakash, S. Naveen, M. A. Sridhar and J. Shashidhara Prasad, *Mol. Cryst. Liq. Cryst.*, 461, 123-130 (2007), Synthesis and Crystal Structure of 1-(4-fluorophenyl)-3-(3,4,5-trimethoxyphenyl)-2-propen-1-one.
- [128] R. F. Bryan, Proceedings of the Pre-Congress Symposium on Organic Crystal Chemistry, Poznan, Poland, 105 (1979); *J. Structural Chem.*, 23, 128 (1982).
- [129] W. Haase and M. A. Athanassopoulou; *Liquid Crystals*, D.M.P. Mingos (Ed.), Springer-Verlag, Vol. I, pp. 139-197 (1999).
- [130] S. Biswas, S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk, and R. Dabrowski; *Cryst. Res. Technol.* 42, 10, 1029-1035 (2007), Crystal structure of a polar nematogen 4-(trans-4-undecylcyclohexyl) isothiocyanatobenzene.
- [131] S. Haldar, S. Biswas, P. K. Mandal, K. Goubitz, H. Schenk and W. Haase; *Mol. Cryst. Liq. Cryst.*, 490, 80-87 (2008), X-Ray Structural Analysis in the Crystalline Phase of a Nematogenic Fluoro-Phenyl Compound.
- [132] S. Haldar, P. K. Mandal, S. J. Prathap, T. N. Guru Row and W. Haase; *Liq. Crys.*, 35, 11, 1307-1312 (2008), X-ray studies of the crystalline and nematic phases of 49-(3,4,5-trifluorophenyl)-4-propylbicyclohexyl.
- [133] P. Das, A. N. Biswas, S. Acharya, A. Choudhury, P. Bandyopadhyay, P. K. Mandal and S. Upreti; *Mol. Cryst. Liq. Cryst.*, 501, 53-61 (2009), Structure of Liquid Crystalline 1-Phenyl-3-{4-[4-(4-octyloxybenzoyloxy)phenyloxycarbonyl]phenyl}triazene-1-oxide at Low Temperature.
- [134] B. K. Vainshtein, *Diffraction of X-rays by Chain Molecules*, Elsevier, Amsterdam, p12 (1966).
- [135] M. F. C. Ladd and R. A. Palmer, *Theory and practice of Direct methods in Crystallography*, Plenum Pub. Co., N. Y., (1980).

- [136] G. Germain, P. Main and M. M. Woolfson; *Acta Cryst.*, B26, 274-285 (1970), On the application of phase relationships to complex structures.
- [137] H. Schenk and C. T. Kiers, Simpel 83, an automatic direct method program package, University of Amsterdam (1983); H. Schenk; *Recl. Trav. Chim. Pays-bas*, 102, 1(1983).
- [138] G. M. Sheldrick, Direct method package program, University of Gottingen, Germany (1993).
- [139] S. R. Hall and J. M. Stewart, XTAL, University of Western Australia and Maryland (1989).
- [140] A. Altomere, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori and M. Camalli, The SIR 92 Programme (1992).
- [141] P. S. White, PC Version of NRCVAX (88), University of New Brunswick, Canada (1988).
- [142] Yao Jia-Xing, Zheng Chao-de, Quian Jin-Zi, Han Fu-Son, Gu Yuan-Zin and Fan Hai-Fu, SAPI: A Computer Program for Automatic Solution of Crystal Structures from X-ray Data, Institute of Physics, Academia Sinica, Beijing (1985).
- [143] C. J. Gilmore, *J. App. Cryst.*, 17, 42-46 (1984), MITHRIL-an integrated direct-methods computer program.
- [144] H. Hauptman and J. Karle, *Acta Cryst.*, 9, 45-55 (1956), Structure invariants and seminvariants for noncentrosymmetric space groups; *ibid.*, 12, 93-97 (1959); Seminvariants for centrosymmetric space group with conventional centered cells.
- [145] J. Karle and H. Hauptman, *Acta Cryst.*, 14, 217-223 (1961), Seminvariants for non-centrosymmetric space groups with conventional centered cells.
- [146] J. Karle and I. L. Karle, *Acta Cryst.*, 21, 849-859 (1966), The symbolic addition procedure for phase determination for centrosymmetric and non-centrosymmetric crystals.
- [147] H. Schenk (Ed), Direct Methods in Solving Crystal Structures, NATO ASI Series, Plenum Press, New York (1991).
- [148] G. H. Stout and L. H. Jensen, *X-ray Structure Determination*, Macmillan, New York p195 (1968).
- [149] A. J. C. Wilson, *Nature*, 150, 152 (1942), Determination of Absolute from Relative X-Ray Intensity Data.
- [150] W. Cochran and M. M. Woolfson, *Acta Cryst.*, 8, 1-12 (1955), "The theory of sign relations between structure factors".
- [151] W. Cochran, *Acta Cryst.*, 8, 473-478 (1955), Relations between the phases of structure factors.

- [152] J. Karle and H. Hauptman, *Acta Cryst.*, 9, 635-651 (1956), A theory of phase determination for the four types of non-centrosymmetric space groups.
- [153] M. J. Buerger, *Crystal Structure Analysis*, Wiley, New York (1960).
- [154] G. H. Stout and L. H. Jensen, *X-ray Structure Determination*, Macmillan, New York, p353 (1968).

## CHAPTER 3:

- [1] S. Gauza, H. Wang, C. H. Wen, S. T. Wu, A. Seed and R. Dabrowski; *Jpn. J. Appl. Phys.*, 42, 3463 (2003), High Birefringence Isothiocyanato Tolane Liquid Crystals.
- [2] J. S. Gasowska, S. J. Cowling, M. C. R. Cockett, M. Hird, R. A. Lewis, E. P. Raynes and J. W. Goodby; *J. Mater. Chem.*, 20, 299 (2010), The influence of an alkenyl terminal group on the mesomorphic behaviour and electro-optic properties of fluorinated terphenyl liquid crystals.
- [3] A. Mori, M. Hashimoto and S. Ujiie; *Liq. Cryst.*, 38, 263 (2011), Effects of a semi-fluorinated side chain and a lateral polar group on mesomorphic properties of 5-cyanotroponeoids and benzonitriles.
- [4] D. Demus, Y. Goto, S. Swada, E. Nakagawa, H. Saito and R. Tarao; *Mol. Cryst. Liq.Cryst.*, 260, 1 (1995), Trifluorinated Liquid Crystals for TFT Displays.
- [5] T. Nishi, A. Matsubara, H. Okada, H. Onnagawa, S. Sugimori, and K. Miyashita; *Jpn. J. Appl. Phys.*, 34, 236 (1995), Relationship between Molecular Structure and Temperature Dependence of Threshold Voltage in Fluorinated Liquid Crystals.
- [6] V. F. Petrov; *Liq. Cryst.*, 19, 729 (1995), Liquid crystals for AMLCD and TFT-PDLCD applications.
- [7] S. Gauza, X. Zhu, W. Piecek, R. Dabrowski, S. T. Wu; *J. Disp. Technol.* 3 (3), 250-252 (2007), Fast Switching Liquid Crystals for Color-Sequential LCDs.
- [8] M. Schadt, W. Helfrich; *Appl. Phys. Lett.*, 18, 127 (1971), Voltage-dependent optical activity of a twisted nematic liquid crystal.
- [9] K. Nishiyama, M. Okita, S. Kawaguchi, K. Teranishi, R. Takamatsu; *SID Tech. Dig.* 36, 132 (2005), 32" WXGA LCD TV using OCB Mode, Low Temperature p-Si TFT and Blinking Backlight Technology.
- [10] G. Harbers, C. Hoelen, *SID Tech. Dig.*; 32, 702 (2001), High Performance LCD Back lighting using High Intensity Red, Green and Blue Light Emitting Diodes.
- [11] D. P. Resler, D. S. Hobbs, R. C. Sharp, L. J. Friedman, T.A. Dorschner; *Opt. Lett.* 21 (9), 689 (1996), High-efficiency liquid-crystal optical phased-array beam steering.
- [12] J. Y. Hardeberg, F. Schmitt, H. Brettel; *Opt. Eng.* 41 (10), 2532 (2002), Multispectral color image capture using a liquid crystal tunable filter.
- [13] H. C. Lin, Y. H. Lin; *Opt. Express*, 20 (3), 2045 (2012), An electrically tunable-focusing liquid crystal lens with a low voltage and simple electrodes.

- [14] T.T. Ru, C.C. Yuan, R.P. Pan, C.L. Pan, X.C. Zhang; IEEE Microwave Wireless Compon. Lett. 14 (2) (2004), Electrically controlled room temperature terahertz phase shifter with liquid crystal.
- [15] R. L. Sutherland, V. P. Tondiglia, L. V. Natarajan; Appl. Phys. Lett. 64, 1074 (1994), Electrically switchable volume gratings in polymer-dispersed liquid crystals.
- [16] N. Mizoshita, K. Hanabusa, T. Kato; Adv. Funct. Mater. 13, 313 (2003), Fast and High-Contrast Electro-optical Switching of Liquid-Crystalline Physical Gels: Formation of Oriented Microphase-Separated Structures.
- [17] Y. H. Fan, H. W. Ren, S. T. Wu; Appl. Phys. Lett. 82, 2945 (2003), Normal-mode anisotropic liquid-crystal gels.
- [18] M. Hird; Chem. Soc. Rev., 2007; 36, 2070, Fluorinated liquid crystals – properties and applications.
- [19] J. A. Malecki and J. Nowak; J. Molecular Liquids, vol. 81, 245-252 (1999), Intermolecular interactions in benzene solutions of 4-heptyl-3'-cyano-biphenyl studied with non-linear dielectric effects.
- [20] S. Gauza, J. Li, S. T. Wu, A. Spadlo, R. Dabrowski, Y. N. Tzeng, K. L. Cheng, Liq. Cryst. 32(8), 1077 (2005), High birefringence and high resistivity isothiocyanate-based nematic liquid crystal mixtures.
- [21] S. Gauza, S. T. Wu, A. Spadlo, R. Dabrowski, J. Disp. Technol. 2(3), 247 (2006), High performance room temperature nematic liquid crystals based on laterally fluorinated isothiocyanato-tolanes.
- [22] D. Sinha, D. Goswami, P. K. Mandal, Ł. Szczucinski, R. Dabrowski; Mol Cryst Liq Cryst., 562, 156 (2012), On the nature of molecular associations, static permittivity and dielectric relaxation in a uniaxial nematic liquid crystal.
- [23] B. Jha, S. Paul, R. Paul, P. Mandal; Phase Transitions, 15, 39 (1989), Order parameters of some homologue cybotactic nematics from X-ray diffraction measurements.
- [24] A. de Vries; Mol Cryst Liq Cryst., 10, 31 (1970), Evidence for the existence of more than one type of nematic phase.
- [25] B. R. Jaishi, P. K. Mandal; Liq Cryst., 33, 753 (2006), Optical microscopy, DSC and X-ray diffraction studies in binary mixtures of 4-pentyloxy-49-cyanobiphenyl with three 4,49-di(alkoxy) azoxybenzenes.
- [26] P.G. de Gennes; Mol. Cryst. Liq. Cryst., 12, 193 (1971), Short range order effects in the isotropic phase of nematics and cholesterics.
- [27] H. E. J. Neugebauer; Canad. J. Phys., 32, 1-8 (1954), Clasius-Mossoti equation for certain types of anisotropic crystals.
- [28] Hyperchem 6.03, Hypercube Inc., Gainesville, FL, USA.
- [29] S. Biswas, S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk and R. Dabrowski, Crystal Research and Technology, Vol. 42, No.10, P. 1029 – 1035 (2007), Crystal structure of a polar nematogen 4-(trans-4-undecylcyclohexyl) isothiocyanatobenzene.
- [30] E. Megnassan and A. Proutier; Mol. Cryst. Liq. Cryst., 108, 245-254 (1984), Dipole Moments and Kerr Constants of 4-n-Alkyl-4'-Cyanobiphenyl Molecules (From 1CB to 12CB) Measured in Cyclohexane Solutions.
- [31] P. Sarkar, P. Mandal, S. Paul, R. Paul; Liq. Cryst, Vol. 30, No. 4, 507–527 (2003), X-ray diffraction, optical birefringence, dielectric and phase transition properties of the long homologous series of nematogens 4-(trans-4'-n-alkylcyclohexyl) isothiocyanatobenzenes.

- [32] A de Vries; Mol Cryst Liq Cryst., 10, 219 (1970), X-ray photographic studies of liquid crystals I. A cybotactic nematic phase.
- [33] P. Sarkar, P. Mandal, S. Paul, R. Dabrowski, K. Czuprynski; Liq.Cryst. 30 (4), 507 (2003), X-ray diffraction, optical birefringence, dielectric and phase transition properties of the long homologous series of nematogens 4-(trans -4'- n -alkylcyclohexyl) isothiocyanatobenzenes.
- [34] S. Haldar, S. Barman, P. K. Mandal, W. Haase, R. Dabrowski; Mol. Cryst. Liq.Cryst., 528, 81-95 (2010), Influence of Molecular Core Structure and Chain Length on the Physical Properties of Nematic Fluorobenzene Derivatives.
- [35] W. L. McMillan; Phys. Rev A, 4, 1238-1246 (1971), Simple molecular model for the smectic A phase of liquid crystals.
- [36] W. L. McMillan; Phys. Rev A, 6, 936 (1972), X-Ray Scattering from Liquid Crystals. I. Cholesteryl Nonanoate and Myristate.
- [37] P. Sarkar, P. Mandal, S. Paul, R. Dabrowski, K. Czuprynski; Liq Cryst. 30 (4), 507 (2003), X-ray diffraction, optical birefringence, dielectric and phase transition properties of the long homologous series of nematogens 4-(trans-4 $\infty$ -n-alkylcyclohexyl)isothiocyanatobenzenes.
- [38] A. J. Leadbetter, R. M. Richardson, C. N. Cooling; J Phys (Paris), 36 (C1), 37 (1975), The structure of a number of nematogens.
- [39] P. K. Sarkar, S. Paul, P. Mandal; Mol Cryst Liq Cryst., 265, 249 (1995), Small angle X-ray scattering studies on smectic and nematic phases of a toluidine compound.
- [40] N. V. Madhusudana, S. Chandrasekhar; Pramana Suppl., 1, 57 (1975), The role of permanent dipoles in nematic order.
- [41] S. Haldar, S. Biswas, P. K. Mandal, K. Goubitz, H. Schenk and W. Haase; Mol. Cryst. Liq. Cryst., 490, 80-87 (2008), X-Ray Structural Analysis in the Crystalline Phase of a Nematic Fluoro-Phenyl Compound.
- [42] S. Haldar, P. K. Mandal, S.J. Prathap, T. N. Guru Row and W. Haase; Liq. Cryst., 35, 1307-1312 (2008), X-ray studies of the crystalline and nematic phases of 49-(3,4,5-trifluorophenyl)-4- propylbicyclohexyl.
- [43] S. Biswas, S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk, R. Dabrowski; Crystal Res Technol., 42(10), 1029-1035 (2007), Crystal structure of a polar nematogen 4-(trans-4-undecylcyclohexyl) isothiocyanatobenzene.
- [44] H. Ishikawa, A. Toda, H. okada, H. Onnagawa, S. Sugimori; Liq. Cryst., 22, 743 (1997), Relationship between order parameter and physical constants in fluorinated liquid crystals.
- [45] S. Urban, P. Kula, A. Spadlo, M. Geppi, A. Marini; Liq.Cryst., 37, 1321 (2010), Dielectric properties of selected laterally fluoro-substituted 4,4"-dialkyl, dialkoxy and alkyl-alkoxy [1:1';4':1"] terphenyls.
- [46] P. Bordewijk, Physica, 75, 146 (1974), Extension of the Kirkwood-Frölich Theory of the Static dielectric Permittivity to anisotropic Liquids.
- [47] N. V. Madhsudana, B. S. Srikanta, R. Subramanya, M. Urs, Mol. Cryst. Liq. Cryst. 108, 19 (1984), Comparative X-ray and dielectric studies on some structurally related smectogenic compounds.

- [48] J. Schacht, P. Zugenmaier, M. Buivydas, L. Komitov, B. Stebler, S. T. Lagerwall, F. Gouda, F. Horii; *Phys Rev E.*, 61, 3926 (2000), Intermolecular and intramolecular reorientations in nonchiral smectic liquid-crystalline phases studied by broadband dielectric spectroscopy.
- [49] S. Urban, R. Dabrowski, J. Dziaduszek, J. Janik, J. K. Moscicki; *Liq. Cryst.* 26, 1817 (1999), Mesomorphic and dielectric properties of fluorosubstituted isothiocyanates with different bridging groups.
- [50] L. Benguigui; *Phys. Rev. A*, 28, 1852 (1983), Dielectric relaxation in the crystalline smectic-B phase.
- [51] M. Gu, Y. Yin, S. V. Shlyanovskii, O. D. Lavrentovich; *Phys Rev E.*, 76, 061702 (2007), Effects of dielectric relaxation on the director dynamics of uniaxial nematic liquid crystals.
- [52] H. Baessler, R. B. Beard, M. M. Labes; *J Chem Phys.*, 52, 2292 (1970), Dipole Relaxation in a Liquid Crystal.
- [53] J. Czub, R. Dabrowski, J. Dziaduszek, S. Urban; *Phase Transition*, 82, 485-495 (2009), Dielectric properties of ten three-ring LC fluorosubstituted isothiocyanates with different mesogenic cores.
- [54] A. C. Diogo, A. F. Martins; *J Phys (Paris)*, 43, 779 (1982), Order parameter and temperature dependence of the hydrodynamic viscosities of nematic liquid crystals.
- [55] S. Arrhenius; *Z Phys Chem.*, 4, 226 (1889), On the reaction rate of the inversion of non-refined sugar upon souring.
- [56] B. Kundu, S. K. Pal, S. Kumar, R. Pratibha, N. V. Madhusudana; *Physical Review E.*, 82, 061703 (2010), Splay and bend elastic constants in the nematic phase of some disulfide bridged dimeric compounds.
- [57] R. Dhar, A. S. Pandey, M. B. Pandey, S. Kumar and R. Dabrowski; *Applied Physics Express* 1, 121501 (2008), Optimization of the display parameters of a room temperature twisted nematic display material by doping single-wall carbon nanotubes.
- [58] M. D. Gupta, A. Mukhopadhyay, S. K. Roy, R. Dabrowski; *J Appl Phys.*, 113, 053516 (2013), High birefringence liquid crystal mixtures: Optical properties and order parameter.
- [59] E. Nowinowski-Kruszelnicki, J. Kedzierski, Z. Raszewski, L. Jaroszewicz, R. Dabrowski, M. Kojdecki, W. Piecek, P. Perkowski, K. Garbat, M. Oliferczuk, M. Sutkowski, Karolina Ogrodnik, P. Morawiak, E. Miszczyk; *Optica Applicata*, Vol. XLII, No. 1, oa120116 (2012), High birefringence liquid crystal mixtures for electro-optical devices.
- [60] H. S. Subramhanyam, C. S. Prabha and D. Krishnamurti; *Mol. Cryst. Liq. Cryst.*, 28, 201 (1974), Optical Anisotropy of Nematic Compounds.
- [61] M. Mitra, S. Paul and R. Paul; *Pramäna J. Phys.*, 29, 409 (1987), Optical Birefringence and Order Parameter of Three Nematogens.
- [62] N. V. Madhusudana, *Mol. Cryst. Liq. Cryst.*, 59, 117 (1980), Polarization Field and Orientational Order in Liquid Crystals.

## CHAPTER 4:

- [1] B. Bahadur, (1991). Liquid Crystals Applications and Uses, World Sci.: Singapore.
- [2] G. H. Heilmeier and L. A. Zanoni; *Appl. Phys. Lett.* 13, 91 (1968), Guest-host interactions in nematic liquid crystals: a new electro-optic effect.
- [3] S. Marino, M. Castriota, V. Bruno, E. Cazzanelli, G. Strangi, C. Versace and N. Scaramuzza; *J. Appl. Phys.*, 97, 013523 (2005), Changes of the electro-optic response of nematic liquid crystal cells due to inserted titania-vanadia films.
- [4] M. Oh-e and K. Kondo; *Appl. Phys. Lett.*, 69, 623 (1996), Response mechanism of nematic liquid crystals using the in-plane switching mode.
- [5] R. Muenster, M. Jarasch, X. Zhuang and Y. R. Shen; *Phys. Rev. Lett.*, 78, 42 (1997), Dye-Induced Enhancement of Optical Nonlinearity in Liquids and Liquid Crystals.
- [6] Y. Goto, T. Ogawa, S. Swada and S. Sugimori; *Mol. Cryst. Liq. Cryst.*, 209, 1–7 (1991), Fluorinated Liquid Crystals for Active Matrix Displays.
- [7] D. Demus, Y. Goto, S. Sawada, E. Nakagawa, H. Saito, R. Tarao; *Mol Cryst Liq Cryst.*, 260, 1–21(1995), Trifluorinated liquid crystals for TFT displays.
- [8] T. Nishi, A. Matsubara, H. Okada, H. Onnagawa, S. Sugimori, K. Miyashita; *Jpn J Appl Phys.*, 34, 236–239 (1995), Relationship between molecular structure and temperature dependence of threshold voltage in fluorinated liquid crystals.
- [9] V. F. Petrov; *Liq Cryst.*, 19, 729–741 (1995), Liquid crystals for AMLCD and TFTLCD applications; V. F. Petrov, Sofia I. Torgova ; Ludmila A. Karamysheva ; Shunsuke Takenaka; *Liq. Cryst.*, 26, 1141-1162 (1999), The trans-1,4-cyclohexylene group as a structural fragment in liquid crystals.
- [10] S. Gauza, H. Wang, C. H. Wen, S. T. Wu, A. Seed, R. Dabrowski; *Jpn J Appl Phys.*, 42, 3463–3466 (2003), High birefringence isothiocyanato tolane liquid crystals.
- [11] J. S. Gasowska, S. J. Cowling, M. C. R. Cockett, M. Hird, R. A. Lewis, E. P. Raynes, J. W. Goodby; *Mater Chem.*, 20, 299–307 (2010), The influence of an alkenyl terminal group on the mesomorphic behavior and electro-optic properties of fluorinated terphenyl liquid crystals.
- [12] H. Ishikawa, A. Toda, H. Okada, H. Onnagawa, S. Sugimori; *Liq Cryst.*, 22, 743–747 (1997), Relationship between order parameter and physical constants in fluorinated liquid crystals.
- [13] M. Hird; *Chem Soc Rev.*, 36, 2070–2095 (2007), Fluorinated liquid crystals – properties and applications.
- [14] M. Klasen, M. Bremer, A. Gotz, A. Manabe, S. Naemura, K. Tarumi; *Jpn J Appl Phys.*, 37, L945–L948 (1998), Calculation of optical and dielectric anisotropy of nematic liquid crystals.
- [15] K. Hori, M. Maeda, M. Yano, M. Kunugi; *Liq Cryst.*, 38, 287–293 (2011), The effect of fluorination (2): dependence of alkyl chain length on the crystal structures of mesogenic alkyl 4-[2-(perfluorohexyl) ethoxy] benzoates.

- [16] M. R. Cargill, G. Sandford, A. J. Tadeusiak, G. D. Love, N. Hollfelder, F. Pleis, G. Nelles, P. Kilickiran; *Liq Cryst.*, 38, 1069–1078 (2011), Highly fluorinated biphenyl ether systems as dopants for fastresponse liquid crystal display applications.
- [17] S. Haldar, K. C. Dey, D. Sinha, P. K. Mandal, W. Haase, P. Kula; *Liq Cryst.*, 39, 1196–1203 (2012), X-ray diffraction and dielectric spectroscopy studies on a partially fluorinated ferroelectric liquid crystal from the family of terphenyl esters.
- [18] E. Bartmann, D. Dorsch, U. Finkenzeller, H. A. Kurmeier and E. Poetsch; Freiburger Arbeitstagung Flüssigkristalle, 19, P8 (1990).
- [19] U. Finkenzeller, A. Kurmeier and E. Poetsch; Freiburger Arbeitstagung Flüssigkristalle, 18, P17 (1989).
- [20] P. Kirsch, S. Naemura and K. Tarumi; Freiburger Arbeitstagung Flüssigkristalle, 27, P45 (1998).
- [21] K. Tabayashi and K. Akasaka; *Liq. Cryst.*, 26, 127-129 (1999), Preliminary communication Natural abundance  $^2\text{H}$  NMR for liquid crystal studies: deuterium isotope effect on microscopic order.
- [22] H. Ishikawa, A. Toda, A. Matsubara, T. Nishi, H. Okada, H. Onnagawa, S. Sugimori and K. Miyashita; 20th Jpn. Symp. Liq. Cryst., Nagoya 1G505 (1994).
- [23] D. A. Dunmur, R. Hanson, H. Okada, H. Onnagawa, S. Sugimori and K. Toriyama; Asia Display' 95Proceeding, S22-2, 563 (1995), Effect of Intermolecular Interactions on Threshold Voltages for TN and TFT LC-Displays: Dipole Association in F-Substituted Cyclohexyl cyclohexyl benzenes.
- [24] S. Biswas, S. Haldar, P. K. Mandal, W. Haase; *Liq Cryst.*, 34, 365–372 (2007), X-ray diffraction and optical birefringence studies on four nematogenic difluorobenzene derivatives.
- [25] S. Haldar, S. Barman, P. K. Mandal, W. Haase, R. Dabrowski; *Mol Cryst Liq Cryst.*, 528, 81–95 (2010), Influence of molecular core structure and chain length on the physical properties of nematogenic fluorobenzene derivatives.
- [26] I. C. Khoo and S. T. Wu; *Optics and Nonlinear Optics of Liquid Crystals*, World Scientific: Singapore, (1993).
- [27] L. M. Blinov, V. G. Chigrinov; *Electro-Optic Effects in Liquid Crystal Materials*, Springer: New York, (1996).
- [28] M. Schadt; *Annu. Rev. Mater. Sci.*, 27, 305–379 (1997), Liquid crystal materials and liquid crystal displays.
- [29] B. Kundu, S. K. Pal, S. Kumar, R. Pratibha, N. V. Madhusudana; *Phys Rev E.*, 82, 061703-1–061703-9 (2010), Splay and bend elastic constants in the nematic phase of some disulfide bridged dimeric compounds.
- [30] H. Kresse; Dynamic dielectric properties of nematics. In: Dunmur, DA. Fukuda, A. Luckhurst, GR, editors. *Physical properties of liquid crystals: nematics*. London: INSPEC; 2001. p. 277–287.
- [31] H. Kresse; Dielectric properties of nematic liquid crystals. In: Demus, D. Goodby, J. Gray, GW. Spiess HW. Vill, V, editors. *Handbook of liquid crystals*. Vol. 2A. Weinheim, FRG: Wiley-VCH, Verlag GmbH, p. 91–112 (1998).
- [32] W. H. de Jeu, T. W. Lathouwers, D. Constants; *ZNaturforsch*, 29a, 905–911 (1974), Molecules structure of nematic liquid crystals. I. Terminally substituted azobenzene and azoxybenzenes.

- [33] J. Schacht, P. Zugenmaier, M. Buivydas, L. Komitov, B. Stebler, S. T. Lagerwall, F. Gouda, F. Horii; Phys Rev E., 61, 3926–3935 (2000), Intermolecular and intramolecular reorientations in nonchiral smectic liquid-crystalline phases studied by broadband dielectric spectroscopy.
- [34] D. A. Dunmur, W. H. Miller; Mol Cryst Liq Cryst., 60, 281–292 (1980), Dipole–Dipole correlation in nematic liquid crystals.
- [35] S. Haldar, D. Sinha, P. K. Mandal, K. Goubitz and R. Peschar; Liquid Crystals, Vol. 40, No. 5, 689–698 (2013), Effect of molecular conformation on the mesogenic properties of a partially fluorinated nematogenic compound investigated by X-ray diffraction and dielectric measurements.
- [36] S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk, W. Haase; Mol Cryst Liq Cryst., 490, 80-87 (2008), X-ray structural analysis in the crystalline phase of a nematogenic fluoro-phenyl compound.
- [37] S. Haldar, P. K. Mandal, S. J. Prathap, T. N. Guru Row, W. Haase; Liq Cryst., 35, 1307–1312 (2008), X-ray studies of the crystalline and nematic phases of 4<sub>-</sub>(3,4,5-trifluorophenyl)-4- propylbicyclohexyl.
- [38] W. Haase, M. A. Athanassopoulou; Vol. I. Berlin: Springer, p. 139–197 (1999), Crystal structures of LC mesogens. In: Mingos, M, editor. Structure and bonding.
- [39] M. Hird and K. J. Toyne; Mol. Cryst. Liq. Cryst., 323, 1-67 (1998), Fluoro Substitution in Thermotropic Liquid Crystals.
- [40] V. Vill; Liq. Cryst. Database, Version 4.4 , LCI Publisher, GmbH, Hamburg (2003) and references therein.
- [41] L. Bata and A. Buka; Mol. Cryst. Liq. Cryst, 63, 307-320 (1981), Dielectric Permittivity and Relaxation Phenomena in smectic Phases.
- [42] R. deGelder, R. A. G. de Graaff, H. Schenk; Acta Cryst., A49, 287–293 (1993), Automatic determination of crystal structures using Karle-Hauptman matrices.
- [43] D. T. Cromer, J. B. Mann; Acta Cryst., 24A, 321–324 (1968), X-ray scattering factors computed from numerical Hartree–Fock wave functions.
- [44] International Union of Crystallography. International tables for X-ray crystallography. Vol. IV, Birmingham: Kynoch Press, pp. 55 (1974).
- [45] D. T. Cromer, D. Liberman; J Chem Phys., 53, 1891–1898 (1970), Relativistic calculation of anomalous scattering factors for X-rays.
- [46] S. R. Hall, D. J. du Boulay, R. Olthof-Hazekamp, editors. XTAL3.7 system. Lamb: University of Western Australia (2000).
- [47] Spek AL. PLATON, An integrated tool for the analysis of the results of a single crystal structure determination. Acta Cryst., A46, C–34 (1990).
- [48] Hyperchem 6.03, Hypercube Inc., Gainesville, FL, USA.
- [49] P. Sarkar, P. K. Mandal, S. Paul, R. Paul; liq. Cryst, Vol. 30, No. 4, 507–527 (2003), X-ray diffraction, optical birefringence, dielectric and phase transition properties of the long homologous series of nematogens 4-(trans-4<sub>∞</sub>-n-alkylcyclohexyl ) isothiocyanatobenzenes.

- [50] E. Megnassan and A. Proutier; Mol. Cryst. liq. Cryst., 108, 245 (1984), Dipole Moments and Kerr Constants of 4-n Alkyl-4'- Cyanobiphenyl Molecules (From 1CB to 12CB) Measured in Cyclohexane Solutions.
- [51] K. P. Gueu, E. Megnassan and A. Proutier; Mol. Cryst. liq. Cryst., 132, 303 (1986), Dipole Moments of 4-n Alkyl-4'-Cyanobiphenyl Molecules (from OCB to 12CB) Measurement in Four Solvents and Theoretical Calculations.
- [52] W. Haase, H. Paulus, R. Pendzialek; Mol. Cryst. Liq. Cryst., 100: 211-221 (1983), Solid State Polymorphism in 4-Cyano-4'-n-Propyl biphenyl and X-Ray Structure Determination of the Higher Melting Modification.
- [53] B. R. Jaishi, P. K. Mandal, K. Goubitz, H. Schenk, R. Dabrowski, K. Czuprynski; Liq.Cryst., 30, 1327-1333 (2003), The molecular and crystal structure of a polar mesogen 4-cyanobiphenyl-4'-hexylbiphenyl carboxylate.
- [54] S. Biswas, S. Haldar, P. K. Mandal, K. Goubitz, H. Schenk, R. Dabrowski; Cryst. Res. Technol., 42, 1029-1035 (2007), Crystal structure of a polar nematogen 4-(trans-4- undecylcyclohexyl) isothiocyanatobenzene.
- [55] L. Walz, W. Haase, R. Eidenschink; Mol. Cryst. Liq. Cryst., 168, 169-182 (1989), The Crystal and Molecular Structures of Four Homologous, Mesogenic trans,trans-4,4'-dialkyl-(1 $\alpha$ ,1'-bicyclohexyl)-4 $\beta$ -carbonitril (CCN's).
- [56] S. Gupta, P. Mandal, S. Paul, K. Goubitz, M. de Wit, H. Schenk; Mol. Cryst. Liq.Cryst., 195, 149-159 (1991), An X-Ray Study of Cyanophenylpyrimidines Part III. Crystal Structure of 5-(trans-4-Heptylcyclohexyl)-2-(4-Cyanophenyl) Pyrimidine.
- [57] A. Nath, S. Gupta, P. Mandal, S. Paul, H. Schenk; Liq. Cryst., 20, 765-770 (1996), Structural analysis by X-ray diffraction of a non-polar alkenyl liquid crystalline compound.
- [58] P. Mandal, S. Paul, H. Schenk, K. Goubitz; Mol. Cryst. Liq. Cryst., 210: 21-30 (1992), Crystal and Molecular Structure of a Cybotactic Nematic Compound bis-(4'-n-Butoxybenzal)-2-Chloro-1,4-Phenylenediamine.
- [59] P. Mandal, S. Paul, H. Schenk, K. Goubitz; Mol. Cryst. Liq. Cryst., 135, 35-48 (1986), Crystal and Molecular Structure of the Nematogenic Compound 4-Cyanophenyl-4'-n-Heptylbenzoate (CPHB).
- [60] P. Mandal, S. Paul; Mol. Cryst. Liq. Cryst., 131, 223-235 (1985), X-Ray Studies on the Mesogen 4'-n-Pentyloxy-4-Biphenylcarbonitrile (5OCB) in the Solid Crystalline State.
- [61] L. Walz, F. Nepveu, W. Haase; Mol. Cryst. Liq. Cryst., 148, 111-121 (1987), Structural Arrangements of the Mesogenic Compounds 4-Ethyl-4'-(4"-pentylcyclohexyl)biphenyl and 4-Ethyl-2'-fluoro-4'-(4"-pentylcyclohexyl)biphenyl (BCH's) in the Crystalline State.
- [62] P. S. Patil, V. Shettigar, S. M. Dharmaprakash, S. Naveen, M. A. Sridhar, J. S. Prasad; Mol. Cryst. Liq. Cryst., 461, 123-130 (2007), Synthesis and Crystal Structure of 1-(4-fluorophenyl)-3-(3,4,5-trimethoxyphenyl)-2-propen-1-one.
- [63] S. Haldar; S. Barman; P. K. Mandal; W. Haase; R. Dabrowski; Mol. Cryst. Liq. Cryst., Vol. 528, 81–95 (2010), Influence of Molecular Core Structure and Chain Length on the Physical Properties of Nematogenic Fluorobenzene Derivatives.
- [64] A. J. Leadbetter, R. M. Richardson, C. N. Colling; 36, C1-37–C1-43 (1975), The structure of a number of nematogens. J Phys (Paris).
- [65] S. Gupta, G-P Chang-Chien, W-S Lee, R. Centore, S. P. Sen Gupta; Liq Cryst., 29, 657–661 (2002), Crystal structure of the mesogenic alkene monomer, 3-[4-(4-ethylbiphenyl)]-1-propene.

- [66] B. R. Ratna, R. Shashidhar; *Mol Cryst Liq Cryst.*, 42, 113–125 (1977), Dielectric studies on liquid crystals of strongly positive dielectric anisotropy.
- [67] B. R. Ratna, R. Shashidhar; *Mol Cryst Liq Cryst.*, 45, 103–116 (1978), Dielectric properties of some nematics of positive dielectric anisotropy.
- [68] M. Schadt; *J Chem Phys.*, 56, 1494–1497 (1972), Dielectric properties of some nematic liquid crystals with strong positive dielectric anisotropy.
- [69] N. V. Madhusudana, S. Chandrasekhar; *Pramana.*, 1, 57–68 (1975), The role of permanent dipoles in nematic order.
- [70] P. Bordewijk; *Physica.*, 75, 146 (1974), Extension of the Kirkwood-Fröhlich theory of the static dielectric permittivity to anisotropic liquids.
- [71] M. Gu, Y. Yin, S. V. Shlyanovskii, O. D. Lavrentovich; *Phys Rev E.*, 76, 061702 (2007), Effects of dielectric relaxation on the director dynamics of uniaxial nematic liquid crystals.
- [72] H. Baessler, R. B. Beard, M. M. Labes; *J Chem Phys.*, 52, 2292 (1970), Dipole Relaxation in a Liquid Crystal.
- [73] A. C. Diogo, A. F. Martins; *J Phys (Paris)*, 43, 779 (1982), Order parameter and temperature dependence of the hydrodynamic viscosities of nematic liquid crystals.
- [74] S. Arrhenius; *Z Phys Chem.*, 4, 226 (1889), On the reaction rate of the inversion of non-refined sugar upon souring.

## CHAPTER 5:

- [1] R. B. Meyer, Paper presented at the 5<sup>th</sup> Int. Liquid Crystal Conf., Stockholm, June, (1974); R. B. Meyer, L. Liebert, L. Strzelecki and P. Kelker, *J Phys (Paris) Lett.*, 36, L69 (1975), Ferroelectric liquid crystals.
- [2] A. D. L. Chandani, Y. Ouchi, H. Takezoe, A. Fukuda, K. Terashima, K. Furukawa and A. Kishi; *Jpn. J. Appl. Phys. Part 2*, 28, L1261 (1989), Novel Phases Exhibiting Tristable Switching.
- [3] Y. Izozaki, T. Fujikawa, H. Takezoe, A. Fukuda, T. Hagiwara, Y. Suzuki and I. Kawamura, *Phys. Rev. B*; 48, 13439 (1993), Devil's staircase formed by competing interactions stabilizing the ferroelectric smectic-C\* phase and the antiferroelectric smectic-C<sub>A</sub>\* phase in liquid crystalline binary mixtures.
- [4] J. P. F. Lagerwall, D. D. Parghi, D. Krugerke, F. Gouda, and P. Jagemalm, *Liq. Cryst.*, 29, 163 (2002), Phases, phase transitions and confinement effects in a series of antiferroelectric liquid crystals.
- [5] J. P. F. Lagerwall, P. Rudquist, and S.T. Lagerwall; *Liq. Cryst.*, 30, 399 (2003), On the phase sequence of antiferroelectric liquid crystals and its relation to orientational and translational order.
- [6] E. Gorecka, D. Pociecha, M. Čepič, B. Žekš, R. Dabrowski; *Phys. Rev. E*, 65, 061703 (2002), Enantiomeric excess dependence of the phase diagram of antiferroelectric liquid crystals.
- [7] I. C. Sage, Applications, in: D. Demus, J. Goodby, G.W. Gray, H.W. Spiess, V. Vill (Eds.), *Handbook of Liquid Crystals Vol. I – Fundamentals*, Wiley-Vch., Weinheim, pp. 731-762 (1998).

- [8] M. Hird; *Liq. Cryst.*, 38, 1467–1493 (2011), Ferroelectricity in liquid crystals-materials, properties and applications.
- [9] P. J. Collings, Ferroelectric liquid crystals: The 2004 Benjamin Franklin Medal in Physics presented to Robert B. Meyer of Brandeis University, *J. Frankl. Inst.* 342, 599–608 (2005).
- [10] J. Wen, M. Tian, Q. Chen, *Liq. Cryst.* 16, 445-453 (1994), Novel fluorinated liquid crystals. II. The synthesis and phase transitions of a novel type of ferroelectric liquid crystals containing 1,4-tetrafluorophenylene moiety.
- [11] Y. Xu, W. Wang, Q. Chen, J. Wen; *Liq. Cryst.*, 21, 65-71 (1996), Synthesis and transition temperatures of novel fluorinated chiral liquid crystals containing 1,4-tetrafluorophenylene units.
- [12] R. Dąbrowski, J. Gąsowska, J. Otón, W. Piecek, J. Przedmojski, M. Tykarska; High tilted antiferroelectric liquid crystalline materials, *Disp.* 25, 9-19 (2004).
- [13] R. Dąbrowski, P. Kula, Z. Raszewski, W. Piecek, J. M. Otòn, A. Spadło; *Ferroelectrics*, 395 116-132 (2010), New orthoconic antiferroelectrics useful for applications.
- [14] D. Ziobro, R. Dąbrowski, M. Tykarska, W. Drzewiński, M. Filipowicz, W. Rejmer, K. Kuśmirek, P. Morowiak, W. Piecek; *Liq. Cryst.* 39, 1011-1032 (2012), Synthesis and properties of new ferroelectric and antiferroelectric liquid crystals with a biphenyl benzoate rigid core.
- [15] H. T. Nguyen, J. C. Rouillon, A. Babeau, J. P. Marcerou, G. Sigaud; *Liq. Cryst.* 26, 1007-1019 (1999), Synthesis, properties and crystal structure of chiral semiperfluorinated liquid crystals with ferro and anticlinic smectic phases.
- [16] R. Dabrowski, *Ferroelectrics*, 243, 1-18 (2000), Liquid crystals with fluorinated terminal chains and antiferroelectric properties.
- [17] S. Seomun, T. Gouda, Y. Takanishi, K. Ishikawa, H. Takezoe; *Liq. Cryst.* 26, 151-161(1999), Bulk optical properties in binary mixtures of antiferroelectric liquid crystal compounds showing V-shaped switching.
- [18] J. V. Selinger, P. J. Collings, R. Shashidhar; *Phys. Rev. E* 64, 061705/1-9 (2001), Field-dependent tilt and birefringence of electroclinic liquid crystals: Theory and experiment.
- [19] W. M. Zoghaib, C. Carboni, A. K. George, S. AL-Manthari, A. Al-Hussaini, F. Al-Futaisi, , *Mol. Cryst. Liq. Cryst.* 542, 123–131 (2011), Novel fluorinated ferroelectric organosiloxane liquid crystals.
- [20] J. Naciri, C. Carboni, A. K. George; *Liq. Cryst.*, 30, 219-225 (2003), Low transition temperature organosiloxane liquid crystals displaying a de Vries smectic A phase.
- [21] J. Naciri, J. Ruth, G. Crawford, R. Shashidhar, B. R. Ratna; *Chem. Mater.*, 7, 1397-1402 (1995), Novel ferroelectric and electroclinic organosiloxane liquid crystals.
- [22] J. Dziaduszek, R. Dabrowski, K. Czuprynski, N. Bennis; *Ferroelectrics.*, 343, 3–9 (2006), Ferroelectric Compounds with Strong Polar Cyano Group in Terminal Position.
- [23] Z. Li, P. Salamon, A. Jakli, K. Wang, C. Qin, Q. Yang, C. Liu and J. Wen; *Liq. Cryst.*, 37 427–433 (2010), Synthesis and mesomorphic properties of resorcylic di[4-(4-alkoxy-2,3-diflorophenyl)ethynyl] benzoate liquid crystals.
- [24] Y. Yang, H. Li, J. Wen; *Liq. Cryst.*, 34, 975-979 (2007), Synthesis and mesomorphic properties of chiral fluorinated liquid crystals.

- [25] N. Shiratori, A. Yoshizawa, I. Nishiyama M. Fukumasa, A. Yokoyama, T. Hirai, M. Yamane; Mol. Cryst. Liq. Cryst. 199, 129-140 (1991), New Ferroelectric Liquid Crystals Having 2-Fluoro-2-Methyl Alkanoyloxy Group.
- [26] A. Fafara, B. Gestblom, S. Wróbel, R. Dabrowski, W. Drzewiński, D. Kilian, W. Haase; Ferroelectrics, 212, 79-90 (1998), Dielectric spectroscopy and electrooptic studies of new MHPOBC analogues.
- [27] D. M. Potukuchi, A. K. George, C. Carboni, S. H. Alharthi, J. Naciri; Ferroelectrics, 300, 79-93 (2004), Low Frequency Dielectric Relaxation, Spontaneous Polarization, Optical Tilt Angle and Response Time Investigations in a Fluorinated Ferroelectric Liquid Crystal, N125F2(R\*).
- [28] D. M. Potukuchi, A. K. George; Mol. Cryst. Liq. Cryst., 487, 92–109 (2008), Phase Transitions and Characterization in a Chiral Smectic-A<sub>de Vries</sub> Liquid Crystal by Low-Frequency Dielectric Spectroscopy.
- [29] P. K. Mandal, S. Haldar, A. Lapanik, W. Haase; Jpn. J. Appl. Phys., 48, 011501-6 (2009), Induction and enhancement of ferroelectric smectic C\* phase in multi-component room temperature mixtures.
- [30] K. Hiraoka, A. Taguchi, Y. Ouchi, H. Takezoe and A. Fukuda; Jpn. J. Appl. Phys., 29, 1473-1476 (1990), Electric-Field-Induced Transitions among Antiferroelectric, Ferrielectric and Ferroelectric Phases in a Chiral Smectic MHPOBC.
- [31] A. M. Biradar, S. Wróbel, and W. Haase; Phys. Rev. A, 39, 2693-2702 (1989), Dielectric relaxation in the smectic-A\* and smectic-C\* phases of a ferroelectric liquid crystal.
- [32] T. Carlsson, B. Zeks, C. Filipic, A. Levstik; Phys. Rev. A, 42, 877–889 (1990), Theoretical model of the frequency and temperature dependence of the complex dielectric constant of ferroelectric liquid crystals near the smectic-C\* - smectic-A phase transition.
- [33] B. Zeks, M. Cepic, in Relaxation Phenomena: Liquid Crystals, Magnetic Systems, Polymers, High-TC Superconductors, Metallic Glasses; W. Haase, S. Wrobel, Eds. Springer Verlag: Berlin–Heidelberg, p 477 (2003).
- [34] M. Buiydas, F. Gouda, S. T. Lagerwall, B. Stebler; Liq. Cryst., 18, 879–886 (1995), The molecular aspect of the double absorption peak in the dielectric spectrum of the antiferroelectric liquid crystal phase.
- [35] Hyperchem 6.03, Hypercube Inc., Gainesville, FL, USA.
- [36] S. Haldar, S. Barman, P. K. Mandal, W. Haase, R. Dabrowski; Mol. Cryst. Liq. Cryst., 528, 81-95 (2010), Influence of molecular core structure and chain length on the physical properties of nematogenic fluorobenzene derivatives.
- [37] P. Sarkar, P. Mandal, S. Paul, R. Paul, R. Dabrowski, K. Czuprynski; Liq. Cryst., 30, 507-527 (2003), X-ray diffraction, optical birefringence, dielectric and phase transition properties of the long homologous series of nematogens 4-(trans-4'-n-alkylcyclohexyl) isothiocyanobenzenes.
- [38] D. Sinha, D. Goswami, P. K. Mandal, Ł. Szczucinski, R. Dabrowski; Mol. Cryst. Liq. Cryst. 562, 156-165 (2012), On the nature of molecular associations, static permittivity and dielectric relaxation in a uniaxial nematic liquid crystal.
- [39] U. Manna, R. M. Richardson, A. Fukuda, J.K. Vij; Phys. Rev. E, 81, 050701–4 (2010), X-ray diffraction study of ferroelectric and antiferroelectric liquid crystal mixtures exhibiting de Vries SmA\*-SmC\* transitions.
- [40] S. T. Lagerwall, Ferroelectric liquid crystals in : D. Demus, J. Goodby, G.W. Gray, H. W. Spiess, V. Vill (Eds.), Handbook of Liquid Crystals Vol. 2B – Low molecular weight liquid crystals, Wiley-Vch., Weinheim, pp 515-664 (1998).

- [41] W. Piecek, Z. Raszewski, P. Perkowski, J. Przedmojski, J. Kedzierski, W. Drzewinski, R. Dabrowski, J. Zielinski; *Ferroelectrics.*, 310, 125–129 (2004), Apparent tilt angle and structural investigations of the fluorinated antiferroelectric liquid crystal material for display application.
- [42] J. P. F. Lagerwall, A. Saipa, F. Giesselmann, R. Dabrowski; *Liq. Cryst.* 31, 1175–1184 (2004), On the origin of high optical director tilt in a partially fluorinated orthoconic antiferroelectric liquid crystal mixture.
- [43] K. D'havé, P. Rudquist, S.T. Lagerwall, H. Pauwels, W. Drzewiski, R. Dabrowski; *Appl. Phys. Lett.*, 76, 3528–3530 (2000), Solution of the dark state problem in antiferroelectric liquid crystal displays.
- [44] W. K. Robinson, C. Carboni, P. Kloess, S. P. Perkins, H. J. Coles; *Liq. Cryst.*, 25, 301–307 (1998), Ferroelectric and antiferroelectric low molar mass organosiloxane liquid crystals.
- [45] E. P. Haridas, S. S. Bawa, A. M. Biradar, S. Chandra, *Jpn. J. Appl. Phys.*, 34, 3602–3606 (1995), Anisotropic surface anchoring conditions for gray-scale capability in high-tilt-angle ferroelectric liquid crystal.
- [46] R. Blinc, B. Zeks; *Phys. Rev. A* 18, 740–745 (1978), Dynamics of helicoidal ferroelectric smectic-C\* liquid crystals.
- [47] T. Carlsson, B. Zeks, C. Filipic, A. Levstik; *Physical Review A*, 42, 877–889 (1990), Theoretical model of the frequency and temperature dependence of the complex dielectric constant of ferroelectric liquid crystals near the smectic–smectic-A phase transition.
- [48] N. A. Clark, S. T. Lagerwall, Introduction to ferroelectric liquid crystals, in: J.W. Goodby, R. Blinc, N. A. Clark, S. T. Lagerwall, M.A. Osipov, S.A. Pikin, T. Sakurai, K. Yoshino, B. Zeks (Eds), *Ferroelectric Liquid Crystals Principles, Properties and Applications*, Gordon and Breach, Philadelphia, 1991, pp. 1–97.
- [49] D. Goswami, P. K. Mandal, R. Dabrowski; to be published.
- [50] S. Sato, J. Hatano, S. Tatemi, H. Uehara, S. Saito, E. Okabe; *Mol. Cryst. Liq. Cryst. A* 328, 411–418 (1999), Antiferroelectricity of a Chiral Smectic Liquid Crystal Having Three Isolated Phenyl Rings in the Core.
- [51] M. Marzec, W. Haase, E. Jakob, M. Pfeiffer, S. Wróbel; *Liquid Crystals*, 14, 1967–1976 (1993), The existence of four dielectric modes in the planar oriented SmC\* phase of a fluorinated substance.
- [52] S. Wróbel, G. Cohen, D. Davidov, W. Haase, M. Marzec, M. Pfeiffer; *Ferroelectrics*, 166, 211–222 (1995), Dielectric, electro optic and X-ray studies of a room temperature ferroelectric mixture.
- [53] J. M. Czerwic, R. Dabrowski, K. Garbat, M. Marzec, M. Tykarska, A. Wawrzyniak, S. Wróbel; *Liquid Crystals*, 39, 1503–1511 (2012), Dielectric and electro-optic behaviour of two chiral compounds and their antiferroelectric mixtures.
- [54] Y. P. Panarin, O. Kalinovskaya, J. K. Vij, and J. W. Goodby; *Phys. Rev. E*, 55, 4345–4353 (1997), Observation and investigation of the ferrielectric subphase with high qT parameter.
- [55] Y. P. Panarin, O. Kalinovskaya, and J. K. Vij; *Liq. Cryst.*, 25, 241–252 (1998), The investigation of the relaxation processes in antiferroelectric liquid crystals by broad band dielectric and electro-optic spectroscopy.
- [56] J. K. Song, U. Manna, A. Fukuda and J. K. Vij; *Appl. Phys. Lett.*, 93, 142903–3 (2008), Antiferroelectric dielectric relaxation processes and the interlayer interaction in antiferroelectric liquid crystals.
- [57] U. Manna, J. K. Song, G. Power, and J. K. Vij; *Phys. Rev. E*, 78, 021711–8 (2008), Effect of cell surfaces on the stability of chiral smectic-C phases.

- [58] U. Manna, J. K. Song, G. Power, J. K. Vij; Phys. Rev. E, 78, 021711-8 (2008), Effect of cell surfaces on the stability of chiral smectic-C phases.
- [59] M. Buivydas, F. Gouda, G. Andersson, S. T. Lagerwall, B. Stembler, J. Bomelburg, G. Heppke, B. Gestblom; Liq. Cryst., 23, 723- 739 (1997), Collective and non-collective excitations in antiferroelectric and ferrielectric liquid crystals studied by dielectric relaxation spectroscopy and electro-optic measurements.
- [60] S. Haldar, K. C. Dey, D. Sinha, P. K. Mandal, W. Haase, P. Kula; Liq. Cryst., 39, 1196-1203 (2012), X-ray diffraction and dielectric spectroscopy studies on a partially fluorinated ferroelectric liquid crystal from the family of terphenyl esters.
- [61] M. Marzec, A. Mikulko, S. Wrobel, R. Dabrowski, M. Darius and W. Haase; Liq. Cryst., 31, 153–159 (2004), Alpha sub-phase in a new ferroelectric fluorinated compound.
- [62] P. Nayek, S. Ghosh, S. Roy, T. P. Majumder, R. Dabrowski; J. Mol. Liq. 175, 91–96 (2012), Electro-optic and dielectric investigations of a perfluorinated compound showing orthoconic antiferroelectric liquid crystal.
- [63] P. Arora, A. Mikulko, F. Podgornov, W. Haase; Mol. Cryst. Liq. Cryst., 502, 1-8 (2009), Dielectric and electro-optic properties of new ferroelectric liquid crystalline mixture doped with carbon nanotubes.
- [64] A. Mikulko, P. Arora, A. Glushchenko, A. Lapanik and W. Haase; Euro Phys. Lett., 87, 27009/1-4 (2009), Complementary studies of BaTiO<sub>3</sub> nanoparticles suspended in a ferroelectric liquid-crystalline mixture.
- [65] P. K. Mandal, B. R. Jaishi, W. Haase, R. Dabrowski, M. Tykarska, P. Kula; Phase Transition, 79, 223–235 (2006), Optical microscopy, DSC and dielectric relaxation spectroscopy studies on a partially fluorinated ferroelectric liquid crystalline compound MHPO(13F)BC.
- [66] W. N. Thurmes, M. D. Wand, R. T. Vohra, K. M. More; SPIE Conf. Proc. 3015, 1-7 (1997), FLC materials for microdisplay applications.
- [67] M. Marzec, S. Wrobel, E. Gondek, R. Dabrowski; Mol. Cryst. Liq. Cryst., 410, 153–161 (2004), Room Temperature Antiferroelectric Phase Studied by Electrooptic Methods.
- [68] F. Gouda, T. Carlsson, G. Andersson, S.T. Lagerwall, B. Stebler; Liq. Cryst., 16, 315-322 (1994), Manifestation of biquadratic coupling in the smectic C\* phase soft mode dielectric response.
- [69] M. Marzec, S. Wröbel, S. Hiller, A. M. Biradar, R. Dabrowski, B. Gestblom, W. Haase; Mol. Cryst. Liq. Cryst., 302, 35-40 (1997), Dynamical properties of two ferroelectric phases of epoxy compound.
- [70] R. Eidenschink, T. Geelhaar, G. Andersson, A. Dahlgren, K. Flatischler, F. Gouda, S. T. Lagerwall, K. Skarp; Ferroelectrics, 84, 167-181 (1988), Parameter characteristics of a ferroelectric liquid crystal with polarization sign reversal.
- [71] A. K. Srivastavaa, R. Dhar, V. K. Agrawal, S. H. Lee, R. Dabrowski; Liq. Cryst., 35, 1101–1108 (2008), Switching and electrical properties of ferro- and antiferroelectric phases of MOPB(H)PBC.
- [72] S. Essid, M. Manai, A. Gharbi, J. P. Marcerou, J. C. Rouillon; Liq. Cryst., 32, 307–313, (2005), Electro-optical switching properties for measuring the parameters of a ferroelectric liquid crystal.

## CHAPTER 6:

- [1] D. Demus and J. Goodby, *Handbook of Liquid Crystals*, Wiley-VCH Weinheim, (1998).
- [2] P. S. Pershan, *Structure of Liquid Crystal Phases*, World Scientific Singapore, (1988).
- [3] R. B. Meyer, L. Liebert, L. Strzelecki, P. Keller; *J. Phys. (France) Lett.*, 36, L-69 (1975), Ferroelectric liquid crystals.
- [4] N. A. Clark and S. T. Lagerwall; *Appl. Phys. Lett.*, 36, 899 (1980), Submicrosecond bistable electro-optic switching in liquid crystals.
- [5] D. C. Wright and N. D. Mermin; *Rev. Mod. Phys.*, 66, 385 (1989), Crystalline liquids: the blue phases.
- [6] C. J. F. Botcher and P. Bordewijk; *Theory of Electric Polarization Vol. I* Elsevier, Amsterdam (1978).
- [7] Wrobel S, Dielectric relaxation spectroscopy, Relaxation phenomena — Liquid crystals, magnetic systems, polymers, high-TC superconductors, metallic glasses, Haase W. and Wrobel S Eds., Ch 1 Springer-Verlag Berlin-Heidelberg (2003).
- [8] D. Goswami, D. Sinha, A. Debnath, P.K. Mandal, S. K. Gupta, W. Haase, D. Ziobro and R. Dabrowski; *J. Mol. Liq.*, 182, 95-101 (2013), Molecular and dynamical properties of a perfluorinated liquid crystal with direct transition from ferroelectric SmC\* phase to isotropic phase.
- [9] K. Miyasato, S. Abe, H. Takezoe, A. Fukuda and E. Kuze; *Jpn. J. Appl. Phys.*, 22, L661 (1983), Direct Method with Triangular Waves for Measuring Spontaneous Polarization in Ferroelectric Liquid Crystals.
- [10] P. K. Mandal, B. R. Jaishi, W. Haase, R. Dabrowski, M. Tykarska and P. Kula; *Phase Trans.*, 79, 223-235 (2006), Optical microscopy, DSC and dielectric relaxation spectroscopy studies on a partially fluorinated ferroelectric liquid crystalline compound MHPO(13F)BC.
- [11] I. Dierking, *Textures of Liquid Crystals* Wiley-VCH, Weinheim, (2003).
- [12] A. J. Leadbetter, J. P. Gaughan, B. Kelly, G. W. Gray and J. Goodby; *J Phys. (France) Colloq.*, 40, C3-178 (1979), Characterisation and structure of some new smectic phases.
- [13] G. W. Gray and J. W. Goodby *Smectic Liquid Crystals, Textures and Structures*, Leonard Hill, Philadelphia, (1984).
- [14] P. A. C. Gane, A. J. Leadbetter and P. G. Wrighton; *Mol. Cryst. Liq. Cryst.*, 66, 247 (1981), Structure and Correlations in Smectic B, F and I Phases.
- [15] H. Stegmeyer, T. Blumel, K. Hiltrop, H. Onnusseit and F. Porsch; *Liq. Cryst. 1*, 3-28 (1986), Thermodynamic, structural and morphological studies on liquid-crystalline blue phases.
- [16] Jan P. F. Lagerwall, Per Rudquist, Sven T. Lagerwall and Frank Gießelmann; *Liq. Cryst.*, 30 399-414 (2003), On the phase sequence of antiferroelectric liquid crystals and its relation to orientational and translational order.
- [17] S. B. Dierker and R. Pindak; *Phys. Rev. Lett.*, 59, 1002 (1987), Dynamics of thin tilted hexatic liquid crystal films.
- [18] E. B. Sirota, P. S. Pershan, L. B. Sorenson and J. Collett; *Phys. Rev. Lett.*, 55, 2039 (1985), X-Ray Studies of Tilted Hexatic Phases in Thin Liquid-Crystal Films.

- [19] E. Grelet, B. Pansu, M-H Li and H. T. Nguyen; Phys. Rev. Lett., 86, 3791 (2001), Structural Investigations on Smectic Blue Phases.
- [20] T. Uemura, Y. Ouchi, K. Ishikawa, H. Takezoe and A. Fukuda; Jpn. J Appl. Phys., 24, L224 (1985), Optical Microscope Observation of Hexagonal Ordering in Surface Stabilized Ferroelectric Liquid Crystal Cells.
- [21] D. Ziobro, R. Dąbrowski, M. Tykarska, W. Drzewiński, M. Filipowicz, W. Rejmer, K. Kuśmirek, P. Morowiak and W. Piecik; Liq. Cryst., 39, 1011 (2012), Synthesis and properties of new ferroelectric and antiferroelectric liquid crystals with a biphenyl benzoate rigid core.
- [22] W. Piecik, Z. Raszewski, P. Perkowski, J. Przedmojski, J. Kedzierski, W. Drzewinski, R. Dabrowski and J. Zielinski; Ferroelectrics, 310, 125 (2004), Apparent Tilt Angle and Structural Investigations of the Fluorinated Antiferroelectric Liquid Crystal Material for Display Application.
- [23] J. P. F. Lagerwall, A. Saipa, F. Giesselmann and R. Dabrowski; Liq. Cryst., 31, 1175 (2004), On the origin of high optical director tilt in a partially fluorinated orthoconic antiferroelectric liquid crystal mixture.
- [24] A. Mikułko, M. Wierzejska, M. Marzec, S. Wrobel, J. Przedmojski and W. Haase; Mol. Cryst. Liq. Cryst., 477, 185 (2007), Ferroelectricity of Hexatic Phases.
- [25] T. Carlsson, B. Zeks, C. Filipic, A. Levstik; Phys. Rev. A, 42, 877 (1990), Theoretical model of the frequency and temperature dependence of the complex dielectric constant of ferroelectric liquid crystals near the smectic-C\*-smectic-A phase transition.
- [26] M. Glogarova and I. Rychetsky; Dielectric relaxation spectroscopy, Relaxation phenomena — Liquid crystals, magnetic systems, polymers, high-TC superconductors, metallic glasses, W. Haase and S. Wrobel, Eds., Ch 5.3 Springer-Verlag Berlin-Heidelberg (2003).
- [27] I. Rychestsky, D. Pociecha , V. Dvorak, J. Mieczkowski, E. Gorecka and M. Glogarova; J. Chem. Phys., 111, 1541 (1999), Dielectric behavior of ferroelectric liquid crystals in the vicinity of the transition into the hexatic phase.