

# LIST OF TABLES

<u>CHAPTERS</u>	<u>TABLE CAPTIONS</u>	<u>PAGE NO.</u>
Chapter-II	Table II.1. Comparison between organic solvents and ionic liquids.	46
	Table II.2. Properties of Ionic Liquids	49-50
Chapter-IV	Table IV.1: Density ( $\rho$ ), viscosity ( $\eta$ ) and relative permittivity ( $\epsilon$ ) of the different solvents dimethyl acetamide (DMA), tetrahydrofuran (THF) and methyl cellosolve (MC) at different temperatures.	224
	Table IV.2: The concentration ( $c$ ) and molar conductance ( $\Lambda$ ) of [bmmim][BF <sub>4</sub> ] in Methyl Cellosolve, Dimethyl acetamide and Tetrahydrofuran at 298.15, 303.15, 308.15 K respectively.	225-227
	Table IV.3: Limiting molar conductance ( $\Lambda_0$ ), association constant ( $K_A$ ), co-sphere diameter ( $R$ ) and standard deviations of experimental $\Lambda$ ( $\delta$ ) obtained from Fuoss conductance equation for 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in Methyl cellosolve and DMA at 298.15, 303.15, 308.15 K respectively.	228
	Table IV.4: walden product ( $\Lambda_0 \cdot \eta$ ) and Gibb's energy change ( $\Delta G^\ddagger$ ) of 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in Methyl cellosolve and DMA at 298.15 K, 303.15 K and 308.15 K respectively are given below.	229
	Table IV.5: Limiting Ionic Conductance ( $\lambda_0^\ddagger$ ), Ionic Walden Product ( $\lambda_0^\ddagger \eta$ ), Stokes' Radii ( $r_s$ ), and Crystallographic Radii ( $r_c$ ) of 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in Methyl cellosolve and DMA at 298.15 K, 303.15 K and 308.15 K respectively are given below.	230-231
	Table IV.6: The calculated limiting molar conductance of ion-pair ( $\Lambda_0$ ), limiting molar conductances of triple ion $\Lambda_0^T$ , experimental slope and intercept obtained from Fuoss-Kraus Equation for 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in THF at 298.15 K, 303.15 K and 308.15 K respectively.	231

## List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	<i>Table IV.7: Salt concentration at the minimum conductivity (<math>C_{min}</math>) along with the ion-pair formation constant (<math>K_P</math>), triple ion formation constant (<math>K_T</math>) for 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in THF at 298.15 K.</i>	232
	<i>Table IV.8: Salt concentration at the minimum conductivity (<math>C_{min}</math>), the ion pair fraction (<math>\alpha</math>), triple ion fraction (<math>\alpha_T</math>), ion pair concentration (<math>C_P</math>) and triple-ion concentration (<math>C_T</math>) for 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in THF at 298.15 K, 303.15 K and 308.15 K respectively.</i>	233
	<i>Table IV.9: Concentration, <math>c</math>, density, <math>\rho</math>, apparent molar volume, <math>\phi_v</math>, limiting apparent molar volume <math>\phi_v^0</math> and experimental slope for 1-butyl-2,3-dimethylimidazolium tetrafluoroborate in Methyl Cellosolve, Dimethyl Acetamide and Tetrahydrofuran at 298.15 K, 303.15 K and 308.15 K respectively.</i>	234-237
	<i>Table IV.10: Values of empirical coefficients (<math>a_0</math>, <math>a_1</math>, and <math>a_2</math>) of Equation 4 for IL in different solvents (MC, DMA, THF) at 298.15K to 308.15K respectively</i>	237
	<i>Table IV.11: Limiting apparent molal expansibilities (<math>\phi_E^0</math>) for IL in different solvents (THF, DMA, THF) at 298.15K to 308.15K respectively</i>	238
	<i>Table IV.12: Concentration, <math>c</math>, viscosity, <math>\eta</math>, <math>\frac{(\eta_r - 1)}{\sqrt{c}}</math>, viscosity A and B coefficients for 1-butyl-2,3-dimethylimidazolium iodide in Methyl Cellosolve, Dimethyl Formamide and Tetrahydrofuran at 298.15 K, 303.15 K and 308.15 K respectively.</i>	239-240
	<i>Table IV.13: Stretching frequencies of the functional groups present in the pure solvent and change of frequency after addition of 0.05(M) concentration of [bmmim][BF<sub>4</sub>] in THF, DMA and MC.</i>	240
Chapter-V	<i>Table V.1: Surface tension (<math>\gamma</math>) with corresponding concentration, and concentration ratio (ratio of inclusion IL:<math>\beta</math>-CD) at the first and second break point of 4-methyl-N-butylpyridinium chloride solution (0.001 M) in aqueous <math>\beta</math>-CD</i>	252

List of Tables

<u>CHAPTERS</u>	<u>TABLE CAPTIONS</u>	<u>PAGE NO.</u>
	<i>Table V.2: Conductance (<math>\Lambda</math>) with corresponding concentration, and concentration ratio (ratio of inclusion IL:<math>\beta</math>-CD) at the first and second break point of 4-methyl-N-butylpyridinium chloride solution (0.001 M) in aqueous <math>\beta</math>-CD</i>	252
	<i>Table V.3: Solvation number (<math>S_n</math>), ionic apparent molar volume (<math>\phi_{\pm}^{\circ}</math>) in deferent mass fractions of aqueous <math>\beta</math>-cyclodextrin mixtures</i>	253
	<i>Table V.S.1: Experimental values of density (<math>\rho</math>), viscosity (<math>\eta</math>), and refractive index (<math>n_D</math>) in deferent mass fraction of aqueous <math>\beta</math>-cyclodextrin mixtures</i>	260
	<i>Table V.S.2: Experimental values of densities (<math>\rho</math>) and viscosities (<math>\eta</math>) corresponding to concentration in different mass fractions of aq. <math>\beta</math>-cyclodextrin at different temperature</i>	261-262
	<i>Table V.S.3: Limiting apparent molar volume (<math>\phi_{\nu}^{\circ}</math>), experimental slope (<math>S_{\nu}^*</math>), viscosity B and A-coefficient, Solvation number (<math>S_n</math>), ionic apparent molar volume (<math>\phi_{\pm}^{\circ}</math>) in deferent mass fractions of aqueous <math>\alpha</math> and <math>\beta</math>-cyclodextrin mixtures</i>	263
	<i>Table V.S.4: Refractive index (<math>n_D</math>), molar refraction (<math>R_m</math>), and limiting molar refraction (<math>R_m^0</math>) for [(bmPy)Cl] in different mass fraction of aqueous <math>\beta</math>-CD at 298.15 K</i>	264-265
Chapter VI	<i>Table VI.1: Sample description</i>	284
	<i>Table VI.2: Values of density (<math>\rho</math>), viscosity (<math>\eta</math>) and relative permittivity (<math>\epsilon_r</math>) at <math>T = 298.15</math> K and pressure <math>p = 101.325</math> kPa of studied pure solvents</i>	284
	<i>Table VI.3: Molar conductance (<math>\Lambda</math>) and the corresponding concentration (<math>m</math>) of [Bu<sub>4</sub>PMS] at <math>T = 298.15</math> K and pressure <math>p = 101.325</math> kPa in different studied solvents</i>	285-286
	<i>Table VI.4: Limiting molar conductivity (<math>\Lambda_0</math>), the association constant (<math>K_A</math>), the distance of closest approach of ions (<math>R</math>), standard deviations <math>\delta</math> of experimental <math>\Lambda</math> from equation 1, Walden product (<math>\Lambda_0\eta</math>) and Gibb's energy change (<math>\Delta G^{\circ}</math>) of [Bu<sub>4</sub>PMS] in different studied solvents at <math>T = 298.15</math> K</i>	287

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	<i>Table VI.5: Limiting ionic conductance (<math>\lambda_o^\pm</math>), ionic Walden product (<math>\lambda_o^\pm \eta</math>), Stokes' radii (<math>r_s</math>) and crystallographic radii (<math>r_c</math>) of [Bu<sub>4</sub>PMS] in different studied solvents at T = 298.15 K</i>	287
	<i>Table VI.6: Diffusion coefficient (<math>D_\pm</math>) and ionic mobility (<math>i_\pm</math>) of Bu<sub>4</sub>P<sup>+</sup> and MS in different studied solvents at 298.15K</i>	288
	<i>Table VI.7: Stretching frequencies of the functional groups present in the solvents and change of frequency after addition of IL, [Bu<sub>4</sub>PMS] in the solvents.</i>	289
Chapter VII	<i>Table VII.1: The values of Density (<math>\rho</math>), Viscosity (<math>\eta</math>), Refractive index (<math>n_D</math>), and Speed of sound (<math>u</math>) in different mass fraction of 1-butylpyridinium bromide solution at 298.15K</i>	305
	<i>Table VII.2: Experimental values of Densities (<math>\rho</math>), Viscosities (<math>\eta</math>), Refractive Index (<math>n_D</math>) and Ultrasonic Speed (<math>u</math>) of L-Glycine, L-Alanine and L-Valine in different mass fraction of 1-butylpyridinium bromide at 298.15K</i>	306-308
	<i>Table VII.3: Molality, apparent molar volume (<math>\phi_V</math>), <math>(\eta/\eta_0 - 1)/m^{1/2}</math>, molar refraction (<math>R</math>), adiabatic compressibility (<math>\beta</math>) and apparent molal adiabatic compressibility (<math>\phi_K</math>) of L-Glycine, L-Alanine, and L-Valine in 1-butylpyridinium bromide at 298.15 K</i>	308-311
	<i>Table VII.4: Limiting apparent molar volumes (<math>\phi_V^0</math>), experimental slopes (<math>S_V^*</math>), A, B coefficients, limiting partial adiabatic compressibility (<math>\phi_K^0</math>), and experimental slope (<math>S_K^*</math>) of L-Glycine, L-Alanine, and L-Valine in aqueous 1-butylpyridinium bromide at 298.15 K</i>	311
Chapter VIII	<i>Table VIII.1: Values of surface tension at the break point (<math>\gamma</math>) with corresponding concentration of IL in different mass fraction of aqueous <math>\beta</math>-cyclodextrin at 298.15K</i>	324

List of Tables

<u>CHAPTERS</u>	<u>TABLE CAPTIONS</u>	<u>PAGE NO.</u>
	<i>Table VIII.2: Values of Specific conductance at the break point (<math>\kappa</math>) with corresponding concentration of IL in different mass fraction of aqueous <math>\beta</math>-cyclodextrin at 298.15K<sup>a</sup></i>	324
	<i>Table VIII.3: Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin host molecules when complexed with amino acid guest molecules in D<sub>2</sub>O at 298.15Ka</i>	325
Chapter IX	<i>Table IX.1: Density (<math>\rho</math>), viscosity (<math>\eta</math>), refractive index (<math>n_D</math>) and relative permittivity (<math>\epsilon</math>) of the different solvents Tetrahydrofuran, 1,4 Dioxane and Acetonitrile.</i>	347
	<i>Table IX.2: The concentration (<math>c</math>) and molar conductance (<math>\Lambda</math>) of [bupy]Br in 1,4 Dioxane, Tetrahydrofuran and Acetonitrile at 298.15 K.</i>	348
	<i>Table IX.3: Limiting molar conductance (<math>\Lambda_0</math>), association constant (<math>K_A</math>), co-sphere diameter (<math>R</math>) and standard deviations of experimental <math>\Lambda</math> (<math>\delta</math>) obtained from Fuoss conductance equation for 1-butyl-pyridinium bromide in Acetonitrile at 298.15 K.</i>	349
	<i>Table IX.4: Walden product (<math>\Lambda_0 \cdot \eta</math>) and Gibb's energy change (<math>\Delta G^\ddagger</math>) of 1-butyl-pyridinium bromide in Acetonitrile at 298.15 K.</i>	349
	<i>Table IX.5: Limiting Ionic Conductance (<math>\lambda_0^\pm</math>), Ionic Walden Product (<math>\lambda_0^\pm \eta</math>), Stokes' Radii (<math>r_s</math>), and Crystallographic Radii (<math>r_c</math>) of [bupy]Br in ACN at 298.15 K.</i>	349
	<i>Table IX.6: The calculated limiting molar conductance of ion-pair (<math>\Lambda_0</math>), limiting molar conductances of triple ion <math>\Lambda_0^T</math>, experimental slope and intercept obtained from Fuoss-Kraus Equation for 1-butyl-puridinium bromide in 1,4 DO and THF at 298.15 K.</i>	350
	<i>Table IX.7: Salt concentration at the minimum conductivity (<math>C_{min}</math>) along with the ion-pair formation constant (<math>K_P</math>), triple ion formation constant (<math>K_T</math>) for 1-butyl-puridinium bromide in 1,4 DO and THF at 298.15 K.</i>	350
	<i>Table IX.8: Salt concentration at the minimum conductivity (<math>C_{min}</math>), the ion pair fraction (<math>\alpha</math>), triple ion fraction (<math>\alpha_T</math>), ion pair concentration (<math>c_P</math>) and triple-ion concentration (<math>c_T</math>) for 1-butyl-puridinium bromide in 1,4 DO and THF at 298.15 K.</i>	351
	<i>Table IX.9: Density (<math>\rho</math>), viscosity (<math>\eta</math>) and refractive index (<math>n_D</math>) of 1-butyl-pyridinium bromide in different mass fraction of 1,4 DO, Tetrahydrofuran, and Acetonitrile at</i>	351-354

List of Tables

<u>CHAPTERS</u>	<u>TABLE CAPTIONS</u>	<u>PAGE NO.</u>
	<i>different temperatures.</i>	
	<i>Table IX.10: Apparent molar volume (<math>\phi_v</math>), <math>\frac{(\eta_r - 1)}{\sqrt{c}}</math> and molar refraction (<math>R_m</math>) for 1-butyl-pyridinium bromide in different mass fraction of 1,4 DO, Tetrahydrofuran, and Acetonitrile at different temperatures.</i>	354-357
	<i>Table IX.11: Limiting apparent molar volume (<math>\phi_v^0</math>), experimental slope, viscosity -A and -B coefficient and limiting molar refraction (<math>R_M^0</math>) for 1-butyl-pyridinium bromide in 1,4 DO, Tetrahydrofuran, and Acetonitrile at different temperatures.</i>	357-358
	<i>Table IX.12: Values of empirical coefficients (<math>a_0</math>, <math>a_1</math>, and <math>a_2</math>) of Equation 4 for IL in different solvents (1,4 DO, THF and ACN) at 298.15K to 308.15K respectively</i>	358
	<i>Table IX.13: Limiting apparent molal expansibilities (<math>\phi_E^0</math>) for IL in different solvents (1,4 DO, THF and ACN) at 298.15K to 308.15K respectively</i>	359
	<i>Table IX.14: Stretching frequencies of the functional groups present in the pure solvents and {solvents+[bupy][Br]}</i>	359