

LIST OF TABLES

Table 1: Table showing the GIS locations of the sampling sites

Table 2: Identification of the plant samples along with the accession no. of the deposited samples

Table 3: Systematic position of the described plant samples

Table 4: Phytochemicals screened in the powdered plant samples

Table 5: Quantification of total phenol and protein content in the crude powdered plant samples

Table 5A and B: ANOVA of data present in Table 5

Table 6: Quantification of total chlorophyll, chlorophyll- a and chlorophyll-b in the powdered plant samples

Table 6A, B and C: ANOVA of data present in Table 6

Table 7: Total lipid and carotenoid content in the crude powdered plant samples

Table 7A and B: ANOVA of data present in Table 7

Table 8: Vitamin C and E content in the dried powdered plant samples

Table 8A & 8B: ANOVA of data presented in Table 8

Table 9: IC₅₀ values of DPPH radical scavenging activity exhibited by the extracts (ME, HWE, EE) of different samples at different concentrations.

Table 9A: ANOVA of data present in Table 9

Table 10: IC₅₀ values of hydrogen peroxide scavenging activity exhibited by the extracts of different samples at different concentrations.

Table 10A: ANOVA of data present in Table 10

Table 11: IC₅₀ values of nitric oxide radical scavenging activity exhibited by the extracts of different samples at different concentrations.

Table 11A: ANOVA of data presented in Table 11

Table 12: IC₅₀ values of superoxide radical scavenging activity exhibited by the extracts of different samples at different concentrations.

Table 12A: ANOVA of data presented in Table 12

Table 13: Ferric reducing antioxidant power (FRAP) assay of the extracts (ME, HWE, EE) of different plants at different concentrations

Table 13A: ANOVA of data presented in Table 13

Table 14: Peak value of standard phenolics

Table 15: Phenolics identified through HPLC in the methanol extracts of different plant samples.

Table 16: Peak table for the data presented in Figure (30A) of *N. cordifolia*

Table 17: Peak table for the data presented in Figure (30B) of *C. dentatus*

Table 18: Peak table for the data presented in Figure (31A) of *D. quercifolia*

Table 19: Peak table for the data presented in Figure (31B) of *P. cuspidatus*

Table 20: Peak table for the data presented in Figure (32A) of *D. linearis*

Table 21: Peak table for the data presented in Figure (32B) of *P. biaurita*

Table 22: Peak table for the data presented in Figure (33A) of *P. vittata*

Table 23: Peak table for the data presented in Figure (33B) of *M. punctatum*

Table 24: Phenolic compounds quantified in the plant samples through HPLC

Table 25: Antimicrobial activity of plant extract against four pathogenic bacterial strains

Table 26: Spore germination bioassay of crude plant extracts and controls against different fungal species

Table 27: Radial growth bioassay of crude plant extracts and controls against different fungal species

Table 28: IC₅₀ values of *in vitro* α -amylase inhibitory activity exhibited by the extracts (ME and HWE) of different samples at different concentrations.

Table 28A: ANOVA of data present in Table 28

Table 29: *In vitro* α -amylase inhibitory activity (%) of different plant extracts (ME and HWE) and acarbose (positive control) at different concentrations

Table 30: Effect of different plant extracts on body weight in diabetic rats

Table 31: Effect of different plant extracts on blood glucose level in streptozotocin induced diabetic rats

Table 32: Effect of different plant extracts on serum lipid profile in normal and streptozotocin induced diabetic rats

Table 33: Effect of different plant extracts on liver enzymes (SGPT and SGOT level) in normal and streptozotocin induced diabetic rats

Table 34: Effect of different plant extracts on serum urea and creatinine level of normal and streptozotocin induced diabetic rats

Table 35: Colour, texture and weight of broad fractions of *N. cordifolia* and *C. dentatus* eluted using different solvent

Table 36: IC₅₀ values of DPPH radical scavenging activity exhibited by the eluted fractions of *C. dentatus* and *N. cordifolia*

Table 36A: ANOVA of data present in Table 36

Table 37: IC₅₀ values of α -amylase inhibitory activity exhibited by the eluted fractions of *C. dentatus* and *N. cordifolia*

Table 37A: ANOVA of data present in Table 37

Table 38: Compounds identified in the CdB fraction of *Cyclosorus dentatus*

Table 39: Compound identified in the CdC fraction of *Cyclosorus dentatus*

Table 40: Compound identified in the CdD fraction of *Cyclosorus dentatus*

Table 41: Compound identified in the CdE fraction of *Cyclosorus dentatus*

Table 42: Compounds identified in the NcB extract of *Nephrolepis cordifolia*

Table 43: Compounds identified in the NcC extract of *Nephrolepis cordifolia*

Table 44: Compounds identified in the NcD extract of *Nephrolepis cordifolia*

Table 45: Compounds identified in the NcE extract of *Nephrolepis cordifolia*

Table 46: Docking score of commercial drug and identified compounds in NcC against IV4S protein

Table 47: Docking score of compounds identified in NcC against IV4S protein

Table 48 & 49: Docking score of compounds identified in *CdC* against IV4S protein

Table 50: Docking score of commercial drug and compounds identified in *NcC* against 2JJK protein

Table 51: Docking score of compounds identified in *NcC* against 2JJK protein

Table 52 & 53: Docking score of compounds identified in *CdC* against 2JJK protein

Table 54: Docking score of commercial drug and compounds identified in *NcC* against 2BEL protein

Table 55: Docking score of compounds identified in *NcC* against 2BEL protein

Table 56 & 57: Docking score of compounds identified in *CdC* against 2BEL protein

Table 58: Docking score of commercial drug and compounds identified in *NcC* against 3DZY protein

Table 59: Docking score of compounds identified in *NcC* against 3DZY protein

Table 60 & 61: Docking score of compounds identified in *CdC* against 3DZY protein