

LIST OF FIGURES

Figure 1A: Collection sites of neighboring areas of Darjeeling town

Figure 1B: Collection sites of Mungpoo subdivisions

Figure 1C: Collection sites of Siliguri (Shibmandir neighboring areas)

Figure 2: GIS location map of the collection sites

Figure 3: Plant samples collected for the study

Figure 4: Dried and powdered frond samples of the study plants

Figure 5: Illustration of maintenance and handling of rats

Figure 6: Total flavonoid and tannin content in the crude powdered samples

Figure 7: Comparison of total and reducing sugar content in the powdered samples

Figure 8: Extraction yield expressed in percentage (%) of different plants in various solvent system

Figure 9&10: DPPH radical scavenging activity (%) exhibited by different plant extracts

Figure 11&12: Hydrogen peroxide scavenging activity (%) exhibited by different plant extracts at varying concentration

Figure 13&14: Nitric oxide radical scavenging activity (%) exhibited by different plant extracts at varying concentration

Figure 15&16: Superoxide radical scavenging activity (%) exhibited by different plants extracts at varying concentration.

Figure 17&18: Ferric reducing antioxidant power exhibited by different plant extracts at varying concentration.

Figure 19&20: Loading plot of the different phytochemicals and DPPH radical scavenging activity

Figure 21&22: Loading plot of the different phytochemicals and hydrogen peroxide scavenging activity

Figure 23&24: Loading plot of the different phytochemicals and superoxide radical scavenging activity

Figure 25&26: Loading plot of the different phytochemicals and nitric oxide radical scavenging activity

Figure 27&28: Loading plot of the different phytochemicals and ferric reducing antioxidant power activity

Figure 29: Standard phenolics detected through HPLC analysis

Figure 30: HPLC fingerprints of the phenolics present in the plant extracts *N. cordifolia* and *C. dentatus*

Figure 31: HPLC fingerprints of the phenolics present in the plant extracts *D. quercifolia* and *P. cuspidatus*

Figure 32: HPLC fingerprints of the phenolics present in the plant extracts *Dicranopteris linearis* and *P. biaurita*

Figure 33: HPLC fingerprints of the phenolics present in the plant extracts *P. vittata* and *M. punctatum*

Figure 34: Antibacterial activity of plant extracts (a: 500mg/ml; b: 250mg/ml; c:100mg/ml) against *Bacillus cereus*.

Figure 35: Antibacterial activity of plant extract (a: 500mg/ml; b: 250mg/ml; c: 100mg/ml) against *Bacillus megaterium*

Figure 36: Antibacterial activity of plant extract (a: 500mg/ml; b: 250mg/ml; c: 100mg/ml) against *Burkholderia symbiont*

Figure 37: Antibacterial activity of plant extract (a: 500mg/ml; b: 250mg/ml; c: 100mg/ml) against *Serratia marcescens*

Figure 38: Antifungal activity against *Alternaria alternata*

Figure 39: Antifungal activity against *Curvularia lunata*

Figure 40: Antifungal activity against *Fusarium oxysporum*

Figure 41: α -Amylase inhibitory activity (%) exhibited by different plant extracts at varying concentration

Figure 42: Illustration of column chromatography and fractions collected

Figure 43: DPPH radical scavenging activity (%) exhibited by the eluted fractions of *C. dentatus* and *N. cordifolia* at varying concentration

Figure 44: α -amylase inhibitory activity (%) exhibited by the eluted fractions of *C. dentatus* and *N. cordifolia* at varying concentration

Figure 45(I-IX): GC-MS Total ion chromatogram (TIC) of *CdB* fraction of *C. dentatus* and compounds identified in *CdB* fraction

Figure 46(I-VIII): GC-MS Total ion chromatogram (TIC) of *CdC* fraction of *C. dentatus* and compounds identified in *CdC* fraction

Figure 47(I-IX): GC-MS Total ion chromatogram (TIC) of *CdD* fraction of *C. dentatus* and compounds identified in *CdD* fraction

Figure 48(I-VI): GC-MS Total ion chromatogram (TIC) of *CdE* fraction of *C. dentatus* and compounds identified in *CdE* fraction

Figure 49(I-VIII): GC-MS Total ion chromatogram (TIC) of *NcB* fraction of *N. cordifolia* and compounds identified in *NcB* fraction

Figure 50(I-VI): GC-MS Total ion chromatogram (TIC) of *NcC* fraction of *N. cordifolia* and compounds identified in *NcC* fraction

Figure 51(I-VI): GC-MS Total ion chromatogram (TIC) of *NcD* fraction of *N. cordifolia* and compounds identified in *NcD* fraction

Figure 52(I-VI): GC-MS Total ion chromatogram (TIC) of *NcE* fraction of *N. cordifolia* and compounds identified in *NcE* fraction

Figure 53: Binding mode with highest binding energy revealing respective interactions of commercial drug and possible potent compound identified in fraction *NcC* and *CdC* into the catalytic binding sites of IV4S. A,B: Metformin C,D: Corynan-17-ol, 18,19- didehydro-10-methoxy-acetate (ester) E,F: Propanoic acid, 2-(3-acetoxy-4,4,14-trimethylandro-8-en-17-yl)- .

Figure 54: Binding mode with highest binding energy revealing respective interactions of commercial drug and possible potent compound identified in fraction *NcC* and *CdC* into the catalytic binding sites of 2JJK. A,B: Metformin C,D: Corynan-17-ol, 18,19- didehydro-10-methoxy-acetate (ester) E,F: Propanoic acid, 2-(3-acetoxy-4,4,14-trimethylandro-8-en-17-yl)- .

Figure 55: Binding mode with highest binding energy revealing respective interactions of commercial drug and possible potent compound identified in fraction *NcC* and *CdC* into the catalytic binding sites of 2BEL. A,B: Metformin C,D: Corynan-17-ol, 18,19- didehydro-10-

methoxy-acetate (ester) E,F: Propanoic acid, 2-(3-acetoxy-4,4,14-trimethylandro-8-en-17-yl)- .

Figure 56: Binding mode with highest binding energy revealing respective interactions of commercial drug and possible potent compound identified in fraction *NcC* and *CdC* into the catalytic binding sites of 3DZY. A,B: Metformin C,D: Corynan-17-ol, 18,19- didehydro-10-methoxy-acetate (ester) E,F: Propanoic acid, 2-(3-acetoxy-4,4,14-trimethylandro-8-en-17-yl)- .