

LIST OF FIGURES

CHAPTERS	FIGURES	PAGE NO.
Chapter IV	<p>Figure 1. Plot of limiting ionic conductance (λ^{\pm}_o) of [emim] CH₃SO₃ (◆), ionic Walden product ($\lambda^{\pm}_o\eta$) of [emim]⁺ (●) and (CH₃SO₃)⁻ (▲) in n-Propanol, n-Butanol and n-Pentanol respectively at 298.15 K</p> <p>Figure 2. Plot of Diffusion Coefficient (<i>D</i>)(solid line) of [emim]⁺(▲) and (CH₃SO₃)⁻ (◆) and ionic mobility (<i>i</i>)(dotted line) of [emim]⁺(Δ) and (CH₃SO₃)⁻ (◇) in n-Propanol, n-Butanol and n-Pentanol respectively at 298.15K.</p> <p>Figure 3. Plot of Molar Refraction (<i>R_M</i>) of [emim]CH₃SO₃ (—■—), in n-Propanol, n-Butanol and n-Pentanol respectively at 298.15K</p>	149-150
Chapter V	<p>Figure 1. Plot of limiting apparent molar volume (ϕ_v^0) for glycine (◆), alanine (■) and valine (▲) against mass fraction of aqueous ionic liquid solution.</p> <p>Figure 2. Plot of Viscosity <i>B</i>-coefficient for glycine (◆), alanine (■) and valine (▲) against mass fraction of aqueous ionic liquid solution.</p> <p>Figure 3. Plot of limiting molar compressibility (ϕ_k^0) for glycine (◆), alanine (■) and valine (▲) against mass fraction of aqueous ionic liquid solution</p>	164-165
Chapter VI	<p>Figure 1. Plot of limiting apparent molar volume (ϕ_v^0) for nicotinic acid at 298.15K (◻), 308.15K (■), 318.15K (▲, and ascorbic acid at 298.15K (◇), 308.15 (□), 318.15K (Δ),</p>	186-188

	<p>against studied temp (T) in $w_I=0.001, 0.003$ and 0.005 mass fraction of aq. IL respectively.</p> <p>Figure 2. Plot of $(\partial\phi_E^0/\partial T)_P$ for nicotinic acid (◻) and ascorbic acid (■) against studied temp (T) in $w_I=0.001, 0.003$ and 0.005 mass fraction of aq. Ionic liquid.</p> <p>Figure 3. The plots of viscosity B-coefficient for nicotinic acid at 298.15K (◆), 308.15K (■), 318.15K (▲), and ascorbic acid at 298.15K (◇), 308.15 (□), 318.15K (Δ), against mass fraction ($w_I=0.001, 0.003$ and 0.005) of aq. IL respectively.</p> <p>Figure 4. The plots of R_M for nicotinic acid (—■—) and ascorbic acid (—◆—) in different mass fractions (w_1) of IL in aqueous mixture at 298.15K.</p> <p>Figure 5. The plots Limiting molar adiabatic compressibilities (ϕ_K^0) for Nicotinic Acid (—◆—) and Ascorbic Acid (—■—) in different mass fractions (w_1) of Ionic Liquid in aqueous mixture at 298.15K.</p>	
Chapter VII	<p>Figure 1. Plot of molar conductance (Λ) versus concentration (c) of [EMIm] NO₃, in DMF (—◻—), DMA (—■—) and DMSO (—▲—) at 298.15K.</p> <p>Figure 2. Plot of Limiting molar conductance (Λ_0) of [EMIm]NO₃ (—◆—), [EMIm] CH₃SO₃ (—■—), [EMIm]OTs (—◇—) and association constant (K_A) of [EMIm]NO₃ (---▲---), [EMIm] CH₃SO₃ (---*---), [EMIm] OTs (---○---) in DMF, DMA and DMSO respectively at 298.15K.</p> <p>Figure 3. Plot of Diffusion Coefficient (D) of NO₃⁻ (—■—), CH₃SO₃⁻ (—x—) and OTs⁻ (—Δ—) in different studied solvents at 298.15K</p> <p>Figure 4. Plot of ionic mobility (i) of NO₃⁻ (—■—), CH₃SO₃⁻ (—x—) and OTs⁻ (—Δ—) in different studied solvents at 298.15K</p>	206-208

	<p>Figure 5: Plot of Molar Refraction (R_M) of [EMIm]NO₃ (—◆—), [EMIm] CH₃SO₃ (—■—), [EMIm] OTs (—▲—) in DMF, DMA and DMSO respectively at 298.15K.</p>	
Chapter VIII	<p>Figure 1. Plot of limiting apparent molar volume (ϕ_V^0) for Histidine and Phenyl alanine at different mass fraction of aq. Nicotinic solution against different temperatures</p> <p>Figure 2. Plot of Viscosity B-coefficient for Phenyl alanine and Histidine at different temperatures vs mass fraction of aq. Nicotinic acid solution</p> <p>Figure 3. Plot of Absorbance vs Wavelength spectra of (a) Aq. Nicotinic acid solution (b) Histidine solution (c) Phenyl alanine solution (d) Aq. Nicotinic acid+Histidine solution (e) Aq. Nicotinic acid+Phenyl alanine solution.</p>	230-231
Chapter IX	<p>Figure 1. Plot of surface tension of ionic liquid corresponding to the added conc. of aq. β-cyclodextrin</p> <p>Figure 2. Plot of conductance of ionic liquid corresponding to the added conc of aq. β-cyclodextrin</p>	243