

LIST OF TABLES

CHAPTERS	TABLES	PAGE NO.
Chapter IV	<p>Table 1. Density (ρ), viscosity (η), refractive index (n_D) and dielectric constant (ϵ) of different pure solvents at 298.15 K.</p> <p>Table 2. Molar conductance (Λ) and the corresponding concentration (c) of the studied IL (1-ethyl-3-methyl imidazolium methanesulfonate) in different solvents at $T = 298.15$ K.</p> <p>Table 3. Limiting molar conductance (Λ_0), association constant (K_A), co-sphere diameter (R) and standard deviations of experimental Λ (δ) obtained of [emim]CH₃SO₃ in different solvents at 298.15 K.</p> <p>Table 4. Walden product ($\Lambda_0 \cdot \eta$) and Gibb's energy change (ΔG°) of [emim] CH₃SO₃ in different solvents at 298.15 K</p> <p>Table 5. Diffusion Coefficient (D) and ionic mobility (i) of [emim]⁺ and X⁻ (CH₃SO₃) in n-Propanol, n-Butanol and n-Pentanol at 298.15K</p> <p>Table 6. Limiting ionic conductance (λ^\pm_0), ionic Walden product ($\lambda^\pm_0 \eta$), crystallographic radii (r_c) and Stoke's radii (r_s) [emim]⁺ and X⁻(CH₃SO₃) n-Propanol, n-Butanol and n-Pentanol at 298.15 K</p> <p>Table 7. Experimental densities, refractive indices and molar refractions of 0.05(M) concentration [emim]X (where X= CH₃SO₃) in n-Propanol, n-Butanol, n-Pentanol at 298.15K.</p>	146-148

<p>Chapter V</p>	<p>Table 1. The values of Density (ρ), Viscosity (η), Refractive index (n_D), and Speed of sound (u) in different mass fraction of 1-ethylpyridinium tetrafluoroborate solution at 298.15K^a</p> <p>Table 2. Experimental values of Densities (ρ), Viscosities (η), Refractive Index (n_D) and Ultrasonic Speed (u) of L-Glycine, L-Alanine and L-Valine in different mass fraction of 1-ethylpyridinium tetrafluoroborate at 298.15K^a</p> <p>Table 3. Molality, apparent molar volume (ϕ_V), $(\eta/\eta_0 - 1)/m^{1/2}$, molar refraction ($R_M$), adiabatic compressibility (β) and apparent molal adiabatic compressibility (ϕ_K) of L-Glycine, L-Alanine, and L-Valine in 1-ethylpyridinium tetrafluoroborate at 298.15 K</p> <p>Table 4. Limiting apparent molar volumes (ϕ_V^0), experimental slopes (S_V^*), A, B coefficients, limiting apparent molar compressibility (ϕ_K^0), and experimental slope (S_K^*) of L-Glycine, L-Alanine, and L-Valine in aqueous 1-ethylpyridinium tetrafluoroborate at 298.15 K</p>	<p>158-163</p>
<p>Chapter VI</p>	<p>Table 1. Values of density (ρ), viscosity (η), refractive index (n_D) and ultrasonic speed (u) of aqueous Ionic Liquid in different mass fraction (w_I), at 298.15K to 318.15K^a</p> <p>Table 2. Experimental values of density (ρ), viscosity (η), refractive index and ultrasonic speed (u) of vitamins in different mass fraction of aqueous Ionic</p>	<p>176-185</p>

	<p>Liquid (w_1) at 298.15K to 318.15K^a respectively</p> <p>Table 3. Molality (m), apparent molar volume (ϕ_V), $(\eta_r - 1)/\sqrt{m}$, molar refraction and apparent molar adiabatic compressibility (ϕ_K) of vitamins in different mass fraction of aqueous Ionic Liquid (w_1) at different temperatures respectively</p> <p>Table 4. Limiting apparent molal volumes (ϕ_V^0), experimental slopes (S_V^*), viscosity A, B-coefficients, limiting partial molal adiabatic compressibilities (ϕ_K^0), and experimental slopes (S_K^*) of vitamins in different mass fraction of aqueous Ionic Liquid (w_1) at 298.15K to 318.15K respectively</p> <p>Table 5. Values of empirical coefficients (a_0, a_1, and a_2) of Equation 4 for vitamins in different mass fraction of aqueous IL (w_1) at 298.15K to 318.15K respectively</p> <p>Table 6. Limiting apparent molal expansibilities (ϕ_E^0) for vitamins in different mass fraction of aqueous IL (w_1) at 298.15K to 318.15K respectively</p> <p>Table 7. Values of dB/dT for the vitamins in different mass fraction of aqueous ionic liquid (w_1) at 298.15 to 318.15K respectively</p>	
<p>Chapter VII</p>	<p>Table 1. Sample provenance and purity</p> <p>Table 2. Experimental and literature [28] values of density (ρ), viscosity (η_0), relative permittivity (ϵ) and refractive index (n_D) of the solvents at 298.15K and 0.101MPa pressure</p> <p>Table 3. The concentration (m) and molar conductance (Λ) of [EMIm]X (where X=NO₃, CH₃SO₃, OTs) in DMF, DMA</p>	<p>199-205</p>

	<p>and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 4. Limiting molar conductance (Λ_0), association constant (K_A), co-sphere diameter (R) and standard deviations of experimental (δ) of [EMIm]X (where X=NO₃, CH₃SO₃, OTs) in DMF, DMA and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 5. Walden product ($\Lambda_0 \cdot \eta_0$) and Gibbs energy change (ΔG^\ddagger) of [EMIm]X (where X=NO₃, CH₃SO₃, OTs) in DMF, DMA and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 6. Ionic limiting molar conductance (λ^\ddagger_0), ionic Walden product ($\lambda^\ddagger_0 \eta_0$), crystallographic radii (r_c) and Stoke's radii (r_s) [EMIm]⁺ and X⁻(NO₃,CH₃SO₃,OTs) DMF, DMA and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 7. Diffusion Coefficient (D) and ionic mobility (i) of [EMIm]⁺ and X⁻ (NO₃,CH₃SO₃,OTs) in DMF, DMA and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 8. Experimental densities, refractive indices and molar refractions of [EMIm]X (where X=NO₃, CH₃SO₃, OTs) in DMF, DMA and DMSO at 298.15K and 0.101MPa pressure</p> <p>Table 9. Stretching frequencies of the functional groups present in the pure solvent and change of frequency of [EMIm]X (where X=NO₃, CH₃SO₃, OTs) in DMF, DMA and DMSO.</p>	
Chapter VIII	<p>Table 1. Values of density (ρ), viscosity (η), at 298.15, 303.15, 308.15 K and refractive index (n_D) at 298.15 K in different mass fraction (w_1) of aqueous NA</p> <p>Table 2. Experimental values of density (ρ) and viscosity (η) at 298.15 K, 303.15 K, 308.15 K and Refractive index (n_D) at 298.15 K of amino acids in different mass fraction of</p>	217-227

	<p>aqueous NA (w_1)</p> <p>Table 3. Molality (m), apparent molar volume (ϕ_V) and $(\eta_r - 1)/\sqrt{m}$ at 298.15 K, 303.15 K, 308.15 K and molar refraction (R) at 298.15 K of amino acids in different mass fraction of aqueous NA (w_1)</p> <p>Table 4. Partial molar volumes (ϕ_V^0), experimental slopes (S_V^*), viscosity A- and B-coefficients of amino acids in different mass fraction of aqueous NA (w_1) at 298.15 K, 303.15 K and 308.15 K respectively</p> <p>Table 5. Values of empirical coefficients (a_0, a_1, and a_2) of Equation 4 for amino acids in different mass fraction of aqueous vitamin (w_1) at 298.15K to 308.15K respectively</p> <p>Table 6. Limiting apparent molal expansibilities (ϕ_E^0) for amino acids in different mass fraction of aqueous vitamin (w_1) at 298.15K to 308.15K respectively</p> <p>Table 7. Values of dB/dT for the amino acids in different mass fraction of aqueous vitamin (w_1) at 298.15 to 308.15K respectively</p>	
<p>Chapter IX</p>	<p>Table 1. Values of surface tension at the break point (γ) with corresponding concentration of IL in different mass fraction of aqueous β-cyclodextrin at 298.15K^a</p> <p>Table 2. Values of Specific conductance at the break point (κ) with corresponding concentration of IL in different mass fraction of aqueous β-cyclodextrin at 298.15K^a</p> <p>Table 3. Values of ϕ_1^0, ϕ_V^0(aqueous), $\mu^{0\#}$, $T\Delta S^\#$, $\Delta H^\#$, for IL in different mass fraction of aqueous β-CD (w_1) at 298.15 to 308.15K respectively</p>	<p>241-242</p>