

LIST OF FIGURES

- Figure 1:** Global scenario of mushroom production
- Figure 2:** Preparation of grain spawn for the cultivation of *Pleurotus* sp.
- Figure 3:** Mushroom production unit for large scale production of *Pleurotus* sp established at Immuno phytopathology laboratory, Department of Botany, NBU
- Figure 4:** Mycelial growth pattern of *Pleurotus* species on potato dextrose agar (A) *P ostreatus*, (B) *P sajor-caju*, (C) *P djamor* and (D) *P florida*
- Figure 5:** Mycelial growth rate of *Pleurotus* species on different media (incubation period 48h., temperature $28 \pm 2^{\circ}$ C)
- Figure 6:** Morphological and anatomical features of *P ostreatus* (A) showing the fruiting body, (B) T.S of gill (10X); (C) basidiospore attached with basidium at 40X; (D) basidium attached with the hymenophore and (E) Basidiospore attached to basidium.
- Figure 7:** Morphological and anatomical features of *P sajor-caju* (A) showing the fruiting body, (B) T.S of gill (10X); (C) basidiospore attached with basidium at 40X; (D) basidium attached with the hymenophore and (E) Basidiospore attached to basidium.
- Figure 8:** Morphological and anatomical features of *P. djamor* (A) showing the fruiting body, (B) T.S of gill 10X; (C) basidiospore attached with basidium at 40X; (D) basidium attached with the hymenophore and (E) Basidiospore attached to basidium.
- Figure 9:** Morphological and anatomical features of *P. florida* (A) showing the fruiting body, (B) T.S of gill 10X; (C) basidiospore attached with basidium at 40X; (D) basidium attached with the hymenophore and (E) Basidiospore attached to basidium
- Figure 10:** Chromatogram and sequence of 18S rDNA region *P ostreatus* (IPL/MC/PO-1) deposited in NCBI Genbank
- Figure 11:** 18S r DNA sequence alignments of *P ostreatus* (IPL/MC/PO-1) with other ex-type isolates. The conserved regions of the gene are demonstrated in different colours

- Figure 12:** Phylogenetic placement of *P. ostreatus* (IPL/MC/PO-1) with other ex-type strain sequences obtained from NCBI Genbank Database
- Figure 13:** Chromatogram and sequence of 18S rDNA region *P. sajor-caju* (IPL/MC/PS-1) deposited in NCBI Genbank
- Figure 14:** 18S r DNA sequence alignments of *P. sajor-caju* (IPL/MC/PS-1) with other ex-type isolates. The conserved regions of the gene are demonstrated in different colours
- Figure 15:** Phylogenetic placement of *P. sajor-caju* (IPL/MC/PS-1) with other ex-type strain sequences obtained from NCBI Genbank Database
- Figure 16:** Chromatogram and sequence of 18S rDNA region *P. djamor* (IPL/MC/PD-1) deposited in NCBI Genbank
- Figure 17:** 18S r DNA sequence alignments of *P. djamor* (IPL/MC/PD-1) with other ex-type isolates. The conserved regions of the gene are demonstrated in different colours
- Figure 18:** Phylogenetic placement of *P. djamor* (IPL/MC/PD-1) with other ex-type strain sequences obtained from NCBI Genbank Database
- Figure 19:** Chromatogram and sequence of 18S rDNA region *P. florida* (IPL/MC/PF-1) deposited in NCBI Genbank
- Figure 20:** 18S r DNA sequence alignments of *P. florida* (IPL/MC/PF-1) with other ex-type isolates. The conserved regions of the gene are demonstrated in different colours
- Figure 21:** Phylogenetic analysis of *P. florida* (IPL/MC/PF-1) with other ex-type strain sequences obtained from NCBI Genbank Database
- Figure 22:** Weather condition of North Bengal during 2013-2015 (A) temperature, (B) relative humidity and (C) rainfall throughout the year
- Figure 23:** Assessment of spawn run period for pin head initiation during cultivation of *P. ostreatus* throughout the year
- Figure 24:** Comparative study on yield of *P. ostreatus* grown in paddy straw, wheat straw and saw dust substrates during 2013 (A), 2014 (B) and 2015 (C)
- Figure 25:** (A-G) Cultivation of *P. ostreatus* in polypropylene bags. (A-D) pinhead stage, (E) early mature stage and (F & G) mature stage.

- Figure 26:** (A-H) Cultivation of *P. ostreatus* in different containers. (A-B): Polypropylene bags; (C-E) plastic bottles (F) broken glass goods; and (G-H) box made up of aluminium sheet
- Figure 27:** Assessment of spawn run period for pin head initiation during cultivation of *P. sajor-caju* throughout the year
- Figure 28:** Comparative study on yield of *P. sajor-caju* grown in paddy straw, wheat straw and saw dust substrates during 2013 (A), 2014 (B) and 2015 (C)
- Figure 29:** (A-I) Cultivation of *P. sajor-caju* (A-C) pinhead, (D-F) early mature stage and (G-I) showing the mature fruiting body.
- Figure 30:** (A-F) Cultivation practices of *P. sajor-caju* using different containers, (A-B) polypropylene bags; (C) plastic bottles (D) waste bottles and (E-F) paper box.
- Figure 31:** Assessment of spawn run period for pin head initiation during cultivation of *P. djamor* throughout the year
- Figure 32:** Comparative study on yield of *P. djamor* grown in paddy straw, wheat straw and saw dust substrates during 2013 (A), 2014 (B) and 2015 (C)
- Figure 33:** (A-K) Cultivation of *P. djamor* (pink oyster mushroom); (A-C) pinhead stage, (D-G) early mature stage and (H-K) mature stage ready for harvesting.
- Figure 34:** (A-G) Growth and development of fruitbody during cultivation of *P. djamor* in different containers; (A-B) polypropylene bags; (C-G) using waste plastic bottles
- Figure 35:** Assessment of spawn run period for pin head initiation during cultivation of *P. florida* throughout the year
- Figure 36:** Comparative study on yield of *P. djamor* grown in paddy straw, wheat straw and saw dust substrates during 2013 (A), 2014 (B) and 2015 (C)
- Figure 37:** (A-I) Cultivation of *P. florida* (A-C) pinhead stage, (D-G) early mature stage and (H&I) showing the mature stage.
- Figure 38:** (A-H) Cultivation practice of *P. florida* using different types of containers (A-B) polypropylene bags; (C-E) waste plastic bottles and (F-H) in paper box.

- Figure 39:** Comparative study of moisture content of (A) *P. ostreatus*, (B) *P. sajor-caju*, (C) *P. djamor* and (D) *P. florida*
- Figure 40:** Total sugar (A), reducing sugar (B) of *P. ostreatus* in different stages of its growth
- Figure 41:** Total soluble protein (A) and total lipid content (B) of *P. ostreatus* in different stages of its growth
- Figure 42:** SDS PAGE analysis of soluble protein of *P. ostreatus* showing the bands on the gel (A) and band intensity analysed using image lab software (B-D).
- Figure 43:** Total sugar (A) and reducing sugar (B) of *P. sajor-caju* in different stages of its growth
- Figure 44:** Total soluble protein (A) and lipid content (B) of *P. sajor-caju* in different stages of its growth
- Figure 45:** SDS PAGE analysis of soluble protein of *P. sajor-caju* showing the bands on the gel (A) and band intensity analysed using image lab software (B-D)
- Figure 46:** Total sugar (A) and reducing sugar (B) of *P. djamor* in different stages of its growth
- Figure 47:** Total soluble protein (A) and lipid content (B) of *P. djamor* in different stages of its growth
- Figure 48:** SDS PAGE analysis of the protein of *P. djamor* showing the bands on the gel (A) and band intensity analysed using image lab software (B-D)
- Figure 49:** Total sugar (A) and reducing sugar (B) of *P. florida* in different stages of its growth
- Figure 50:** Total soluble protein (A) and lipid content (B) of *P. florida* in different stages of its growth
- Figure 51:** SDS PAGE analysis of the protein of *P. florida* showing the bands on the gel (A) and band intensity analysed using image lab software (B-D)
- Figure 52:** Cultivation of *P. ostreatus* using pruned tea leaves; (A) fully colonized bag, (B) pinhead formation, (C) young stage and (D) mature fruiting body
- Figure 53:** (A-D) Cultivation of *P. ostreatus* on tea waste substrate in combination with paddy straw (A) fully colonized, (B) pinhead stage, (C) young fruiting body and (D) mature fruiting body

- Figure 54:** Cultivation of *Pleurotus sajor-caju* on pruned tea leaves (A-D) and in combine with paddy straw (E and F)
- Figure 55:** Moisture content of *P. ostreatus* grown on tea waste
- Figure 56:** Total sugar (A) reducing sugar (B) and total protein (C) content of *P. ostreatus* grown on tea waste
- Figure 57:** Antioxidant activity of *Pleurotus* species; (A) DPPH scavenging activity (B) Free radicle antioxidant power assay and (C) flavonoid activity
- Figure 58:** Induction of hyperglycaemia in white albino rats; (A &B) selected healthy rats of same size and weight kept in grouped, (C) induction of hyperglycaemia using STZ intraperitoneally, (D) oral treatment of mushroom, (E) treatment after taking blood from tail vein.
- Figure 59:** Evaluation of antihyperglycemic activity of *Pleurotus* species on induced rats (A) effect on body weight and (B) blood glucose level following induction
- Figure 60:** Changes in creatinine level following oral treatment of *Pleurotus* species in hyperglycaemic rats
- Figure 61:** Changes in blood urea level following oral treatment of *Pleurotus* species in hyperglycaemic rats
- Figure 62:** Liver enzyme serum glutamic pyruvate transaminase (SGPT) activity of STZ treated diabetic rats following oral treatment of *Pleurotus* species in relation to control
- Figure 63:** Liver enzyme serum glutamic oxaloacetic transaminase (SGOT) activity of STZ treated diabetic rats following oral treatment of *Pleurotus* species in relation to control
- Figure 64:** Effect of spent substrate of oyster mushroom on growth of *Capsicum chinense*; (A) control plants and (B) plants grown in spent substrate amended soil [after 15 days]; (C) untreated control and (D) plants grown spent substrates amended soil [after 45 days]; (E) flower, (F) developmental stages of fruit

- Figure 65:** Effect of spent mushroom substrates on the growth of *Capsicum annuum* L. (A) control; (B) fresh spent oyster mushroom substrate; (C) spent oyster mushroom substrate leachate; (D) spent weathered oyster mushroom substrate; (E) fresh button mushroom compost; (F) spent button mushroom compost leachate; (G) spent weathered button mushroom compost; (H) combined treatment of spent oyster mushroom and button mushroom substrate
- Figure 66:** Effect of spent mushroom substrates of oyster mushroom on *Solanum lycopersicum* (A& D) untreated control and (B &C) SMS treated and *Amaranthus* (C) Untreated control and (D) SMS treated
- Figure 67:** Effect of spent mushroom substrate on total chlorophyll (A) and carotenoid content (B) of *C. chinense*.
- Figure 68:** Effect of spent mushroom substrate on soluble protein (A) and carotenoid contents (B) of *Capsicum annuum*
- Figure 69:** Schematic diagram of post-harvest processing of *Pleurotus* species
- Figure 70:** Harvesting and packaging of oyster mushroom
- Figure 71:** Different recipes of oyster mushroom (A) mushroom tomato curry, (B) vegetable mushroom, (C) egg mushroom scramble (D) onion mushroom pakoda, and (E-F) mushroom soup.
- Figure 72:** Dry mushroom of different *Pleurotus* species
- Figure 73:** Mushroom powder prepared from dried *Pleurotus* species
- Figure 74:** Mushroom biscuit prepared using dried oyster mushroom powder
- Figure 75:** Evaluation of biochemical constituents of oyster mushroom powder (A) total sugar, (B) reducing sugar
- Figure 76:** Evaluation of biochemical constituents of oyster mushroom powder (A) total protein, (B) total lipid and (C) dietary fibre
- Figure 77:** Ink Cap disease caused by *Coprinus* sp.
- Figure 78:** Cobweb disease during cultivation of oyster mushroom
- Figure 79:** Green Mould disease caused by *Trichoderma* sp.
- Figure 80:** Yellow mould disease during cultivation of *Pleurotus* species

Figure 81: Bacterial blotch disease

Figure 82: Pest of mushroom Sciarid fly (A), phorid fly (B) and beetle fly (C&D)

Figure 83: Protective measures against insects and pests during mushroom cultivation practice inside the production unit

Figure 84: GPS location of mushroom farms established by the rural people after proper training in different districts of North Bengal (A) Mainaguri; (B) Belakoba; (C) Dhupguri; (D) Ambari; (E) Naxalbari, (F) Bagdogra, (G) Jalpaiguri and (H) Batabasi

Figure 85: Promotion of oyster mushroom cultivation in different parts of North Bengal (A) Supak Singha Naxalbari, (B) Baburam Sarkar, Maynaguri, (C) Topu Barman, Belakoba, (D) Amit Kumar Sarkar, Ambari and (E) Jagannath Baroi, Bagdogra

Figure 86: Demonstration of oyster mushroom cultivation practices during Krishi Mela organized by the Department of Agriculture, Govt. of West Bengal at Atharakhai Gram Panchayet, Shivmandir

Figure 87: Hands on training of cultivation process of oyster mushroom to the post-graduate students and unemployed youth (A) training for the unemployed youth, (B) students of Ananda Chandra College, Jalpaiguri, (C-D) Post-graduate students of NBU.

Figure 88: Mushroom sell at local markets; (A) fulbari haat, (B) Naxalbari, (C) Batabasi haat, (D) Barivasa bazar and (E &F) departmental stores