

BIBLIOGRAPHY

Chapter 1.

- [1] G.M. Cragg, D.J. Newman, Biodiversity: A continuing source of novel drug leads, *Pure Appl Chem.* 77 (2005) 7-24.
- [2] H.F. Ji, X.J. Li, H.Y. Zhang, Natural products and drug discovery, *EMBO rep.* 10 (2009) 194-200.
- [3] D.J. Newman, Natural products as leads to potential drugs: an old process or the new hope for drug discovery?, *J Med Chem.* 51 (2008) 2589-2599.
- [4] W.J. Zhang, G.H. Liu, Coevolution: A synergy in biology and ecology, *Selforganizology.* 2 (2015) 35-38.
- [5] S.J. Cutler, H.G. Cutler, *Biologically Active Natural Products: Pharmaceuticals*, CRC Press, New York, 2000.
- [6] B. delasHeras, B. Rodríguez, L. Boscá, A.M. Villar, Terpenoids: Sources, Structure Elucidation and Therapeutic Potential in Inflammation, *Curr Top Med Chem.* 3 (2003) 171-185
- [7] D. Tholl, Biosynthesis and biological functions of terpenoids in plants, *Adv Biochem Eng Biotechnol.* 148 (2015) 63-106.
- [8] Z.L. Ruzicka, The isoprene rule and the biogenesis of terpenic compounds, *Experientia.* 9 (1953) 357–367.
- [9] J. Gershenzon, W. Kreis, Biosynthesis of monoterpenes, sesquiterpenes, diterpenes, sterols, cardiac glycosides and steroid saponins, *Biochemistry of plant secondary metabolites,*

Annual Plant Reviews, in: M. Wink (Ed), Sheffield Academic Press: Sheffield, UK, Vol. 2, 1999, pp. 222-299.

- [10] J. Gershenzon, N. Dudareva, The function of terpene natural products in the natural world, *Nat Chem Biol.* 3 (2007) 408–414.
- [11] M. Rohmer, The Discovery of a Mevalonate-Independent Pathway for Isoprenoid Biosynthesis in Bacteria, Algae and Higher Plants, *Nat Prod Rep.* 16 (1999) 565-574.
- [12] I.L Finar, Stereochemistry and the chemistry of natural products, Pearson Education, Singapore, Vol. 2, 2004.
- [13] P. Crabbé, Ord and Cd in chemistry and biochemistry, Academic Press, London, 1972.
- [14] N. Harada, K. Nakanishi, Circular dichroic spectroscopy, exciton coupling in organic stereochemistry, University Science Books, Mill Valley, CA, 1983.
- [15] E.L. Eliel, S.H. Wilen, Stereochemistry of organic compounds, Wiley-Interscience, New York, 1994.
- [16] A.A. Saddiq, S.A. Khayyat, Chemical and antimicrobial studies of monoterpenes: Citral, *Pestic Biochem Physiol.* 98 (2010) 89–93.
- [17] C. de Bona da Silva, S.S. Guterres, V. Weisheimer, E.E.S. Schapoval, Antifungal activity of the lemongrass oil and citral against *Candida* spp., *Braz J Infect Dis.* 12 (2008) 63-66.
- [18] K. Ghosh, Anticancer effect of lemongrass oil and citral on cervical cancer cell lines, *Phcog Commn.* 3 (2013) 41-48.

- [19] W. Chaouki, D.Y. Leger, B. Liagre, J.L. Beneytout, M. Hmamouchi, Citral inhibits cell proliferation and induces apoptosis and cell cycle arrest in MCF-7 cells, *Fundam Clin Pharmacol.* 23 (2009) 549–556.
- [20] H. Xia, W. Liang, Q. Song, X. Chen, X. Chen, J. Hong, . The in vitro study of apoptosis in NB4 cell induced by citral, *Cytotechnology* 65 (2013) 49-57.
- [21] S. Zeng, A. Kapur, M.S. Patankar, M.P. Xiong, Formulation, characterization, and antitumor properties of trans- and cis-citral in the 4T1 breast cancer Xenograft mouse model, *Pharm Res.* 32 (2015) 2548-2558.
- [22] P.L. Crowell, Monoterpenes in breast cancer chemoprevention, *Breast Cancer Res Treat.* 46 (1997) 191–197.
- [23] P.L. Crowell, Prevention and therapy of cancer by dietary monoterpenes, *J Nutr.* 129 (1999) 775S–778S.
- [24] M.N. Gould, Cancer chemoprevention and therapy by monoterpenes, *Environ. Health Perspect.* 105 (1997) 977–979.
- [25] K.H. Wagner, I. Elmadfa, Biological relevance of terpenoids-overview focusing on mono, di and tetraterpenes, *Ann Nutr Metab.* 47 (2003) 95-106.
- [26] H.S. Choi, H.S. Song, H. Ukeda, M. Sawamura, Radical scavenging activities of citrus essential oils and their components: detection using 1,1-diphenyl-2-picrylhydrazyl, *J Agric Food Chem.* 48 (2000) 4156–4161.
- [27] L. Caputi, E. Aprea, Use of terpenoids as natural flavouring compounds in food industry, *Recent Pat Food Nutr Agric.* 3 (2011) 9-16.

- [28] J.H. Joo, A.M. Jetten, Molecular mechanisms involved in farnesol-induced apoptosis, *Cancer Lett.* 287 (2010) 123-135.
- [29] L. Kromidas, E. Perrier, J. Flanagan, R. Rivero, I. Bonnet, Release of antimicrobial actives from microcapsules by the action of axillary bacteria, *Int J Cosmet Sci.* 28 (2006) 103-108.
- [30] A. Bahi, S.A Mansouri, E.A Memari, M.A Ameri, S.M Nurulain, S. Ojha, β -Caryophyllene, a CB₂ receptor agonist produces multiple change relevant to anxiety and depression in mice, *Physiol Behav.* 135 (2014) 119-124.
- [31] M. Seo, T. Koshiba, Complex regulation of ABA biosynthesis in plants, *trends in plant sci.* 7 (2002) 41-48.
- [32] T.R. Evans, S.B. Kaye, Retinoids: Present role and future potential, *Br J Cancer.* 80 (1999) 1-8.
- [33] R. Gupta, S.K. Chakrabarty, Gibberellic acid in plant Still a mystery unresolved, *Plant Signal Behav.* 8 (2013) e25504.
- [34] J.J. Johnson, Carnosol: A promising anti-cancer and anti-inflammatory agent, *Cancer Lett.* 305 (2011) 1-7.
- [35] S. Birtić, P. Dussort, F.X. Pierre, A.C. Bily, M. Roller, Carnosic acid, *Phytochemistry.* 115 (2015) 9-19.
- [36] G.E. Bartley, P.A. Scolnik, Plant carotenoids: pigments for photoprotection, visual Attraction, and human health, *Plant Cell.* 7 (1995) 1027-1038.

- [37] J. Grassmann, Terpenoids as plant antioxidants, *Vitam Horm.* 72 (2005) 505-535.
- [38] A.G. Atanasov, B. Waltenberger, E.M.P. Wenzig , T. Linder , C. Wawrosch, P. Uhrin, V. Temml, L. Wang, S. Schwaiger, E. Heiss, J.M. Rollinger, D. Schuster, J.M. Breuss, V. Bochkov, M.D. Mihovilovic, B. Kopp, R. Bauer, V.M. Dirsch, H. Stuppner, Discovery and resupply of pharmacologically active plant-derived natural products: a review, *Biotechnol Adv.* 33 (2015) 1582-1614.
- [39] M. Bollen, S. Keppens, W. Stalmans, Specific features of glycogen metabolism in the liver, *Biochem J.* 336 (1998) 19-31.
- [40] H.G. Hers, The control of glycogen metabolism in the liver, *Annu Rev Biochem.* 45 (1976) 167-190.
- [41] W. Stalmans, M. Laloux, H.G. Hers, The interaction of liver phosphorylase a with glucose and AMP, *Eur J Biochem.* 49 (1974) 415-427.
- [42] W. Pimenta, N. Nurjhan, P.A. Jansson, M. Stumvoll, J. Gerich, M. Korytkowski, Glycogen: its mode of formation and contribution to hepatic glucose output in postabsorptive humans, *Diabetologia.* 37 (1994) 697-702.
- [43] M.K. Hellerstein, R.A. Neese, P. Linfoot, M. Christiannsen, S. Turner, A. Letscher, Hepatic gluconeogenic fluxes and glycogen turnover during fasting in humans. A stable isotope study, *J Clin Invest.* 100 (1997) 1305-1319.
- [44] N.G. Oikonomakos, E.D. Chrysina, M.N. Kosmopoulou, D.D. Leonidas, Crystal structure of rabbit muscle glycogen phosphorylase a in complex with a potential hypoglycaemic drug at 2.0 Å resolution, *Biochim Biophys Acta.* 1647 (2003) 325-332.

- [45] V.L. Rath, M. Ammirati, D.E. Danley, J.L. Ekstrom, E.M. Gibbs, T.R. Hynes, A.M. Mathiowitz, R.K. McPherson, T.V. Olson, J.L. Treadway, D.J. Hoover, Human liver glycogen phosphorylase inhibitors bind at a new allosteric site, *Chem Biol.* 7 (2000) 677-682.
- [46] R. Kurukulasuriya, J.T. Link, D.J. Madar, Z. Pei, S.J. Richards, J.J. Rohde, A.J. Souers, B.G. Szczepankiewicz, Potential drug targets and progress towards pharmacologic inhibition of hepatic glucose production, *Curr Med Chem.* 10 (2003) 123-153.
- [47] Z. Liang, L. Zhang, L. Li, J. Liu, H. Li, L. Zhang, L. Chen, K. Cheng, M. Zheng, X. Wen, P. Zhang, J. Hao, Y. Gong, X. Zhang, X. Zhu, J. Chen, H. Liu, H. Jiang, C. Luo, H. Sun, Identification of pentacyclic triterpenes derivatives as potent inhibitors against glycogen phosphorylase based on 3D-QSAR studies, *Eur J Med Chem.* 46 (2011) 2011-2021.
- [48] X. Wen, H. Sun, J. Liu, K. Cheng, P. Zhang, L. Zhang, J. Hao, L. Zhang, P. Ni, S.E. Zographos, D.D. Leonidas, K.M. Alexacou, T. Gimisis, J.M. Hayes, N.G. Oikonomakos, Naturally occurring pentacyclic triterpenes as inhibitors of glycogen phosphorylase: synthesis, structure-activity relationships, and X-ray crystallographic studies, *J Med Chem.* 51 (2008) 3540-3554.
- [49] T. Hunter, Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling, *Cell.* 80 (1995) 225-236.
- [50] A. Alonso, J. Sasin, N. Bottini, I. Friedberg, I. Friedberg, A. Osterman, A. Godzik, T. Hunter, J. Dixon, T. Mustelin, Protein tyrosine phosphatases in the human genome, *Cell.* 117 (2004) 699-711.

- [51] Z.Y. Zhang, S.Y. Lee, PTP1B inhibitors as potential therapeutics in the treatment of type 2 diabetes and obesity, *Expert Opin Investig Drugs.* 12 (2003) 223-233.
- [52] A.P. Combs, Recent advances in the discovery of competitive protein tyrosine phosphatase 1B inhibitors for the treatment of diabetes, obesity, and cancer, *J Med Chem.* 53 (2010) 2333-2344.
- [53] S. Thareja, S. Aggarwal, T.R. Bhardwaj, M. Kumar, Protein tyrosine phosphatase 1B inhibitors: a molecular level legitimate approach for the management of diabetes mellitus, *Med Res Rev.* 32 (2012) 459-517.
- [54] R.E. Cebula, J.L. Blanchard, M.D. Boisclair, K. Pal, N.J. Bockovich, Synthesis and phosphatase inhibitory activity of analogs of sulfircin, *Bioorg Med Chem Lett.* 7 (1997) 2015-2020.
- [55] R.M. Chen, L.H. Hu, T.Y. An, J. Li, Q. Shen, Natural PTP1B inhibitors from *Broussonetia papyrifera*, *Bioorg Med Chem Lett.* 12 (2002) 3387-3390.
- [56] C.S. Jiang, L.F. Liang, Y.W. Guo, Natural products possessing protein tyrosine phosphatase 1B (PTP1B) inhibitory activity found in the last decades, *Acta Pharmacologica Sinica.* 33 (2012) 1217-1245.
- [57] J.J. Ramírez-Espinosa, M.Y. Rios, S.L. Martínez, F.L. Vallejo, J.L. Medina-Franco, P. Paoli, G. Camici, G. Navarrete-Vázquez, R. Ortiz-Andrade, S. Estrada-Soto, Antidiabetic activity of some pentacyclic acid triterpenoids, role of PTP-1B: in vitro, in silico, and in vivo approaches, *Eur J Med Chem.* 46 (2011) 2243-2251.

- [58] E.I. Solomon, J. Zhou, F. Neese, E.G. Pavel, New insights from spectroscopy into the structure/function relationships of lipoxygenases, *Chem Biol.* 4 (1997) 795-808.
- [59] H.W. Gardner, Biological roles and biochemistry of the lipoxygenase pathway, *Hortscience.* 30 (1995) 197-205.
- [60] R. Wisastr, F.J. Dekker, Inflammation, cancer and oxidative lipoxygenase activity are intimately linked, *Cancers.* 6 (2014) 1500-1521.
- [61] L.A. Dailey, P. Imming, 12-Lipoxygenase: classification, possible therapeutic benefits from inhibition, and inhibitors, *Curr Med Chem.* 6 (1999) 389-398.
- [62] V.E. Steele, C.A. Holmes, E.T. Hawk, L. Kopelovich, R.A. Lubet, J.A. Crowell, C.C. Sigman, G.J. Kelloff, Lipoxygenase inhibitors as potential cancer chemopreventives, *Cancer Epidem Biomar.* 8 (1999) 467-483.
- [63] B. Samuelsson, S.E. Dahlen, J.A. Lindgren, C.A. Rouzer, C.N. Serhan, Leukotrienes and lipoxins: structures, biosynthesis, and biological effects, *Science.* 237 (1987) 1171-1176.
- [64] X.Z. Ding, W.G. Tong, T.E. Adrian, 12-Lipoxygenase metabolite 12(S)-HETE stimulates human pancreatic cancer cell proliferation via protein tyrosine phosphorylation and ERK activation, *Int J Cancer.* 94 (2001) 630-636.
- [65] J.A. Cornicelli, B.K. Trivedi, 15-Lipoxygenase and its inhibition: a novel therapeutic target for vascular disease, *Curr Pharm Des.* 5 (1999) 11-20.
- [66] U.P. Kelavkar, J.B. Nixon, C. Cohen, D. Dillehay, T.E. Eling, K.F. Badr, Overexpression of 15-lipoxygenase-1 in PC-3 human prostate cancer cells increases tumorigenesis, *Carcinogenesis.* 22 (2001) 1765-1773.

- [67] B.D. Roebuck, D.S. Longnecker, K.J. Baumgartner, C.D. Thron, Carcinogen-induced lesions in the rat pancreas: effects of varying levels of essential fatty acid, *Cancer Res.* 45 (1985) 5252-5256.
- [68] J. Zhang, V.L. Go, High fat diet, lipid peroxidation, and pancreatic carcinogenesis, *Adv Exp Med Biol.* 399 (1996) 165-172.
- [69] D.P. Rose, Dietary fatty acids and cancer, *Am J Clin Nutr.* 66 (1997) 998S-1003S.
- [70] P.L. Zock, M.B. Katan, Linoleic acid intake and cancer risk: a review and meta-analysis, *Am J Clin Nutr.* 68 (1998) 142-153.
- [71] W.G. Tong, X.Z. Ding, R.C. Witt, T.E. Adrian, Lipoxygenase inhibitors attenuate growth of human pancreatic cancer xenografts and induce apoptosis through the mitochondrial pathway, *Mol Cancer Therapeutics.* 1 (2002) 929-935.
- [72] C. Kemal, P. Louis-Flamberg, R. Krupinski-Olsen, A.L. Shorter, Reductive inactivation soybean lipoxygenase-1 by catechols: A possible mechanism for regulation of lipoxygenase activity, *Biochemistry.* 26 (1987) 7064-7072.
- [73] R. Mogul, E. Johansen, T.R. Holman, Oleyl sulfate reveals allosteric inhibition of soybean lipoxygenase-1 and human 15-lipoxygenase, *Biochemistry.* 39 (2000) 4801-4807.
- [74] A.F.A. Cros, Ph.D. dissertation thesis: Action de l'alcool amylique sur l'organisme, University of Strasbourg, Strasbourg, 1863.
- [75] A. Crum-Brown, T.R. Fraser, On the connection between chemical constitution and physiological action. Part 1. On the physiological action of the ammonium bases, derived

from strychia, brucia, thebaia, codeia, morphia and nicotia, Trans R Soc Edinburgh. 25 (1868) 151-203.

[76] C. Richet, On the relationship between the toxicity and the physical properties of substances, Compt Rendus Seances Soc Biol. 9 (1893) 775-776.

[77] E. Overton, Osmotic properties of cells in the bearing on toxicology and pharmacology, Z Physik Chem. 22 (1897) 189-209.

[78] H. Meyer, On the theory of alcohol narcosis I. Which property of anesthetics gives them their narcotic activity?, Arch Exp Pathol Pharmakol. 42 (1899) 109-118.

[79] L.P. Hammett, Some relations between reaction rates and equilibrium constants, Chem Rev. 17 (1935) 125-136.

[80] L.P. Hammett, The effect of structure upon the reactions of organic compounds. benzene derivatives, J Am Chem Soc. 59 (1937) 96-103.

[81] R.W. Taft, Polar and steric substituent constants for aliphatic and o- Benzoate groups from rates of esterification and hydrolysis of esters¹, J Am Chem Soc. 74 (1952) 3120-3128.

[82] C. Hansch, T. Fujita, p-σ-π Analysis. A method for the correlation of biological activity and chemical structure, J Am Chem Soc. 86 (1964) 1616-1626.

[83] C. Hansch, Quantitative approach to biochemical structure-activity relationships, Acc Chem Res. 2 (1969) 232-239.

- [84] S.M. Free Jr, J.W. Wilson, A Mathematical contribution to structure-activity studies, *J Med Chem.* 7 (1964) 395-399.
- [85] T. Fujita, T. Ban, Structure-activity study of phenethylamines as substrates of biosynthetic enzymes of sympathetic transmitters, *J Med Chem.* 14 (1971) 148-152.
- [86] L.B. Kier, Molecular orbital theory in drug research, Academic Press, New York, 1971.
- [87] A.T. Balaban, F. Harary, The characteristic polynomial does not uniquely determine the topology of a molecule, *J Chem Doc.* 11 (1971) 258-259.
- [88] M. Randić, On the recognition of identical graphs representing molecular topology, *J Chem Phys.* 60 (1974) 3920-3928.
- [89] L.B. Kier, L.H. Hall, W.J. Murray, M. Randic, Molecular connectivity I: relationship to nonspecific local anesthesia, *J Pharm Sci.* 64 (1975) 1971-1974.
- [90] R.H. Rohrbaugh, P.C. Jurs, Descriptions of molecular shape applied in studies of structure/activity and structure/property relationships, *Anal Chim Acta.* 199 (1987) 99-109.
- [91] D.T. Stanton, P.C. Jurs, Development and use of charged partial surface area structural descriptors in computer-assisted quantitative structure-property relationship studies, *Anal Chem.* 62 (1990) 2323-2329.
- [92] R. Todeschini, M. Lasagni, E. Marengo, New molecular descriptors for 2D- and 3D- structures, *Theory J Chemom.* 8 (1994) 263-273.

- [93] A.R. Katritzky, L. Mu, V.S. Lobanov, M. Karelson, Correlation of boiling points with molecular structure. 1. A training set of 298 diverse organics and a test set of 9 simple inorganics, *J Phys Chem.* 100 (1996) 10400-10407.
- [94] A.M. Ferguson, T.W. Heritage, P. Jonathon, S.E. Pack, L. Phillips, J. Rogan, P.J. Snaith, EVA: a new theoretically based molecular descriptor for use in QSAR/QSPR analysis, *J Comput Aided Mol Des.* 11 (1997) 143-152.
- [95] J. Schuur, P. Selzer, J. Gasteiger, The coding of the three-dimensional structure of molecules by molecular transforms and its application to structure-spectra correlations and studies of biological activity, *J Chem Inf Comput Sci.* 36 (1996) 334-344.
- [96] V. Consonni, R. Todeschini, M. Pavan, Structure/response correlations and similarity/diversity analysis by GETAWAY descriptors. Part 1. Theory of the novel 3D molecular descriptors, *J Chem Inf Comput Sci.* 42 (2002) 682-692.
- [97] R.D. Cramer, D.E. Patterson, J.D. Bunce, Comparative molecular field analysis (CoMFA). 1. Effect of shape on binding of steroids to carrier proteins, *J Am Chem Soc.* 110 (1988) 5959-5967.
- [98] J. Gasteiger, Handbook of chemoinformatics. From data to knowledge in 4 volumes, Wiley-VCH, Weinheim, 2003.
- [99] T.I. Oprea, 3D QSAR modeling in drug design, in: P. Bultinck, H. De Winter, W. Langenaeker, J.P. Tollenaere (Eds.), Computational medicinal chemistry for drug discovery. Marcel Dekker, New York, 2004.
- [100] R. Franke, Theoretical Drug Design Methods, Elsevier, Amsterdam, 1984.

- [101] K. Osmialowski, J. Halkiewicz, A. Radecki, R. Kalisz, Quantum chemical parameters in correlation analysis of gas-liquid chromatographic retention indices of amines, *J Chromatogr.* 346 (1985) 53-60.
- [102] K. Fukui, *Theory of Orientation and Stereoselection*, Springer- Verlag, New York, 1975.
- [103] H. Sklenar, J. Jäger, Molecular structure-biological activity relationships on the basis of quantum-chemical calculations, *Int J Quant Chem.* 16 (1979) 467-484.
- [104] K. Tuppurainen, S. Lötzönen, R. Laatikainen, T. Vartiainen, U. Maran, M. Strandberg, T. Tamm, About the mutagenicity of chlorine-substituted furanones and halopropenals. A QSAR study using molecular orbital indices, *Mutat Res Fund Mol Mech Mut.* 247 (1991) 97-102.
- [105] I. Fleming, *Frontier orbitals and organic chemical reactions*, John Wiley & Sons, New York, 1976.
- [106] D.F.V. Lewis, C. Ioannides, D.V. Parke, Interaction of a series of nitriles with the alcohol inducible isoform of P450- computer analysis of structure-activity relationships, *Xenobiotica.* 24 (1994) 401-408.
- [107] M. Karelson, V.S. Lobanov, Quantum-Chemical Descriptors in QSAR/QSPR Studies, *Chem Rev.* 96 (1996) 1027-1043.

Chapter 2.

- [1] S.B. Mahato, S. Sen, Advances in triterpenoid research, 1990-1994, *Phytochemistry*. 44 (1997) 1185-1236.
- [2] I.H. Hall, K.H. Lee, C.O. Starnes, Y. Sumida, R.Y. Wu, T.G. Waddell, J.W. Cochran, K.G. Gerhart, Anti- inflammatory activity of sesquiterpene lactones and related compounds, *J Pharm Sci.* 68 (1979) 537-542.
- [3] C.H. Chen, L.M. Yang, T.T. Lee, Y.C. Shen, D.C. Zhang, D.J. Pan, A.T. McPhail, D.R. McPhail, S.Y. Liu, D.H. Li, Antitumor agents-CLI. Bis(helenalanyl)glutarate and bis(isoalantodiol- B)glutarate, potent inhibitors of human DNA topoisomerase II, *Bioorg Med Chem.* 2 (1994) 137-145.
- [4] K. Hayashi, H. Hayashi, N. Hiraoka, Y. Ikeshiro, Inhibitory activity of soyasaponin II on virus replication in vitro, *Planta Med.* 63 (1997) 102-105.
- [5] M.C. Recio, R.M. Giner, L. Uriburu, S. Manez, M. Cerda, J.R. De la Fuente, J.L. Rios, In vivo activity of pseudoguaianolide sesquiterpene lactones in acute and chronic inflammation, *Life Sci.* 66 (2000) 2509-2518.
- [6] C.L. Cantrell, S.G. Franzblau, N.H. Fischer, Antimycobacterial plant terpenoids, *Planta Med.* 67 (2001) 685-694.
- [7] N. Osaki, T. Koyano, T. Kowithayakorn, M. Hayashi, K. Komiyama, M. Ishibashi, Sesquiterpenoids and plasmin-inhibitory flavonoids from *Blumea balsamifera*, *J Nat Prod.* 68 (2005) 447-449.
- [8] D.B. Stierle, A.A. Stierle, T. Girtsman, K. McIntyre, J. Nichols, Caspase-1 and -3 inhibiting drimane sesquiterpenoids from the extremophilic fungus *Penicillium solitum*, *J Nat Prod.* 75 (2012) 262-266.

- [9] Q.Y. Ma, Y.C. Chen, S.Z. Huang, Z.K. Guo, H.F. Dai, Y. Hua, Y.X. Zhao, Two new guaiane sesquiterpenoids from *Daphne holosericea* (Diels) Hamaya, *Molecules.* 19 (2014) 14266-14272.
- [10] Q.Q. Tao, K. Ma, L. Bao, K. Wang, J.J. Han, J.X. Zhang, C.Y. Huang, H.W. Liu, New sesquiterpenoids from the edible mushroom *Pleurotus cystidiosus* and their inhibitory activity against α -glucosidase and PTP1B, *Fitoterapia.* 111 (2016) 29-35.
- [11] K. Dimas, C. Demetzos, M. Marsellos, R. Sotiriadou, M. Malamas, D. Kokkinopoulos, Cytotoxic activity of labdane typed diterpenes against human leukemic cell lines in vitro, *Planta Med.* 64 (1998) 208-211.
- [12] R. Batista, E. Chiari, A.B. de Oliveira, Trypanosomicidal kaurane diterpenes from *Wedelia paludosa*, *Planta Med.* 65 (1999) 283-284.
- [13] M. Singh, M. Pal, R.P. Sharma, Biological activity of the labdane diterpenes, *Planta Med.* 65 (1999) 2-8.
- [14] M. DellaGreca, A. Fiorentino, M. Isidori, P. Monaco, A. Zarrelli, Antialgal ent-labdane diterpenes from *Ruppia maritime*, *Phytochemistry.* 55 (2000) 909-913.
- [15] R. Tanaka, H. Ohtsu, M. Iwamoto, T. Minami, H. Tokuda, H. Nishino, S. Matsunaga, A. Yoshitake, Cancer chemopreventive agents, labdane diterpenoids from the stem bark of *Thuja standishii* (Gord.) Carr, *Cancer Lett.* 161 (2000) 165-170.
- [16] A. Navarro, B. de las Heras, A.M. Villar, Andalusol, a diterpenoid with anti-inflammatory activity from *Sideritis foetens* Clemen, *Z Naturforsch.* 52c (1997) 844-849.
- [17] M.A. Fernandez, M.P. Tornos, M.D. Garcia, B. de las Heras, A.M. Villar, M.T. Saenz, Anti-inflammatory activity of abietic acid, a diterpene isolated from *Pimenta racemosa* var. *grissea*, *J Pharm Pharmacol.* 53 (2001) 867-872.

- [18] X. Chen, J. Ding, Y.M. Ye, J.S. Zhang, Bioactive abietane and seco-abietane diterpenoids from *Salvia prionitis*, *J Nat Prod.* 65 (2002) 1016-1020.
- [19] M. Na, W.K. Oh, Y.H. Kim, X.F. Cai, S. Kim, B.Y. Kim, J.S. Ahn, Inhibition of protein tyrosine phosphatase 1B by diterpenoids isolated from *Acanthopanax koreanum*, *Bioorg Med Chem Lett.* 16 (2006) 3061-3064.
- [20] A. Leverrier, M.T. Martin, C. Servy, J. Ouazzani, P. Retailleau, K. Awang, M.R. Mukhtar, F. Guérinne, M. Litaudon, Rearranged diterpenoids from the biotransformation of ent-trachyloban-18-oic acid by *Rhizopus arrhizus*, *J Nat Prod.* 73 (2010) 1121-1125.
- [21] L. Yang, L. Qiao, C. Ji, D. Xie, N.B. Gong, Y. Lu, J. Zhang, J. Dai, S. Guo, Antidepressant abietane diterpenoids from chinese eaglewood, *J Nat Prod.* 76 (2013) 216-222.
- [22] J.X. Zhao, C.P. Liu, W.Y. Qi, M.L. Han, Y.S. Han, M.A. Wainberg, J.M. Yue, Eurifoloids A-R, structurally diverse diterpenoids from *Euphorbia neriifolia*, *J Nat Prod.* 77 (2014) 2224-2233.
- [23] G. Luo, Q.Ye, B. Du, F. Wang, G. Zhang, Y. Luo, Iridoid glucosides and diterpenoids from *Caryopteris glutinosa*, *J Nat Prod.* 79 (2016) 886-893.
- [24] H. Nishino, K. Yoshioka, A. Iwashima, H. Takizawa, S. Konishi, H. Okamoto, H. Okabe, S. Shibata, H. Fujiki, T. Sugimura, Glycyrrhetic acid inhibits tumor promoting activity of teleocidin and 12-O-tetradecanoylphorbol-13-acetate in two-stage mouse skin carcinogenesis, *Jpn J Cancer Res.* 77 (1986) 33-38.
- [25] H. Tokuda, H. Ohigashi, K. Koshimizu, Y. Ito, Inhibitory effects of ursolic and oleanolic acid on skin tumor promotion by 12-O-tetradecanoylphorbol-13-acetate, *Cancer Lett.* 33 (1986) 279-285.

- [26] M.C. Recio, R.M. Giner, S. Manez, J.L. Rios, Structural requirements for the anti-inflammatory activity of natural triterpenoids, *Planta Med.* 61 (1995) 182-185.
- [27] F. Soler, C. Poujade, M. Evers, J.C. Carry, Y. Henin, A. Bousseau, T. Huet, R. Pauwels, E. De Clercq, J.F. Mayaux, J.B. Le Pecq, N. Dereu, Betulinic acid derivatives: a new class of specific inhibitors of human immunodeficiency virus type 1 entry, *J Med Chem.* 39 (1996) 1069-1083.
- [28] H. Ito, S. Onoue, Y. Miyake, T. Yoshida, Iridal-type triterpenoids with ichthyotoxic activity from *Belamcanda chinensis*, *J Nat Prod.* 62 (1999) 89-93.
- [29] A. Rajic, G. Kweifio-Okai, T. Macrides, R.M. Sandeman, D.S. Chandler, G.M. Polya, Inhibition of serine proteases by anti-inflammatory triterpenoids, *Planta Med.* 66 (2000) 206-210.
- [30] C.L. Cantrell, S.G. Franzblau, N.H. Fischer, Antimycobacterial plant terpenoids, *Planta Med.* 67 (2001) 685- 694.
- [31] T. Fujioka, Y. Kashiwada, R.E. Kilkuskie, L.M. Cosentino, L.M. Ballas, J.B. Jiang, W.P. Janzen, I.S. Chen, K.H. Lee, Anti-AIDS agents, 11. Betulinic acid and platanic acid as anti-HIV principles from *Syzygium claviflorum*, and the anti-HIV activity of structurally related triterpenoids, *J Nat Prod.* 57 (1994) 243-247.
- [32] Z.Z. Ma, Y. Hano, T. Nomura, Y.J. Chen, Three new triterpenoids from *Peganum nigellastrum*, *J Nat Prod.* 63 (2000) 390-392.
- [33] S. Wada, A. Iida, R. Tanaka, Screening of triterpenoids isolated from *Phyllanthus flexuosus* for DNA topoisomerase inhibitory activity, *J Nat Prod.* 64 (2001) 1545-1547.
- [34] K. Yoshikawa, M. Inoue, Y. Matsumoto, C. Sakakibara, H. Miyatake, H. Matsumoto, S. Arihara, Lanostane triterpenoids and triterpene glycosides from the fruit body of

Fomitopsis pinicola and their inhibitory activity against COX-1 and COX-2, J Nat Prod. 68 (2005) 69-73.

- [35] P.T. Thuong, C.H. Lee, T.T. Dao, P.H. Nguyen, W.G. Kim, S. J. Lee, W.K. Oh, Triterpenoids from the leaves of *Diospyros kaki* (persimmon) and their inhibitory effects on protein tyrosine phosphatase 1B, J Nat Prod. 71 (2008) 1775-1778.
- [36] M. Litaudon, C. Jolly, C.L. Callonec, D. D. Cuong, P. Retailleau, O. Nosjean, V.H. Nguyen, B. Pfeiffer, J.A. Boutin, F. Guérinne, Cytotoxic pentacyclic triterpenoids from *Combretum sundaicum* and *Lantana camara* as inhibitors of Bcl-xL/BakBH3 domain peptide interaction, J Nat Prod. 72 (2009) 1314-1320.
- [37] C. Chen, S. Qiang, L. Lou, W. Zhao, Cucurbitane type triterpenoids from the stems of *Cucumis melo*, J Nat Prod. 72 (2009) 824-829.
- [38] P. Wang, S. Ownby, Z. Zhang, W. Yuan, S. Li, Cytotoxicity and inhibition of DNA topoisomerase I of polyhydroxylated triterpenoids and triterpenoid glycosides, Bioorg Med Chem Lett. 20 (2010) 2790-2796.
- [39] J.J. Ramírez-Espinosa, M.Y. Rios, S.L. Martínez, F.L. Vallejo, J.L. Medina-Franco, P. Paoli, G. Camici, G. Navarrete-Vázquez, R. Ortiz-Andrade, S. Estrada-Soto, Antidiabetic activity of some pentacyclic acid triterpenoids, role of PTP-1B: in vitro, in silico, and in vivo approaches, Eur J Med Chem. 46 (2011) 2243-2251.
- [40] J.D.S. Mpetga, Y. Shen, P. Tane, S.F. Li, H.P. He, H.K. Wabo, M. Tene, Y. Leng, X.J. Hao, Cycloartane and friedelane triterpenoids from the leaves of *Caloncoba glauca* and their evaluation for inhibition of 11 β - hydroxysteroid dehydrogenases, J Nat Prod. 75 (2012) 599-604.

- [41] L. Xiong, M. Zhu, C. Zhu, S. Lin, Y. Yang, J. Shi, Structure and bioassay of triterpenoids and steroids isolated from *Sinocalamus affinis*, *J Nat Prod.* 75 (2012) 1160-1166.
- [42] C.L. Zhang, Y. Wang, Y.F. Liu, G. Ni, D. Liang, H. Luo, X.Y. Song, W.Q. Zhang, R.Y. Chen, N.H. Chen, D.Q. Yu, Iridal-type triterpenoids with neuroprotective activities from *Iris tectorum*, *J Nat Prod.* 77 (2014) 411-415.
- [43] G.W. Wang, C. Lv, X. Fang, X.H. Tian, J. Ye, H.L. Li, L. Shan, Y.H. Shen, W.D. Zhang, Eight pairs of epimeric triterpenoids involving a characteristic spiro-E/F ring from *Abies faxoniana*, *J Nat Prod.* 78 (2015) 50-60.
- [44] L.P. Hammett, Some relations between reaction rates and equilibrium constants, *Chem Rev.* 17 (1935) 125-136.
- [45] L.P. Hammett, *Physical Organic Chemistry*, 2nd ed., McGraw-Hill, New York, 1970.
- [46] R.W. Taft Jr., Polar and steric substituent constants for aliphatic and o-benzoate groups from rates of esterification and hydrolysis of esters¹, *J Am Chem Soc.* 74 (1952) 3120-3128.
- [47] C. Hansch, P.P. Maloney, T. Fujita, R.M. Muir, Correlation of biological activity of phenoxyacetic acids with Hammett substituent constants and partition coefficients, *Nature.* 194 (1962) 178-180.
- [48] C. Hansch, T. Fujita, ρ - σ - π Analysis. A method for the correlation of biological activity and chemical structure, *J Am Chem Soc.* 86 (1964) 1616-1626.
- [49] C. Hansch, Quantitative approach to biochemical structure-activity relationships, *Acc Chem Res.* 2 (1969) 232-239.
- [50] S.M. Free Jr., J.W. Wilson, A mathematical contribution to structure-activity studies, *J Med Chem.* 7(1964) 395-399.

- [51] Z. Simon, Specific interactions. Intermolecular forces, steric requirements, and molecular size, *Angew Chem Int Ed Eng.* 13 (1974) 719-727.
- [52] L.H. Hall, L.B. Kier, Structure-activity studies using valence molecular connectivity, *J Pharm Sci.* 66 (1977) 642-644.
- [53] L.B. Kier, L.H. Hall, Molecular structure description: the electrotopological state, Academic Press, San Diego, CA, 1999.
- [54] W. Tong, D.R. Lowis, R. Perkins, Y. Chen, W.J. Welsh, D.W. Goddette, T.W. Heritage, D.M. Sleehan, Evaluation of quantitative structure-activity relationship methods for large-scale prediction of chemicals binding to the estrogen receptor, *J Chem Inf Comput Sci.* 38 (1998) 669-677.
- [55] S.J. Cho, W. Zheng, A. Tropsha, Focus-2D: a new approach to the design of targeted combinatorial chemical libraries, *Pac Symp Biocomput.* (1998) 305-316.
- [56] H. Gao, J. Bajorath, Comparison of binary and 2D QSAR analyses using inhibitors of human carbonic anhydrase II as a test case, *J Mol Diversity.* 4 (1998) 115-130.
- [57] H. Gao, C. Williams, P. Labute, and J. Bajorath, Binary quantitative structure-activity relationship (QSAR) analysis of estrogen receptor ligands, *J Chem Inf Comput Sci.* 39 (1999) 164-168.
- [58] R.D. Cramer III, D.E. Patterson, J.D. Bunce, Comparative molecular field analysis (CoMFA). 1. Effect of shape on binding of steroids to carrier proteins, *J Am Chem Soc.* 110 (1988) 5959- 5967.
- [59] H. Kubinyi, G. Folkers, Y.C. Martin (Eds.), 3D QSAR in drug design, ligand-protein interactions and molecular similarity (Vol. 2) and recent advances (Vol. 3), Kluwer Academic Publishers, the Netherlands, 1997.

- [60] G. Klebe, U. Abraham, T. Mietzner, Molecular similarity indices in a comparative analysis (CoMSIA) of drug molecules to correlate and predict their biological activity, *J Med Chem.* 37 (1994) 4130-4146.
- [61] D.D. Robinson, P.J. Winn, P.D. Lyne, W.G Richards, Self-organizing molecular field analysis: a tool for structure-activity studies, *J Med Chem.* 42 (1999) 573-583.
- [62] H.L. Jiang, K.X. Chen, H.W. Wang, Y. Tang, J.Z. Chen, R.Y. Ji, 3D-QSAR study on ether and ester analogs of artemisinin with comparative molecular field analysis, *Zhongguo Yao Li Xue Bao.* 15 (1994) 481-487.
- [63] J.Z. Chen, L.H. Hu, H.L. Jiang, J.D. Gu, W. L. Zhu, Z.L. Chen, K.X. Chen, R.Y. Ji, A 3D-QSAR study on ginkgolides and their analogues with comparative molecular field analysis, *Bioorg Med Chem Lett.* 8 (1998) 1291-1296.
- [64] J.R. Woolfrey, M.A. Avery, A.M. Doweyko, Comparison of 3D quantitative structure-activity relationship methods: analysis of the in vitro antimalarial activity of 154 artemisinin analogues by hypothetical active-site lattice and comparative molecular field analysis, *J Comput Aided Mol Des.* 12 (1998) 165-181.
- [65] M. Jalali-Heravi, M.H. Fatemi, Artificial neural network modeling of Kováts retention indices for noncyclic and monocyclic terpenes, *J Chromatogr A.* 915 (2001) 177-183.
- [66] J.A. Grodnitzky, J.R. Coats, QSAR evaluation of monoterpenoids' insecticidal activity, *J Agric Food Chem.* 50 (2002) 4576-4580.
- [67] F. Cheng, J. Shen, X. Luo, W. Zhu, J. Gu, R. Ji, H. Jiang, K. Chen, Molecular docking and 3-D-QSAR studies on the possible antimalarial mechanism of artemisinin analogues, *Bioorg Med Chem.* 10 (2002) 2883-2891.

- [68] T. Kuriyama, T.J. Schmidt, E. Okuyama, Y. Ozoe, Structure-activity relationships of seco-prezizaane terpenoids in gamma-aminobutyric acid receptors of houseflies and rats, *Bioorg Med Chem.* 10 (2002) 1873-1881.
- [69] M.A. Avery, M. Alvim-Gaston, C.R. Rodrigues, E.J. Barreiro, F.E. Cohen, Y.A. Sabnis, J.R. Woolfrey, Structure-activity relationships of the antimalarial agent artemisinin. 6. The development of predictive in vitro potency models using CoMFA and HQSAR methodologies, *J Med Chem.* 45 (2002) 292-303.
- [70] T. Ghafourian, P. Zandasrar, H. Hamishekar, A. Nokhodchi, The effect of penetration enhancers on drug delivery through skin: a QSAR study, *J Control Release.* 99 (2004) 113-125.
- [71] R. Guha, P.C. Jurs, Development of QSAR models to predict and interpret the biological activity of artemisinin analogues, *J Chem Inf Comput Sci.* 44 (2004) 1440-1449.
- [72] T.J. Schmidt, M. Gurrath, Y. Ozoe, Structure-activity relationships of seco-prezizaane and picrotoxane/picrodendrane terpenoids by Quasar receptor-surface modeling, *Bioorg Med Chem.* 12 (2004) 4159-4167.
- [73] F. Cortés-Selva, M. Campillo, C.P. Reyes, I.A. Jiménez, S. Castanys, I.L. Bazzocchi, L. Pardo, F. Gamarro, A.G. Ravelo, SAR studies of dihydro-betaagarofuran sesquiterpenes as inhibitors of the multidrug-resistance phenotype in a *Leishmania tropica* line overexpressing a P-glycoprotein-like transporter, *J Med Chem.* 47 (2004) 576-587.
- [74] B. Siedle, A.J. García-Piñeres, R. Murillo, J. Schulte-Mönting, V. Castro, P. Rüngeler, C.A. Klaas, F.B. Da Costa, W. Kisiel, I. Merfort, Quantitative structure-activity relationship of sesquiterpene lactones as inhibitors of the transcription factor NF-kappaB, *J Med Chem.* 47 (2004) 6042-6054.

- [75] F.A. Macías, R.F. Velasco, D. Castellano, J.C. Galindo, Application of Hansch's model to guaianolide ester derivatives: a quantitative structure-activity relationship study, *J Agric Food Chem.* 53 (2005) 3530-3539.
- [76] W. Zhu, G. Chen, L. Hu, X. Luo, C. Gui, C. Luo, C.M. Puah, K. Chen, J.H. Jiang, QSAR analyses on ginkgolides and their analogues using CoMFA, CoMSIA, and HQSAR, *Bioorg Med Chem.* 13 (2005) 313-322.
- [77] M. Ishihara, H. Sakagami, W.K. Liu, Quantitative structure-cytotoxicity relationship analysis of betulinic acid and its derivatives by semi-empirical molecular-orbital method, *Anticancer Res.* 25 (2005) 3951-3955.
- [78] S.L. Cunningham, A.R. Cunningham, B.W. Day, CoMFA, HQSAR and molecular docking studies of butitaxel analogues with beta-tubulin, *J Mol Model.* 11 (2005) 48-54.
- [79] H. Parra-Delgado, C.M. Compadre, T. Ramírez-Apan, M.J. Muñoz-Fambuena, R.L. Compadre, P. Ostrosky-Wegman, M. Martínez-Vázquez, Synthesis and comparative molecular field analysis (CoMFA) of argentatin B derivatives as growth inhibitors of human cancer cell lines, *Bioorg Med Chem.* 14 (2006) 1889-1901.
- [80] Y.H. Chung, S.C. Jang, S.J. Kim, N.D. Sung, 2D-QSAR and HQSAR on the inhibition activity of protein tyrosine phosphatase 1B with oleanolic acid analogues, *J Appl Biol Chem.* 50 (2007) 52-57.
- [81] B.F. Rasulev, A.I. Saidkhodzhaev, S.S. Nazrullaev, K.S. Akhmedkhodzhaeva, Z.A. Khushbaktova, J. Leszczynski, Molecular modelling and QSAR analysis of the estrogenic activity of terpenoids isolated from Ferula plants, *SAR QSAR Environ Res.* 18 (2007) 663-673.

- [82] B. Hemmateenejad, K. Javadnia, M. Elyasi, Quantitative structure-retention relationship for the Kovats retention indices of a large set of terpenes: a combined data splitting-feature selection strategy, *Anal Chim Acta.* 592 (2007) 72-81.
- [83] M.T. Scotti, M.B. Fernandes, M.J. Ferreira, V.P. Emerenciano, Quantitative structure-activity relationship of sesquiterpene lactones with cytotoxic activity, *Bioorg Med Chem.* 15 (2007) 2927-2934.
- [84] H.J. Chang, H.J. Kim, H.S. Chun, Quantitative structure-activity relationship (QSAR) for neuroprotective activity of terpenoids, *Life Sci.* 80 (2007) 835-841.
- [85] C.P. Reyes, F. Muñoz-Martínez, I.R. Torrecillas, C.R. Mendoza, F. Gamarro, I.L. Bazzocchi, M.J. Núñez, L. Pardo, S. Castanys, M. Campillo, I.A.E Jiménez, Biological evaluation, structure-activity relationships, and three-dimensional quantitative structure-activity relationship studies of dihydro-betaagarofuran sesquiterpenes as modulators of P-glycoprotein-dependent multidrug resistance, *J Med Chem.* 50 (2007) 4808-4817.
- [86] L. Kang, C.W. Yap, P.F. Lim, Y.Z. Chen, P.C. Ho, Y.W. Chan, G.P. Wong, S.Y. Chan, Formulation development of transdermal dosage forms: quantitative structure-activity relationship model for predicting activities of terpenes that enhance drug penetration through human skin, *J Control Release.* 120 (2007) 211-219.
- [87] W.N. Setzer, Non-intercalative triterpenoid inhibitors of topoisomerase II: A molecular docking study, *TOBCJ.* 1 (2008) 13-17.
- [88] R.P. Verma, C. Hansch, Taxane analogues against breast cancer: a quantitative structure-activity relationship study, *ChemMedChem.* 3 (2008) 642-652.
- [89] C. Hansch, R.P. Verma, Understanding tubulin/microtubule-taxane interactions: a quantitative structure-activity relationship study, *Mol Pharm.* 5 (2008) 151-161.

- [90] F.J. Cardoso, A.F. de Figueiredo, M. da Silva Lobato, R.M. de Miranda, R.C. de Almeida, J.C. Pinheiro, A study on antimalarial artemisinin derivatives using MEP maps and multivariate QSAR, *J Mol Model.* 14 (2008) 39-48.
- [91] Z. Wang, J. Song, Z. Han, Z. Jiang, W. Zheng, J. Chen, Z. Song, S. Shang, Quantitative structure-activity relationship of terpenoid aphid antifeedants, *J Agric Food Chem.* 56 (2008) 11361-11366.
- [92] W.N. Setzer, A theoretical investigation of cytotoxic activity of celastroid triterpenoids, *J Mol Model.* 15 (2009) 197-201.
- [93] R.P. Verma, C. Hansch, Taxane analogues against lung cancer: a quantitative structure-activity relationship study, *Chem Biol Drug Des.* 73 (2009) 627-636.
- [94] W.E. Steinmetz, C.B. Rodarte, A. Lin, 3D QSAR study of the toxicity of trichothecene mycotoxins, *Eur J Med Chem.* 44 (2009) 4485-4489.
- [95] T.J. Schmidt, A.M. Nour, S.A. Khalid, M. Kaiser, R. Brun, Quantitative structure-antiprotozoal activity relationships of sesquiterpene lactones, *Molecules.* 14 (2009) 2062-2076.
- [96] C. Hansch, R.P. Verma, Overcoming tumor drug resistance with C2-modified 10-deacetyl-7-propionyl cephalomannines: a QSAR study, *Mol Pharm.* 6 (2009) 849-860.
- [97] L. Saiz-Urra, J.C. Racero, A.J. Macías-Sánchez, R. Hernández-Galán, J.R. Hanson, M. Pérez-González, I.G. Collado, Synthesis and quantitative structure-antifungal activity relationships of clovane derivatives against *Botrytis cinerea*, *J Agric Food Chem.* 57 (2009) 2420-2428.
- [98] R.J. Little, A.A. Pestano, Z. Parra, Modeling of peroxide activation in artemisinin derivatives by serial docking, *J Mol Model.* 15 (2009) 847-858.

- [99] J. Xu, S. Huang, H. Luo, G. Li, J. Bao, S. Cai, Y. Wang, QSAR Studies on andrographolide derivatives as α -glucosidase inhibitors, *Int J Mol Sci.* 11 (2010) 880-895.
- [100] M.E. Badawy, S.A. El-Arami, S.A. Abdelgaleil, Acaricidal and quantitative structure activity relationship of monoterpenes against the two-spotted spider mite, *Tetranychus urticae*, *Exp Appl Acarol.* 52 (2010) 261-274.
- [101] R.P. Verma, C. Hansch, QSAR modeling of taxane analogues against colon cancer, *Eur J Med Chem.* 45 (2010) 1470-1477.
- [102] D.L. McGovern, P.D. Mosier, B.L. Roth, R.B. Westkaemper, CoMFA analyses of C-2 position salvinorin A analogs at the kappa-opioid receptor provides insights into epimer selectivity, *J Mol Graph Model.* 28 (2010) 612-625.
- [103] H.M. Hassan, A.Y. Elnagar, M.A. Khanfar, A.A. Sallam, R. Mohammed, L.A. Shaala, D.T. Youssef, M.S. Hifnawy, K.A. El Sayed, Design of semisynthetic analogues and 3D-QSAR study of eunicellin-based diterpenoids as prostate cancer migration and invasion inhibitors, *Eur J Med Chem.* 46 (2011) 1122-1130.
- [104] S.B. Bharate, I.P. Singh, Quantitative structure-activity relationship study of phloroglucinol-terpene adducts as anti-leishmanial agents, *Bioorg Med Chem Lett.* 21 (2011) 4310-4315.
- [105] P. Lan, W.N. Chen, P.H. Sun, W.M. Chen, 3D-QSAR studies on betulinic acid and betulin derivatives as anti-HIV-1 agents using CoMFA and CoMSIA, *Med Chem Res.* 20 (2011) 1247-1259.
- [106] N.S. Moorthy, M.J. Ramos, P.A. Fernandes, Prediction of the relationship between the structural features of andrographolide derivatives and α -glucosidase inhibitory activity: a

quantitative structure-activity relationship (QSAR) study, *J Enzyme Inhib Med Chem.* 26 (2011) 78-87.

- [107] P. Lan, W.N. Chen, Z.J. Huang, P.H. Sun, W.M. Chen, Understanding the structure-activity relationship of betulinic acid derivatives as anti-HIV-1 agents by using 3D-QSAR and docking, *J Mol Model.* 17 (2011) 1643-1659.
- [108] S.P. Wei, Z.Q. Ji, H.X. Zhang, J.W. Zhang, Y.H. Wang, W.J. Wu, Isolation, biological evaluation and 3D-QSAR studies of insecticidal/narcotic sesquiterpene polyol esters, *J Mol Model.* 17 (2011) 681-693.
- [109] M.L. Zhao, J.J. Yin, M.L. Li, Y. Xue, Y. Guo, QSAR study for cytotoxicity of diterpenoid tanshinones, *Interdiscip Sci.* 3 (2011) 121-127.
- [110] J. Bartalis, F.T. Halaweish, In vitro and QSAR studies of cucurbitacins on HepG2 and HSC-T6 liver cell lines, *Bioorg Med Chem.* 19 (2011) 2757-2766.
- [111] Z. Liang, L. Zhang, L. Li, J. Liu, H. Li, L. Zhang, L. Chen, K. Cheng, M. Zheng, X. Wen, P. Zhang, J. Hao, Y. Gong, X. Zhang, X. Zhu, J. Chen, H. Liu, H. Jiang, C. Luo, H. Sun, Identification of pentacyclic triterpenes derivatives as potent inhibitors against glycogen phosphorylase based on 3D-QSAR studies, *Eur J Med Chem.* 46 (2011) 2011-2021.
- [112] A. Maurya, F. Khan, D.U. Bawankule, D.K. Yadav, S.K. Srivastava, QSAR, docking and in vivo studies for immunomodulatory activity of isolated triterpenoids from *Eucalyptus tereticornis* and *Gentiana kurroo*, *Eur J Pharm Sci.* 47 (2012) 152-161.
- [113] F. Tong, J.R. Coats, Quantitative structure-activity relationships of monoterpenoid binding activities to the housefly GABA receptor, *Pest Manag Sci.* 68 (2012) 1122-1129.

- [114] K. Kalani, D.K. Yadav, F. Khan, S.K. Srivastava, N. Suri, Pharmacophore, QSAR, and ADME based semisynthesis and in vitro evaluation of ursolic acid analogs for anticancer activity, *J Mol Model.* 18 (2012) 3389-3413.
- [115] F. Abbasitabar, V. Zare-Shahabadi, Development predictive QSAR models for artemisinin analogues by various feature selection methods: a comparative study, *SAR QSAR Environ Res.* 23 (2012) 1-15.
- [116] W. Ding, M. Sun, S. Luo, T. Xu, Y. Cao, X. Yan, Y. Wang, A 3D QSAR study of betulinic acid derivatives as anti-tumor agents using topomer CoMFA: model building studies and experimental verification, *Molecules.* 18 (2013) 10228-10241.
- [117] J. Song, Z. Wang, A. Findlater, Z. Han, Z. Jiang, J. Chen, W. Zheng, S. Hyde, Terpenoid mosquito repellents: A combined DFT and QSAR study, *Bioorg Med Chem Lett.* 23 (2013) 1245-1248.
- [118] I.J. Sousa, M.J. Ferreira, J. Molnár, M.X. Fernandes, QSAR studies of macrocyclic diterpenes with P-glycoprotein inhibitory activity, *Eur J Pharm Sci.* 48 (2013) 542-553.
- [119] S. Rajkhowa, I. Hussain, K.K. Hazarika, P. Sarmah, R.C. Deka, Quantitative structure-activity relationships of the antimalarial agent artemisinin and some of its derivatives - a DFT approach, *Comb Chem High Throughput Screen.* 16 (2013) 590-602.
- [120] C. Schomburg, W. Schuehly, F.B. Da Costa, K.H. Klempnauer, T.J. Schmidt, Natural sesquiterpene lactones as inhibitors of Myb-dependent gene expression: structure-activity relationships, *Eur J Med Chem.* 63 (2013) 313-320.
- [121] S. Liao, J. Song, Z. Wang, J. Chen, G. Fan, Z. Song, S. Shang, S. Chen, P. Wang, Molecular interactions between terpenoid mosquito repellents and human-secreted attractants, *Bioorg Med Chem Lett.* 24 (2014) 773-779.

- [122] A.I. Foudah, A.A. Sallam, M.R. Akl, K.A. El Sayed, Optimization, pharmacophore modeling and 3D-QSAR studies of sipholanes as breast cancer migration and proliferation inhibitors, *Eur J Med Chem.* 73 (2014) 310-324.
- [123] J. Vinholesa, A. Rudnitskayab, P. Gonçalvesc, F. Martelc, M.A. Coimbraa, S.M. Rocha, Hepatoprotection of sesquiterpenoids: a quantitative structure–activity relationship (QSAR) approach, *Food Chem.* 146 (2014) 78-84.
- [124] H.G. Trossini, V.G. Maltarollo, T.J. Schmidt, Hologram QSAR studies of antiprotozoal activities of sesquiterpene lactones, *Molecules.* 19 (2014) 10546-10562.
- [125] J.L. Cheng, M. Zheng, T.T. Yao, X.L. Li, J.H. Zhao, M. Xia, G.N. Zhu, Synthesis, antifungal activity, and QSAR study of novel trichodermin derivatives, *J Asian Nat Prod.* 17 (2015) 47-55.
- [126] K. Muthusamy, P. Kirubakaran, G. Krishnasamy, R.R. Thanashankar, Computational insights into the inhibition of influenza viruses by rupestonic acid derivatives: pharmacophore modeling, 3D-QSAR, CoMFA and COMSIA studies, *Comb Chem High Throughput Screen.* 18 (2015) 63-74.
- [127] M. Appell M, W.B. Bosma, Assessment of the electronic structure and properties of trichothecene toxins using density functional theory, *J Hazard Mater.* 288 (2015) 113-123.
- [128] A. Hazra, C. Mondal, D. Chakraborty, A.K. Halder, Y.P. Bharitkar, S.K. Mondal, S. Banerjee, T. Jha, N.B. Mondal, Towards the development of anticancer drugs from andrographolide: semisynthesis, bioevaluation, QSARanalysis and pharmacokinetic studies, *Curr Top Med Chem.* 15 (2015) 1013-1026.

- [129] M.E. Saeed, O. Kadioglu, E.J. Seo, H.J. Greten, R. Brenk, T. Efferth, Quantitative structure-activity relationship and molecular docking of artemisinin derivatives to vascular endothelial growth factor receptor 1, *Anticancer Res.* 35 (2015) 1929-1934.
- [130] P. Tiwari, P. Sharma, F. Khan, N.S. Sangwan, B.N. Mishra, R.S. Sangwan, Structure Activity Relationship Studies of Gymnemic Acid Analogues for Antidiabetic Activity Targeting PPAR γ , *Curr Comput Aided Drug Des.* 11 (2015) 57-71.
- [131] R. Bonikowski, P. Świtakowska, M. Sienkiewicz, M. Zakłos-Szyda, Selected compounds structurally related to acyclic sesquiterpenoids and their antibacterial and cytotoxic activity, *Molecules.* 20 (2015) 11272-11296.
- [132] S. Andrade-Ochoa, G.V. Nevárez-Moorillón, L.E. Sánchez-Torres, M. Villanueva-García, B.E. Sánchez-Ramírez, L.M. Rodríguez-Valdez, B.E. Rivera-Chavira, Quantitative structure-activity relationship of molecules constituent of different essential oils with antimycobacterial activity against *Mycobacterium tuberculosis* and *Mycobacterium bovis*, *BMC Complement Altern Med.* 15 (2015) 332.

Chapter 3.

- [1] H. Wiener, Structural determination of paraffin boiling points, *J Am Chem Soc.* 69 (1947) 17-20.
- [2] F. Harary, Graph theory. Addison-Wesley, Reading MA, 1969.
- [3] L.B. Kier, L.H. Hall, Molecular connectivity in structure activity analysis, Research studies press Letchworth, Hertfordshire, U. K., 1986.
- [4] M. Randić, Characterization of molecular branching, *J Am Chem Soc.* 97 (1975) 6609-6615.

- [5] S.C. Basak, Use of molecular complexity indices in predictive pharmacology and toxicology: a QSAR approach, *Med Sci Res.* 15 (1987) 605-609.
- [6] C.E. Shannon, A mathematical theory of communication, *Bell Syst Tech J.* 27 (1948) 379-423.
- [7] S.C. Basak, S. Bertelsen, G.D. Grunwald, Application of graph theoretical parameters in quantifying molecular similarity and structure-activity studies, *J Chem Inf Comput Sci.* 34 (1994) 270-276.
- [8] S.C. Basak, G.D. Grunwald, Predicting mutagenicity of chemicals using topological and quantum chemical parameters: a similarity based study, *Chemosphere.* 31 (1995) 2529-2546.
- [9] R. Todeschini, V. Consonni, *Handbook of molecular descriptors*, Wiley, New York, 2008.
- [10] B. Lee, F.M. Richards, The interpretation of protein structures: estimation of static accessibility, *J Mol Biol.* 55 (1971) 379-400.
- [11] M.L. Connolly, Analytical molecular surface calculation, *J Appl Cryst.* 16 (1983) 548-558.
- [12] M.F. Sanner, Ph.D. dissertation thesis: Modeling and applications of molecular surfaces, Université de Haute-Alsace, France, 1992.
- [13] M.F. Sanner, A.J. Olson, Reduced surface: an efficient way to compute molecular surfaces, *Biopolymers.* 38 (1996) 305-320.
- [14] J. Kujawski, H. Popielarska, A. Myka, B. Drabińska, M.K. Bernard, The log P parameter as a molecular descriptor in the computer-aided drug design – an overview, *CMST.* 18 (2012) 81-88.

- [15] D. Eros, I. Kovács, L. Orfi, K. Takacs-Novak, G. Acsády, G. Kéri, Reliability of logP predictions based on calculated molecular descriptors: a critical review, *Curr Med Chem.* 9 (2002) 1819-1829.
- [16] M.J.S. Dewar, E.G. Zoebisch, E.F. Healy, J.J.P. Stewart, Development and use of quantum mechanical molecular models. 76. AM1: a new general purpose quantum mechanical molecular model, *J Am Chem Soc.* 107 (1985) 3902-3909.
- [17] J.J.P. Stewart, Optimization of parameters for semiempirical methods. 1. method, *J Comput Chem.* 10 (1989) 209-220.
- [18] J.J.P. Stewart, Comments on a comparison of AM1 with the recently developed PM3 method- reply, *J Comput Chem.* 11 (1990) 543-544.
- [19] J.J.P. Stewart, Optimization of parameters for semiempirical methods. III Extension of PM3 to Be, Mg, Zn, Ga, Ge, As, Se, Cd, In, Sn, Sb, Te, Hg, Tl, Pb, and Bi, *J Comput Chem.* 12 (1991) 320-341.
- [20] J.B. Foresman, A. Frisch, Exploring chemistry with electronic structure methods, Gaussian Inc., Pittsburgh, 1996.
- [21] W. Koch, M.C. Holthausen, A chemist's guide to density functional theory, Wiley-VCH, Weinheim, 2000.
- [22] C. Gruber, V. Buss, Quantum-mechanically calculated properties for the development of quantitative structure-activity relationships (QSAR'S). pKa-values of phenols and aromatic and aliphatic carboxylic acids, *Chemosphere.* 19 (1989) 1595-1609.
- [23] T. Sotomatsu, Y. Murata, T.J. Fujita, Correlation analysis of substituent effects on the acidity of benzoic acids by the AM1 method, *J Comput Chem.* 10 (1989) 94-98.

- [24] M. Karelson, V.S. Lobanov, A.R. Katritzky, Quantum-chemical descriptors in QSAR/QSPR studies, *Chem Rev.* 96 (1996) 1027-1043.
- [25] D.F. Lewis, C. Ioannides, D.V. Parke, Interaction of a series of nitriles with the alcohol-inducible isoform of P450: computer analysis of structure-activity relationships, *Xenobiotica.* 24 (1994) 401-408.
- [26] R.G. Parr, R.A. Donnelly, M. Levy, W.E. Palke, Electronegativity: the density functional viewpoint, *J Chem Phys.* 68 (1978) 3801-3807.
- [27] R.P. Iczkowski, J.L. Margrave, Electronegativity, *J Am Chem Soc.* 83 (1961) 3547-3551.
- [28] I.N Levine, Quantum chemistry, Pearson Education, Singapore, 2000.
- [29] I.N Levine, Physical chemistry, Mc-Graw Hill, New York, 2011.
- [30] B.J. McConkey, V. Sobolev, M. Edelman, The performance of current methods in ligand-protein docking, *Curr Sci.* 83 (2002) 845-855.
- [31] G.M. Morris, D.S. Goodsell, R.S. Halliday, R. Huey, W.E. Hart, R.K. Belew, A.J. Olson, Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function, *J Comput Chem.* 19 (1998) 1639-1662.
- [32] M.L. Verdonk, J.C. Cole, M.J. Hartshorn, C.W. Murray, R.D. Taylor, Improved protein-ligand docking using GOLD, *Proteins.* 52 (2003) 609-623.
- [33] T.J. Ewing, S. Makino, A.G. Skillman, I.D. Kuntz, DOCK 4.0: search strategies for automated molecular docking of flexible molecule databases, *J Comput Aided Mol Des.* 15 (2001) 411-428.
- [34] R.A. Friesner, J.L. Banks, R.B. Murphy, T.A. Halgren, J.J. Klicic, D.T. Mainz, M.P. Repasky, E.H. Knoll, M. Shelley, J.K. Perry, D.E. Shaw, P. Francis, P.S. Shenkin, Glide:

a new approach for rapid, accurate docking and scoring. 1 Method and assessment of docking accuracy, J Med Chem. 47 (2004) 1739-1749.

[35] R. Abagyan, M. Totrov, D. Kuznetsov, ICM-A new method for protein modeling and design: applications to docking and structure prediction from the distorted native conformation. J Comput Chem. 15 (1994) 488-506.

[36] M. Rarey, B. Kramer, T. Lengauer, G. Klebe, A fast flexible docking method using an incremental construction algorithm, J Mol Biol. 261 (1996) 470-489.

[37] S.J. Weiner, P.A. Kollman, D.A. Case, U.C. Singh, C. Ghio, G. Alagona, S. Profeta, P. Weiner, A new force field for molecular mechanical simulation of nucleic acids and proteins, J Am Chem Soc. 106 (1984) 765-784.

[38] R. Huey, G.M. Morris, A.J. Olson, D.S. Goodsell, A semiempirical free energy force field with charge-based desolvation, J Comput Chem. 28 (2007) 1145-1152.

[39] T.W. Anderson, An introduction to multivariate analysis, John Wiley, New York, 1958.

[40] A.M. Kshirsagar, Multivariate analysis, Marcel Dekker, New York, 1972.

[41] C.R. Rao, Advanced statistical inference and its applications, Wiley Eastern, New Delhi, 1973.

Chapter 4.

[1] M. Bollen, S. Keppens, W. Stalmans, Specific features of glycogen metabolism in the liver, Biochem J. 336 (1998) 19-31.

[2] H.G. Hers, The control of glycogen metabolism in the liver, Annu Rev Biochem. 45 (1976) 167-190.

- [3] W. Stalmans, The interaction of liver phosphorylase *a* with glucose and AMP, Eur J Biochem. 49 (1974) 415-427.
- [4] W. Pimenta, N. Nurjhan, P.A. Jansson, M. Stumvoll, J. Gerich, M. Korytkowski, Glycogen: its mode of formation and contribution to hepatic glucose output in postabsorptive humans, Diabetologia. 37 (1994) 697-702.
- [5] M.K. Hellerstein, R.A. Neese, P. Linfoot, M. Christiannsen, S. Turner, A. Letscher, Hepatic gluconeogenic fluxes and glycogen turnover during fasting in humans. A stable isotope study, J Clin Invest. 100 (1997) 1305-1319.
- [6] N.G. Oikonomakos, E.D. Chrysina, M N. Kosmopoulou, D.D. Leonidas, Crystal structure of rabbit muscle glycogen phosphorylase *a* in complex with a potential hypoglycaemic drug at 2.0 Å resolution, Biochim Biophys Acta. 1647 (2003) 325-332.
- [7] V.L. Rath, M. Ammirati, D.E. Danley, J.L. Ekstrom, E.M. Gibbs, T.R. Hynes, A.M. Mathiowitz, R.K. Mcpherson, T.V. Olson, J.L. Treadway, D.J. Hoover, Human liver glycogen phosphorylase inhibitors bind at a new allosteric site, Chem Biol. 7 (2000) 677-682.
- [8] R. Kurukulasuriya, J.T. Link, D.J. Madar, Z. Pei, S.J. Richards, J.J. Rohde, A.J. Souers, B.G. Szczepankiewicz, Potential drug targets and progress towards pharmacologic inhibition of hepatic glucose production, Curr Med Chem. 10 (2003) 123-153.
- [9] Z. Liang, L. Zhang, L. Li, J. Liu, H. Li, L. Zhang, L. Chen, K. Cheng, M. Zheng, X. Wen, P. Zhang, J. Hao, Y. Gong, X. Zhang, X. Zhu, J. Chen, H. Liu, H. Jiang, C. Luo, H. Sun,

Identification of pentacyclic triterpenes derivatives as potent inhibitors against glycogen phosphorylase based on 3D-QSAR studies, Eur J Med Chem. 46 (2011) 2011-2021.

- [10] X. Wen, H. Sun, J. Liu, K. Cheng, P. Zhang, L. Zhang, J. Hao, L. Zhang, P. Ni, S.E. Zographos, D.D. Leonidas, K.M. Alexacou, T. Gimisis, J.M. Hayes, N.G. Oikonomakos, Naturally occurring pentacyclic triterpenes as inhibitors of glycogen phosphorylase: synthesis, structure-activity relationships, and X-ray crystallographic studies, J Med Chem. 51 (2008) 3540-3554.
- [11] P. Zhu, Y. Bi, J. Xu, Z. Li, J. Liu, L. Zhang, W. Ye, X. Wu, Terpenoids. III: Synthesis and biological evaluation of 23-hydroxybetulinic acid derivatives as novel inhibitors of glycogen phosphorylase, Bioorg Med Chem Lett. 19 (2009) 6966–6969.
- [12] P. Lan, J. Wang, D.M. Zhang, C. Shu, H.H. Cao, P.H. Sun, X.M. Wu, W.C. Ye, W.M. Chen, Synthesis and antiproliferative evaluation of 23-hydroxybetulinic acid derivatives, Eur J Med Chem. 46 (2011) 2490-2502.
- [13] Y. Bi, J. Xu, F. Sun, X. Wu, W. Ye, Y. Sun, W. Huang, Synthesis and biological activity of 23-hydroxybetulinic acid C-28 ester derivatives as antitumor agent candidates, Molecules. 17 (2012) 8832-8841.
- [14] W. Ding, M. Sun, S. Luo, T. Xu, Y. Cao, X. Yan, Y. Wang, A 3D QSAR study of betulinic acid derivatives as anti-tumor agents using topomer CoMFA: model building studies and experimental verification, Molecules. 18 (2013) 10228-10241.
- [15] S. Fulda, K.M. Debatin, Betulinic acid induces apoptosis through a direct effect on mitochondria in neuroectodermal tumors, Med Pediatric Oncol. 35 (2000) 616–618.

- [16] Z.N. Ji, W.C. Ye, G.G. Liu, W.L. Wendy Hsiao, 23-Hydroxybetulinic acid-mediated apoptosis is accompanied by decreases in Bcl-2 expression and telomerase activity in HL-60 Cells, *Life Sci.* 72 (2002) 1-9.
- [17] M. Mandal, R. Kumar, Bcl-2 Modulates telomerase activity, *J Biol Chem.* 272 (1997) 14183-14187.
- [18] Y Zheng, F. Zhou, X. Wu, X. Wen, Y. Li, B. Yan, J. Zhang, G. Hao, W. Ye, G. Wang, 23-Hydroxybetulinic acid from *Pulsatilla chinensis* (Bunge) Regel synergizes the antitumor activities of doxorubicin in vitro and in vivo, *J Ethnopharmacol.* 128 (2010) 615–622.
- [19] ACD/ChemSketch version 12.01, Advanced Chemistry Development, Inc., Toronto, Ontario, 2009.
- [20] E. Eroglu, Some QSAR Studies for a Group of Sulfonamide Schiff Base as Carbonic Anhydrase CA II Inhibitors, *Int J Mol Sci.* 9 (2008) 181-197.
- [21] L.B. Kier, Use of molecular negentropy to encode structure governing biological activity, *J Pharm Sci.* 69 (1980) 807-810.
- [22] S.C. Basak, D.K. Harriss, V.R. Magnuson, Comparative study of lipophilicity versus topological molecular descriptors in biological correlations, *J Pharm Sci.* 73 (1984) 429-437.
- [23] H. Wiener, Structural determination of paraffin boiling points, *J Am Chem Soc.* 69 (1947) 17-20.

- [24] D. Playsic, S. Nikolic, N. Trinajstic, Z. Mihalic, On the Harary index for the characterization of chemical graphs, *J Math Chem.* 12 (1993) 235-250.
- [25] M. Randiac, On the characterization of molecular branching, *J Am Chem Soc.* 97 (1975) 6609-6615.
- [26] L.B. Kier, L.H. Hall, Molecular connectivity in structure-activity analysis, Research studies press: Letchworth, Hertfordshire, U K, 1986.
- [27] M.W. Schmidt, K.K. Baldridge, J.A. Boatz, S.T. Elbert, M.S. Gordon, J.H. Jensen, GAMESS Version= 24 Mar 2007 (R1) from Iowa State University, *J Comput Chem.* 14 (1993) 1347-1363.
- [28] G.M. Morris, R. Huey, W. Lindstrom, M.F. Sanner, R.K. Belew, D.S. Goodsell, A.J. Olson, AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility, *J Comput Chem.* 30 (2009) 2785-2791.
- [29] R. Huey, G.M. Morris, A.J. Olson, D.S. Goodsell, A semiempirical free energy force field with charge-based desolvation, *J Comput Chem.* 28 (2007) 1145-1152.
- [30] R. Huey, D.S. Goodsell, G.M. Morris, A.J. Olson, Grid-based hydrogen bond potentials with improved directionality, *Lett Drug Des Discov.* 1 (2004) 178-183.
- [31] R. Thomsen, M.H. Christensen, MolDock: a new technique for high-accuracy molecular docking, *J Med Chem.* 49 (2006) 3315-3321.

Chapter 5.

- [1] E.I. Solomon, J. Zhou, F. Neese, E.G. Pavel, New insights from spectroscopy into the structure/function relationships of lipoxygenases, *Chem Biol.* 4 (1997) 795-808.

- [2] H.W. Gardner, Biological roles and biochemistry of the lipoxygenase pathway, HortScience. 30 (1995) 197-205.
- [3] R. Wisastra, F.J. Dekker, Inflammation, cancer and oxidative lipoxygenase activity are intimately linked, Cancers. 6 (2014) 1500-1521.
- [4] L.A. Dailey, P. Imming, 12-Lipoxygenase: classification, possible therapeutic benefits from inhibition, and inhibitors, Curr Med Chem. 6 (1999) 389-398.
- [5] V.E. Steele, C.A. Holmes, E.T. Hawk, L. Kopelovich, R.A. Lubet, J.A. Crowell, C.C. Sigman, G.J. Kelloff, Lipoxygenase inhibitors as potential cancer chemopreventives, Cancer Epidemiol Biomarkers Prev. 8 (1999) 467-483.
- [6] B. Samuelsson, S.E. Dahlen, J.A. Lindgren, C.A. Rouzer, C.N. Serhan, Leukotrienes and lipoxins: structures, biosynthesis, and biological effects, Science. 237 (1987) 1171-1176.
- [7] X.Z. Ding, W.G. Tong, T.E. Adrian, 12-Lipoxygenase metabolite 12(S)-HETE stimulates human pancreatic cancer cell proliferation via protein tyrosine phosphorylation and ERK activation, Int J Cancer. 94 (2001) 630-636.
- [8] J.A. Cornicelli, B.K. Trivedi, 15-Lipoxygenase and its inhibition: a novel therapeutic target for vascular disease, Curr Pharm Des. 5 (1999) 11-20.
- [9] U.P. Kelavkar, J.B. Nixon, C. Cohen, D. Dillehay, T.E. Eling, K.F. Badr, Overexpression of 15-lipoxygenase-1 in PC-3 human prostate cancer cells increases tumorigenesis, Carcinogenesis. 22 (2001) 1765-1773.
- [10] C. Kemal, P. Louis-Flamberg, R. Krupinski-Olsen, A.L. Shorter, Reductive inactivation of soybean lipoxygenase-1 by catechols: a possible mechanism for regulation of lipoxygenase activity, Biochemistry. 26 (1987) 7064-7072.

- [11] R. Mogul, E. Johansen, T.R. Holman, Oleyl sulfate reveals allosteric inhibition of soybean lipoxygenase-1 and human 15-lipoxygenase, *Biochemistry*. 39 (2000) 4801-4807.
- [12] T. Amagata, S. Whitman, T.A. Johnson, C.C. Stessman, C.P. Loo, E.L. Jonclardy, P. Crews, T.R. Holman, Exploring sponge-derived terpenoids for their potency and selectivity against 12-human, 15-human, and 15-soybean lipoxygenases, *J Nat Prod.* 66 (2003) 230-235
- [13] A.A. Granovsky, Firefly version 8, [www](http://classic.chem.msu.su/gran/firefly/index.html)
<http://classic.chem.msu.su/gran/firefly/index.html>.
- [14] Y.H. Jang, W.A. Goddard III, K.T. Noyes, L.C. Sowers, S. Hwang, D.S. Chung, pKa Values of guanine in water: density functional theory calculations combined with poisson-boltzmann continuum-solvation model, *J Phys Chem B.* 107 (2003) 344-357.
- [15] J.E. Bartmess, Thermodynamics of the electron and the proton, *J Phys Chem.* 98 (1994) 6420-6424.
- [16] M.D. Liptak, K.G. Gross, P.G. Seybold, S. Feldgus, G.C. Shields, Absolute pK(a) determinations for substituted phenols, *J Am Chem Soc.* 124 (2002) 6421-6427.
- [17] P. Winget, E.J. Weber, C.J. Cramer, D.G. Truhlar, Computational electrochemistry: aqueous one-electron oxidation potentials for substituted anilines, *Phys Chem Chem Phys.* 2 (2000) 1231-1239.
- [18] N.S. Babu, Computational studies for oxidation reduction reactions of cinnoline - 4(1H)-one, in aqueous phase by density functional theory, *BJAST.* 4(2014) 465-476.
- [19] M.J. Nelson, D.G. Batt, J.S. Thompson, S.W. Wright, Reduction of the active-site iron by potent inhibitors of lipoxygenases, *J Biol Chem.* 266 (1991) 8225-8229.

Chapter 6.

- [1] H. Harada, Y. Kamei, Dose-dependent selective cytotoxicity of extracts from marine green alga, *Cladophoropsis vaucheriaeformis*, against mouse leukemia L1210 cells, *Biol Pharm Bull.* 21 (1998) 386-389.
- [2] K.E. Apt, P.W. Behrens, Commercial developments in microalgal biotechnology, *J Phycol.* 35 (1999) 215-26.
- [3] K. Harada, M. Suomalainen, H. Uchida, H. Masui, K. Ohmura, J. Kiviranta, M.L. Niku-Paavola, T. Ikemoto, Insecticidal compounds against mosquito larvae from *Oscillatoria agardhii* strain 27, *Environ Toxicol.* 15 (2000) 114-119.
- [4] V.H. Argandoña, J. Rovirosa, A. San-Martín, A. Riquelme, A.R. Díaz-Marrero, M. Cueto, J. Darias, O. Santana, A. Guadaño, A. González-Coloma, Antifeedant effects of marine halogenated monoterpenes, *J Agric Food Chem.* 50 (2002) 7029-7033.
- [5] C.D. Inés, V.H. Argandoña, J. Rovirosa, A. San-Martín, A.R. Díaz-Marrero, M. Cueto, A.G. Coloma, Cytotoxic activity of halogenated monoterpenes from *Plocamium cartilagineum*, *Z Naturforsch.* 59c (2004) 339-344.
- [6] ACD/ChemSketch version 12.01, Advanced Chemistry Development, Inc., Toronto, Ontario, 2009.
- [7] A.A. Granovsky, Firefly version 8, www <http://classic.chem.msu.su/gran/firefly/index.html>.

Chapter 7.

- [1] D. Barford, A.J. Flint, N.K. Tonks, Crystal structure of human protein tyrosine phosphatase 1B, *Science*. 263 (1994) 1397-1404.
- [2] Z.Y. Zhang, Structure, mechanism, and specificity of protein-tyrosine phosphatases, *Curr Top Cell Regul.* 35 (1997) 21-68.
- [3] D. Barford, A.K. Das, M.P. Egloff, The structure and mechanism of protein phosphatases: insights into catalysis and regulation, *Annu Rev Biophys Biomol Struct.* 27 (1998) 133-164.
- [4] Z.Y. Zhang, Protein-tyrosine phosphatases: biological function, structural characteristics, and mechanism of catalysis, *Crit Rev Biochem Mol Biol.* 33 (1998) 1-52.
- [5] A. Alonso, J. Sasin, N. Bottini, I. Friedberg, I. Friedberg, A. Osterman, A. Godzik, T. Hunter, J. Dixon, T. Mustelin, Protein tyrosine phosphatases in the human genome, *Cell*. 117 (2004) 699-711.
- [6] K.A. Kenner, E. Anyanwu, J.M. Olefsky, J. Kusari, Protein-tyrosine phosphatase 1B is a negative regulator of insulin- and insulin-like growth factor-I-stimulated signaling, *J Biol Chem.* 271 (1996) 19810-19816.
- [7] J.R. Wiener, B.J. Kerns, E.L. Harvey, M.R. Conaway, J.D. Igglehart, A. Berchuck, R.C. Bast Jr, Overexpression of the protein tyrosine phosphatase PTP1B in human breast cancer: association with p185c-erbB-2 protein expression, *J Natl Cancer Inst.* 86 (1994) 372-378.

- [8] M. Elchebly, P. Payette, E. Michaliszyn, W. Cromlish, S. Collins, A.L. Loy, D. Normandin, A. Cheng, J. Himms-Hagen, C.C. Chan, C. Ramachandran, M.J. Gresser, M.L. Tremblay, B.P. Kennedy, Increased insulin sensitivity and obesity resistance in mice lacking the protein tyrosine phosphatase-1B gene, *Science*. 283 (1999) 1544-1548.
- [9] J. Liu, Pharmacology of oleanolic acid and ursolic acid, *J Ethnopharmacol*. 49 (1995) 57-68.
- [10] J. Liu, Oleanolic acid and ursolic acid: research perspective, *J Ethnopharmacol*. 100 (2005) 92-94.
- [11] P. Dzubak, M. Hajduch, D. Vydra, A. Hustova, M. Kyasnica, D. Biedermann, L. Markova, M. Urban, J. Sarek, Pharmacological activities of natural triterpenoids and their therapeutic implications, *Nat Prod Rep*. 23 (2006) 394-411.
- [12] R. Martín, J. Carvalho-Tavares, M. Hernández, M. Arnés, V. Ruiz-Gutiérrez, M.L. Nieto, Beneficial actions of oleanolic acid in an experimental model of multiple sclerosis: a potential therapeutic role, *Biochem Pharmacol*. 79 (2010) 198-208.
- [13] J.J. Ramírez-Espinosa, M.Y. Rios, S.L. Martínez, F.L. Vallejo, J.L. Medina-Franco, P. Paoli, G. Camici, G. Navarrete-Vázquez, R. Ortiz-Andrade, S. Estrada-Soto, Antidiabetic activity of some pentacyclic acid triterpenoids, role of PTP-1B: in vitro, in silico, and in vivo approaches, *Eur J Med Chem*. 46 (2011) 2243-2251.
- [14] Y.A. Puius, Y.U. Zhao, M. Sullivan, D.S. Lawrence, S.C. Almo, Z.Y. Zhang, Identification of a second aryl phosphate-binding site in protein-tyrosine phosphatase 1B: a paradigm for inhibitor design, *Proc Natl Acad Sci USA*. 94 (1997) 13420-13425.

[15] J.M. Castellano, A. Guinda, T. Delgado, M. Rada, J.A. Cayuela, Biochemical basis of the antidiabetic activity of oleanolic acid and related pentacyclic triterpenes, *Diabetes*. 62 (2013) 1791-1799.

[16] Y.N. Zhang, W. Zhang, D. Hong, L. Shi, Q. Shen, J.Y. Li, J. Li, L.H. Hu, Oleanolic acid and its derivatives: new inhibitor of protein tyrosine phosphatase 1B with cellular activities, *Bioorg Med Chem*. 16 (2008) 8697-8705.

[17] ACD/ChemSketch version 12.01, Advanced Chemistry Development, Inc., Toronto, Ontario, 2009.

[18] G.M. Morris, D.S. Goodsell, R.S. Halliday, R. Huey, W.E. Hart, R.K. Belew, A.J. Olson, Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function, *J Comput Chem*. 19 (1998) 1639-1662.

[19] Molegro molecular viewer – version 2.5.0. <http://www.molegro.com/index.php>.

[20] A.A. Granovsky, Firefly version 8, [www](http://classic.chem.msu.su/gran/firefly/index.html)
<http://classic.chem.msu.su/gran/firefly/index.html>.

[21] M.F. Sanner, Python: a programming language for software integration and development, *J Mol Graphics Mod*. 17 (1999) 57-61.

Chapter 8.

[1] R.W. Fuller, J.H. Cardellina II, Y. Kato, L.S. Brinen, J. Clardy, K.M. Snader, M.R. Boyd, A pentahalogenated monoterpene from the red alga Portieria hornemannii produces a novel cytotoxicity profile against a diverse panel of human tumor cell lines, *J Med Chem*. 35 (1992) 3007-3011.

- [2] R.W. Fuller, J.H. Cardellina II, J. Jurek, P.J. Scheuer, B.A. Lindner, M. McGuire, G.N. Gray, J.R. Steiner, J. Clardy, E. Menez, R.H. Shoemaker, D.J. Newman, K.M. Snader, M.R. Boyd, Isolation and structure/activity features of halomon - related antitumor monoterpenes from the red alga Portieria hornemannii, *J Med Chem.* 37 (1994) 4407-4411.
- [3] M.J. Egorin, D.L. Sentz, D.M. Rosen, M.F. Ballesteros, C.M. Kearns, P.S. Callery, J.L. Eiseman, Plasma pharmacokinetics, bioavailability, and tissue distribution in CD₂F₁ mice of halomon, an antitumor halogenated monoterpene isolated from the red alga Portieria hornemannii, *Cancer Chemother Pharmacol.* 39 (1996) 51-60.
- [4] M.J. Egorin, M.D. Rosen, S.E. Benjamin, P.S. Callery, D.L. Sentz, J.L. Eiseman, In vitro metabolism of mouse and human liver preparations of halomon, an antitumor halogenated monoterpene, *Cancer Chemother Pharmacol.* 41 (1997) 9-14.
- [5] M.E. Jung, M.H. Parker, Synthesis of several naturally occurring polyhalogenated monoterpene of the halomon class, *J Org Chem.* 62 (1997) 7094-7095.
- [6] T. Sotokawa, T. Noda, S. Pi, M.A. Hirama, A three step synthesis of halomon, *Angew Chem Int Ed Engl.* 39 (2000) 3430-3432.
- [7] H.D. Yoo, S.O. Ketchum, D. France, K. Bair, W.H. Gerwick, Vidalenolone , a novel phenolic metabolite from the tropical red alga Vidalia sp., *J Nat Prod.* 65 (2002) 51-53.
- [8] E.H. Andrianasolo, D. France, S.C. Kennon, W.H. Gerwick, DNA Methyl Transferase Inhibiting Halogenated Monoterpenes from the Madagascar Red Marine Alga Portieria hornemannii, *J Nat Prod.* 69 (2006) 576-579.

[9] J.J.P. Stewart, MOPAC2016, Stewart Computational Chemistry, Colorado Springs, CO, USA, HTTP://OpenMOPAC.net (2016).

[10] A.A. Granovsky, Firefly version 8, www
<http://classic.chem.msu.su/gran/firefly/index.html>.

[11] A.R. Allouche, Gabedit-a graphical user interface for computational chemistry softwares, J Comput Chem. 32 (2011) 174-182.

Chapter 9.

[1] M.R. Orofino, O. Kreuger, S. Grootjans, M.W. Biavatti, P. Vandenabeele, K. D'Herde, Sesquiterpene lactones as drugs with multiple targets in cancer treatment: focus on parthenolide, Anticancer Drugs. 23 (2012) 883-896.

[2] Y.M. Zhao, M.L. Zhang, Q.W. Shi, H. Kiyota, Chemical constituents of plants from the genus Inula, Chem Biodivers. 3 (2006) 371-384.

[3] X. Cheng, Q. Zeng, J. Ren, J. Qin, S. Zhang, Y. Shen, J. Zhu, F. Zhang, R. Chang, Y. Zhu, W. Zhang, H. Jin, Sesquiterpene lactones from Inula falconeri, a plant endemic to the Himalayas, as potential anti-inflammatory agents, Eur J Med Chem. 46 (2011) 5408-5415.

[4] ACD/ChemSketch version 12.01, Advanced Chemistry Development, Inc., Toronto, Ontario, 2009.

[5] B. Bagchi, A. Chatterjee, P. Ghosh, A.K. Bothra, A theoretical investigation of cytotoxic activity of halogenated monoterpenoids from plocamium cartilagineum, JOCPR. 4 (2012) 5076-5080.

[6] A.A. Granovsky, Firefly version 8, www
<http://classic.chem.msu.su/gran/firefly/index.html>.

- [7] E. Eroglu, Some QSAR studies for a group of sulfonamide schiff base as carbonic anhydrase CA II inhibitors, *Int J mol Sci.* 9 (2008) 181-197.
- [8] L.B. Kier, Use of molecular negentropy to encode structure governing biological activity, *J Pharm Sci.* 69 (1980) 807-810.
- [9] S.C. Basak, D.K. Harriss, V.R. Magnuson, Comparative study of lipophilicity versus topological molecular descriptors in biological correlations, *J Pharm Sci.* 73 (1984) 429-437.
- [10] J.J.P. Stewart, MOPAC2012, Stewart Computational Chemistry, Colorado Springs, CO, USA.
- [11] E. Eroğlu, H. Türkmen, S. Güler, S. Palaz, O. Oltulu, A DFT-based QSARs study of acetazolamide/sulfanilamide derivatives with carbonic anhydrase (CA-II) isozyme inhibitory activity, *Int J Mol Sci.* 8 (2007) 145-155.
- [12] S. Sharma, B. Bagchi, S. Mukhopadhyay, A.K. Bothra, Theoretical study of lysophosphatidic acid acyltransferase 2 inhibitors, *JOCPR.* 5 (2013) 348-355.
- [13] B. Das, P. Ghosh, A.K. Bothra, QSAR studies of anthrax lethal inhibitors through quantum chemical indices, *JOCPR.* 3 (2011) 443-449.