

LIST OF FIGURES

CHAPTERS	FIGURES	PAGE NO.
Chapter-IV	<p>Figure 1: Transfer volume (Δw_v^0) of amino acids varying with increasing temperatures (T/K) with increasing concentrations of 18C6 (0.001, 0.002 & 0.003 mass fraction)</p> <p>Figure 2. Plot of limiting molar refraction (R_M^0) for glycine, L-alanine and L-isoleucine in different mass fractions (w_1) of aq. 18C6 at 298.15 K</p>	162
Chapter-V	<p>Figure 1. Plot of surface tension of ionic liquid corresponding to the added conc. of aq. α-cyclodextrin</p> <p>Figure 2. Job plot of IL-cyclodextrin systems at λ_{\max} (nm) = 210 and at 298.15 K</p> <p>Figure 3. Benesi-Hildebrand double reciprocal plot for the effect of α-CD on the absorbance of IL at 210 nm at 298.15 K</p> <p>Figure 4. Plot of conductance of ionic liquid corresponding to the added conc. of aq. α-cyclodextrin</p> <p>Figure 5. ^1H NMR spectra of α-CD and 1:1 molar ratio of α-CD & IL in D_2O at 298.15K</p>	175-177
Chapter-VI	<p>Figure 1: Plot of φv^0 as a function of different mass fraction (w_1) of aqueous uric acid solutions, () $T=298.15\text{ K}$; () $T=303.15\text{ K}$ and () $T=308.15\text{ K}$</p> <p>Figure 2: Plot of φv^0 as a function of temperature (T/K) of different mass fraction (w_1) of aqueous uric acid solutions, () $w_1=0.00001$; () $w_1=0.00002$ and () $w_1=0.00003$</p>	199-201

List of Figures

	<p>Figure 3: Plot of B as a function of different mass fraction (w_1) of aqueous uric acid solutions, () $T=298.15\text{ K}$; () $T=303.15\text{ K}$ and () $T=308.15\text{ K}$</p> <p>Figure 4: Plot of B as a function of temperature (T/K) of different mass fraction (w_1) of aqueous uric acid solutions, () $w_1=0.00001$; () $w_1=0.00002$ and () $w_1=0.00003$</p> <p>Figure 5: Plot of B against ϕ_{V^0} of different mass fraction (w_1) of aqueous uric acid solutions, () $w_1=0.00001$; () $w_1=0.00002$ and () $w_1=0.00003$</p> <p>Figure 6: Plot of molar refractive index (R_M) against square root of concentration (\sqrt{m}) for citric acid in mass fraction (w_1) of aqueous uric acid solution, () $w_1=0.00001$; () $w_1=0.00002$ and () $w_1=0.00003$</p>	
Chapter-VII	<p>Figure 1. Variation of conductivity of aqueous (a) 15-C-5 and (b) 18-C-6 with mole ratio of CE/IL at 298.15 K</p> <p>Figure 2. Variation of surface tension of aqueous (a) 15-C-5 and (b) 18-C-6 with mole ratio of CE/IL at 298.15 K</p> <p>Figure 3. ^1H NMR Spectra of 15-C-5, [EMIm]Ts and 1:1 molar ratio of 15-C-5 + [EMIm]Ts in D_2O in 298.15K</p> <p>Figure 4. ^1H NMR Spectra of 18-C-6, [EMIm]Ts and 1:1 molar ratio of 18-C-6 + [EMIm]Ts in D_2O in 298.15K</p>	217-218
Chapter-VIII	<p>Figure 1. Plot of limiting molar volume (ϕ_{V^0}) vs mass fraction for L-Serine (♦), L-Proline (▲), L-Histidine (●) in aq. [bmyrr]PF₆ respectively</p> <p>Figure 2. Plot of viscosity B-coefficient vs mass fraction for L-Serine (♦), L-Proline (▲), L-Histidine (●) in aq. [bmyrr]PF₆ respectively</p> <p>Figure 3. Plot of viscosity B-coefficient vs limiting molar volume (ϕ_{V^0}) for L-Serine (♦), L-Proline (▲), L-Histidine (●)</p>	236-237

	<p>in aq. [bmpyrr]PF₆ respectively</p> <p>Figure 4. Plot of limiting molar refraction vs mass fraction for L-Serine (♦), L-Proline (○), L-Histidine (●) in aq. [bmpyrr]PF₆ respectively</p>	
Chapter-IX	<p>Figure 1. Plot of molar conductance (Λ) versus \sqrt{C} for [BTMA]Cl in 1-Hexanol at 298.15 K</p> <p>Figure 2. Plot of molar conductance (Λ) versus \sqrt{C} for [BTMA]Cl in 1-Heptanol at 298.15 K</p> <p>Figure 3. Plot of molar conductance (Λ) versus \sqrt{C} for [BTMA]Cl in 1-Octanol at 298.15K</p> <p>Figure 4: Plot of φV^ρ as a function of temperature (T/K) in different solvent systems</p> <p>Figure 5: Plot of B as a function of temperature (T/K) in different solvent systems</p> <p>Figure 6: Plot of molar refractive index (R_M) against square root of concentration (\sqrt{m}) for [BTMA]Cl in different solvent systems, (○) 1-Hexanol; (○) 1-Heptanol and (○) 1-Octanol</p>	261-263