

## List of Tables

- Table1:** Serological and molecular techniques for detection of plant pathogens
- Table2:** Different immunological formats used for detection of plant pathogenic fungi
- Table3:** *In situ* analysis of phytopathogenic fungi using immunofluorescence
- Table4:** Mycelial growth of *P. disseminata* isolates in different solid media
- Table5:** Mycelial growth of *P. disseminata* isolates in Richard's solution
- Table6:** Mycelial growth of *C. gloeosporioides* isolates in different solid media
- Table7:** Mycelial growth of *C. gloeosporioides* isolates in Richard's solution
- Table8:** Spore characters of *C. gloeosporioides* isolates
- Table9:** Pathogenicity test of *P. disseminata* isolate on *Persea bombycina* following detached leaf inoculation technique
- Table10:** Pathogenicity test of *C. gloeosporioides* on *Persea bombycina* following detached leaf inoculation technique
- Table11:** Grey blight and leaf blight disease incidence of som following whole plant inoculation with *P. disseminata* and *C. gloeosporioides*
- Table12:** Protein content of *C. gloeosporioides* isolates
- Table13:** Protein content of *P. disseminata* isolates
- Table14:** Indirect ELISA values ( $A_{405}$ ) of mycelial antigens reacted with PABs of *C. gloeosporioides* and *P. disseminata*
- Table15:** Dot immunobinding assay of mycelial antigens reacted with PABs of *C. gloeosporioides* and *P. disseminata*
- Table16:** Indirect ELISA values ( $A_{405}$ ) of som leaf antigens reacted with PABs of *C. gloeosporioides*
- Table17:** Indirect ELISA values ( $A_{405}$ ) of som leaf antigens reacted with PABs of *P. disseminata*
- Table18:** Detection of pathogen in naturally leaf blight infected som leaves using PTA-ELISA formats
- Table19:** Detection of pathogen in naturally grey blight infected som leaves using PTA-ELISA formats

- Table 20:** Dot immunobinding assay of healthy and leaf blight infected leaf antigens of som plants using PAb of *C. gloeosporioides*
- Table21:** Dot immunobinding assay of healthy and grey blight infected leaf antigens of som plants using PAb of *P. disseminata*
- Table22:** PTA-ELISA values ( $A_{405}$ ) showing reaction of PABs of *C.gloeosporioides* and *P.disseminata* with antigens of healthy and inoculated som leaves
- Table23:** Detection of foliar fungal pathogens in artificially inoculated Som leaves using Dot immunobinding assay
- Table24:** Microscopic characters of AMF spores associated with som root
- Table25:** AMF spore population in field grown Som plants
- Table26:** GenBank accession numbers and geographic location of ex-type strains of *Colletotrichum* species that showed homology with isolate SOM/CL/02
- Table27:** GenBank accession numbers and geographic location of ex-type strains of *Pestalotiopsis* species that showed homology with isolate IPL/SOM/P/01
- Table28:** Total number of polymorphic bands produced by different RAPD primers
- Table29:** Total and Ortho-phenol content in healthy and infected leaves
- Table30:** Effect of leaf diffusate of som morphotypes on spore germination and appressoria formation of *C. gloeosporioides*
- Table31:** Retention time and peak heights of phenolics from healthy and infected leaf sample of S7 morphotype analyzed by HPLC
- Table32:** *In vitro* antagonistic test of selected PGPR isolates against foliar fungal pathogens
- Table33:** *In vitro* antagonistic test of selected PGPF isolates against foliar fungal pathogens
- Table34:** Growth promotion of pot grown som plants following treatment with PGPR and AMF
- Table 35:** Protein and phenol content in som plants following treatment with PGPR and AMF
- Table36:** Growth promotion in som plants following treatment with Vermicompost, PGPR and AMF, singly and jointly

**Table 37:** Growth promotion in different morphotypes of som plants following application of PGPR, AMF and PGPF

**Table 38:** Activity of defense enzymes in som plants following treatment with bioinoculants and artificial inoculation with *P.disseminata*

**Table39:** Growth promotion of field grown som plants following application of bioformulation

**Table 40:** Effect of joint application of AMF, PGPF and PGPR on biochemical components and defense enzymes of som plants

**Table 41:** Percent lesion production in som plants after joint application with AMF, PGPF and PGPR following detached leaf inoculation with *C. gloeosporioides*