

## List of Figures

- Figure 1:** Maintenance of *Persea bombycina* morphotypes in glass house condition
- Figure 2:** Maintenance of *Persea bombycina* morphotypes in field condition
- Figure 3:** Leaf blight disease incidence of nursery grown (A,C,E) and field grown (B,D,F) som plants. Healthy leaf (B), Naturally infected leaf (A,C,D,E,F); Morphotype: (A) – S3, (B, D, F) – S6, (C) – S5, (E) – S4
- Figure 4:** Grey blight disease incidence in field grown som plants. Healthy leaf (A), naturally infected (B-D); Morphotype: (A) – S3, (B) – S5,(C) – S6 and (D) – S7
- Figure 5:** Percent Disease Index (PDI) of leaf blight and grey blight disease in nursery grown som plants
- Figure 6:** Percentage sporulation of *P. disseminata* isolates in three different solid media (PDA – Potato Dextrose Agar, RM – Richard’s Medium, OMA – Oat Meal Agar)
- Figure 7:** Scanning electron microscopic observations of *P. disseminata* spores (isolate IPL/SOM/P/01)
- Figure 8:** Growth of *P. disseminata* isolates in PDA media (A-C) and spore characteristics (D-F); (A,D) – IPL/SOM/P/01; (B,E) – IPL/SOM/P/02; (C,F) – IPL/SOM/P/03
- Figure 9:** Percentage sporulation of *C. gloeosporioides* isolates in three different solid media (PDA – Potato Dextrose Agar, RM – Richard’s Medium, OMA – Oat Meal Agar)
- Figure 10:** Growth of *C. gloeosporioides* isolates in PDA media (A-C) and spore characteristics (D-F); (A,D) – SOM/CI/01; (B,E) – SOM/CI/02; (C,F) – SOM/CI/03
- Figure 11:** Pathogenicity test of *P. disseminata* following detached leaf inoculation technique. (A) – Experimental setup after 48h of inoculation, (B,C) – Leaves of S5 morphotype showing disease symptoms after 96h of inoculation, (D,E) – Leaves of S6 morphotype showing disease symptoms after 96h of inoculation
- Figure 12:** Pathogenicity test of *C. gloeosporioides* following detached leaf inoculation technique. (A) – Experimental setup after 48h of inoculation, (B,C) – Leaves of S5 morphotype showing disease symptoms after 96h of inoculation, (D,E) – Leaves of S6 morphotype showing disease symptoms after 96h of inoculation
- Figure 13:** Screening of resistance of eight morphotypes of som plants against (A) *P. disseminata* and (B) *C. gloeosporioides* following whole plant inoculation

- Figure 14:** Optimization of ELISA by considering two variable, (A) dilution of the antiserum and (B) dilution of the antigen extract
- Figure 15:** Serological assay of *Pestalotiopsis disseminata*; (A) - Dot blot; (B) - Immunofluorescence of young mycelia of *P. disseminata* reacted with PAb-Pt and labelled with FITC; C- Western blot analysis of mycelial antigen of *P. disseminata* probed with PAb-Pt on nitrocellulose membrane using NBT/BCIP as substrate
- Figure 16:** Spore morphology of *P. disseminata*. (A) – Spore under bright field, (B) – Scanning electron micrograph of spores, (C) – Indirect immunofluorescence of spores of *P. disseminata* treated with PAb-Pt and labelled with FITC
- Figure 17:** Serological assay of *Colletotrichum gloeosporioides*; (A) – Dot blot; (B) - Immunofluorescence of young mycelia of *C.gloeosporioides* reacted with PAb-Cg labelled with FITC; C - Western blot analysis of mycelial antigen of *C. gloeosporioides* probed with PAb-Cg on nitrocellulose membrane using NBT/BCIP as substrate
- Figure 18:** Spore morphology of *C. gloeosporioides*; (A) – Spore characteristics under bright field microscope, (B) – Indirect immunofluorescence of spores of *C.gloeosporioides* probed with PAb-Cg and labelled with FITC
- Figure 19:** Cellular localization of CRA shared by som plant and *C. gloeosporioides*. (A) Autofluorescence; (B) Cellular localization of CRA in som leaf tissue treated with PAb-Cg and labelled with FITC
- Figure 20:** Cellular localization of CRA shared by som plant and *P. disseminata* (A) Autofluorescence; (B) Cellular localization of CRA in som leaf tissue treated with PAb-Pt and labelled with FITC
- Figure 21:** Transmission electron micrographs showing immunogold localization of CRA in som leaf tissue reacted with PAb-Cg and labelled with antirabbit-IgG (whole molecule) gold conjugate
- Figure 22:** Transmission electron micrographs showing immunogold localization of CRA in som leaf tissue reacted with PAb-Pt and labelled with antirabbit-IgG (whole molecule) gold conjugate
- Figure 23:** Immunofluorescence of naturally infected som leaf tissue treated with PAb-Cg and labelled with FITC; Transverse section of (A) Healthy and (B and C) infected leaf
- Figure 24:** Immunofluorescence of naturally infected som leaf tissue treated with PAb-Pt and labelled with FITC; Transverse section of (A) Healthy and (B and C) infected leaf

- Figure 25:** Transmission electron micrograph of infected leaf tissue treated with PAb-Cg and labelled with antirabbit-IgG (whole molecule) gold conjugate
- Figure 26:** Transmission electron micrograph of infected leaf tissue treated with PAb-Pt and labelled with antirabbit-IgG (whole molecule) gold conjugate
- Figure 27:** Mycorrhizal association with eight morphotypes of Som plants grown in the experimental field
- Figure 28:** Root colonization (in percent) with arbuscular mycorrhizal fungi in som plants
- Figure 29:** Root colonization of different morphotypes of Som plant with AM fungi in experimental field of NBU showing intraradical hyphae (S-1 and S-7), arbuscules and vesicles embedded in root cells (S-2 and S-8), branched vesicles (S-3, S-5 and S-6) and spore with long hyphal attachment (S-4).
- Figure 30:** Indirect immunofluorescence of som root tissue treated with PAb of AMF and labelled with FITC
- Figure 31:** Transmission electron micrograph of som root tissue treated with PAb-AMF and labelled with antirabbit-IgG (whole molecule) gold conjugate
- Figure 32:** Chromatogram and sequence deposition of 18S r DNA region of *C.gloeosporioides* (KM491736).
- Figure 33:** Multiple sequence alignment of *C. gloeosporioides* (KM491736) with other extypes isolate. The conserved regions of the gene are demonstrated in different colours
- Figure 34:** Phylogenetic placement of *C.gloeosporioides*(KM491736) with other ex-type strain sequences obtained from NCBI GenBank Database
- Figure 35:** PCR amplification of genomic DNA of foliar fungal pathogens, (A) using ITS1/ITS4 primer (M – DNA ladder, CI/02 – *C. gloeosporioides* isolate, P/01 – *P. disseminata* isolate) (B) using primer pair CgINT/ITS4 (Lane 1-4: *Pestalotiopsis* isolates, Lane 5-10: *Bipolaris* isolate; Lane 11- 14: *Colletotrichum* isolates)
- Figure 36:** Chromatogram and sequence deposition of 18S rDNA region of *P.disseminata* (KT697994)
- Figure 37:** Multiple sequence alignment of *Pestalotiopsi* sp (KT697994) with other extypes isolate. The conserved regions of the gene are demonstrated in different colour
- Figure 38:** Phylogenetic placement of *Pestalotiopsis* sp (KT697994) with other ex-type strain sequences obtained from NCBI GenBank Database

- Figure 39:** RAPD-PCR analysis of *Colletotrichum* and *Pestalotiopsis* isolates of som plant using primers (A) – OPD-12, (B) - OPA-1 and (C) - OPD-6. Lane M – DNA ladder, Lane 1-4 *C.gloeosporioides* isolates (SOM/CI/01, SOM/CI/02, SOM/CI/03, SOM/CI/04), Lane 5 – 8 *P. disseminata* isolates (IPL/SOM/P/01, IPL/SOM/P/02, IPL/SOM/P/03, IPL/SOM/P/04)
- Figure 40:** Denature Gradient Gel Electrophoresis of the ITS-PCR amplified products of isolates of *Colletotrichum gloeosporioides* (Lane 1-4; SOM/CI/01, SOM/CI/02, SOM/CI/03, SOM/CI/04) and *Pestalotiopsis disseminata* (Lane 5-8; IPL/SOM/P/01, IPL/SOM/P/02, IPL/SOM/P/03, IPL/SOM/P/04). (A) – ITS-PCR products, 300bp; (B) – Gradient 20-60%, 8hrs run, 100V
- Figure 41:** Analysis of defense enzymes in healthy and *C. gloeosporioides* infected leaves of som morphotypes; A – Peroxidase; B – PAL; C – Chitinase, D -  $\beta$  1,3Glucanase
- Figure 42:** (A and B) – Peroxyzyme analysis of healthy and infected leaves of som morphotypes.
- Figure 43:** (A-E) HPLC profile of standard phenolic acids; A – Resorcinol, B – Catechol, C – Salicylic acid, D – Chlorogenic acid, E – Ferulic acid; (F-G ) HPLC analysis of phenolic acids in leaves of Som plant; F – Healthy leaf, G – Infected leaf (1 – Resorcinol, 2 – Catechol, 3 – Chlorogenic acid, 4 – Morin, 5 and 7 – Ferulic acid, 6 – Salicylic acid)
- Figure 44:** *In vitro* antagonistic test of PGPR and PGPF against *C. gloeosporioides* isolate (SOM/CI/02), (A) – *C. gloeosporioides* culture grown alone in PDA, (B) – interaction with *B. pumilus*, (C) interaction with *T. harzianum*
- Figure 45:** *In vitro* antagonistic test of PGPR and PGPF against against *P. disseminata* isolate (IPL/SOM/P/01), (A) – *P. disseminata* culture grown alone in PDA, (B) – interaction with *B. pumilus*, (C) interaction with *T. harzianum*
- Figure 46:** Growth enhancement in som plants in field following treatment with AMF and PGPR singly and jointly
- Figure 47:** Growth promotion of som plants in glass house conditions after 45days of treatment with *Bacillus altitudinus*
- Figure 48:** Growth promotion in som plants after 45 days of treatment with bioinoculants [Row 1 – Vermi+*B.altitudinus*+*G.constrictum*; Row 2 – Vermi+*B.altitudinus*; Row 3 - Vermi; Row 4 – *B.altitudinus*; Row 5 – *G.constrictum*; Row 6 – Control]
- Figure 49:** Total soluble protein (A) and Total phenol (B) contents in som plants following treatments with Vermicompost, PGPR and AMF

- Figure 50:** Activities of defense enzymes (A) PAL, (B) POX, (C) CHT and (D) GLU in leaves of som plants following treatment with value added Vermicompost
- Figure 51:** Analysis of Total Phenol (A), Orthophenol (B) and Total Soluble protein (C) content in som plants after treatment under nursery conditions
- Figure 52:** Activities of defense enzymes, PAL (A), Peroxidase (B), Chitinase (C) and  $\beta$ -1,3Glucanase (D) in control and treated som plants under nursery conditions.
- Figure 53:** Percent disease index in som plants after treatment with bioinoculants and inoculated with pathogen [UI=Untreated Inoculated, Cg = *C. gloeosporioides*]
- Figure 54:** Activity of defense enzymes in bioinoculants treated som plants following artificial inoculation with pathogen (A – Peroxidase, B – PAL, C- Chitinase , D –  $\beta$ -1,3 Glucanase, Cg – Treated with *C. gloeosporioides*)
- Figure 55:** Percent Disease Index in som plants following treatment and artificial inoculation with *P. disseminata*; UI – untreated inoculated, T1 – Vermicompost, T2 – *B. altitudinus*, T3 – *G.constrictum*, T4 – Vermi+*B. altitudinus*, T5 – Vermi+*B. altitudinus*+*G. constrictum*
- Figure 56:** Growth promotion following application of bioinoculants in 18 month old som plant under field condition. A&B – Morphotype S5; C&D – Morphotype S6; A&C – Untreated Control; B&D – AMF, PGPF and PGPR treated plants
- Figure 57:** HPLC analysis of the phenolic acid content in control and treated som plants under field condition
- Figure 58:** Changes in levels of defense enzymes: Peroxidase (A & B), PAL (C & D), Chitinase (E & F) and Glucanase (G & H) as well as changes in total phenol content (I & J) in S5 and S6 plants following artificial inoculation with *C.gloeosporioides*
- Figure 59:** Cellular localization of glucanase in som leaf tissues following treatment with bioinoculants, probed with PAb of glucanase and labelled with FITC (A&B)
- Figure 60:** Cellular localization of glucanase in roots following treatment with bioinoculants, probed with PAb of glucanase and labelled with FITC (A&B)
- Figure 61:** Transmission electron micrographs of bioinoculant treated leaf tissues of som plant (S6) reacted with PAb of glucanase and labelled with antirabbit goat IgG (whole molecule) gold conjugate

**Figure 62:** Transmission electron micrographs of bioinoculant treated root tissues of som plant (S6) reacted with PAb of glucanase and labelled with antirabbit goat IgG (whole molecule) gold conjugate

**Figure 63:** Cellular localization of chitinase in som leaf tissues following treatment with bioinoculants, probed with PAb of chitinase and labelled with FITC (A&B)

**Figure 64:** Cellular localization of chitinase in roots following treatment with bioinoculants, probed with PAb of chitinase and labelled with FITC (A&B)

**Figure 65:** Transmission electron micrographs of bioinoculant treated leaf tissues of som plant (S6) reacted with PAb of chitinase and labelled with antirabbit goat IgG (whole molecule) gold conjugate

**Figure 66:** Transmission electron micrographs of bioinoculant treated root tissues of som plant (S6) reacted with PAb of chitinase and labelled with antirabbit goat IgG (whole molecule) gold conjugate