

LIST OF TABLES

Table	Caption	Page
Table 1.2.1.	Susceptibility functions ($\chi^{(2)}$ and $\chi^{(3)}$) for various types for interacting field components, their effects and applications.	9
Table 3.1.	HOMO/LUMO energies and β_{tot} values of NiTDT and its derivative computed in gas-phase at UB3LYP/6-311++g(d,p) level of theory.	45
Table 3.2.	TDDFT calculated transitions for thienyl (1a) and phenyl (2a) expanded analogues of NiTDT core and its derivative (1b, 1c & 2b, 2c) at CAM-B3LYP/6-311g(d,p) level of theory.	46
Table 3.3.	Transition energies (eV, nm), their oscillator strengths (f) and transition dipole moment densities for three strong transitions for Ph-NiTDT. The purple color implies charge increase upon excitation and the turquoise color implies charge depletion. The isovalue is 4×10^{-4} in a.u.	49
Table 3.4.	Major transition energies (eV, nm), their oscillator strengths (f), Δr index and eigenvalues for three strong transitions for two sets of complexes (1a, 1b, 1c & 2a, 2b, 2c) calculated at CAM-B3LYP/6-311g(d,p)	49
Table 4.1.	Calculated absolute energies in atomic units (au), $\langle S^2 \rangle$ values, intramolecular magnetic exchange coupling constants (J in cm^{-1}) between two radical centers for the designed diradicals at UB3LYP/6-311++g (2df, 2p) level.	61
Table 4.2.	The spin-spin ZFS parameter D_{Total} (cm^{-1}), static ZFS magnitude a_2 (cm^{-1}), rhombic ZFS parameter (E) for each ferromagnetic diradical at BPW91/EPR-II and two photon absorption cross section (σ_{TPA} in GM), for the same diradicals at UB3LYP/6-311++g(d,p) level.	63
Table 4.3.	The magnetic exchange coupling constants (J) and ZFS parameter (D) values for the proposed ferromagnetic diradical systems in gas phase, water and blood plasma media, computed using PCM model using UB3LYP/6-311++g(2df,2p) and BPW91/EPR-II method.	64
Table 5.1.	Bond distance between Cr and axial ligand (Cr- L_{ax}), Cr and equatorial ligand (Cr- L_{eq}) and energies of the $[\text{Cr}^{III}(\text{NH}_3)_6]^{3+}$ at quartet ground state (Q), quartet excited state (Q'), and doublet ground state (D).	70
Table 5.2.	Comparison of absorption and emission wavelengths in nm (energies in cm^{-1}) with experimental value in $[\text{Cr}^{III}(\text{NH}_3)_6]^{3+}$.	71

Table 5.3.	Bond distance between Cr and axial ligand (Cr-L _{ax}), Cr and equatorial ligand (Cr-L _{eq}) and energies of the complex [Cr ^{III} (NH ₃) ₄ (X) ₂] ⁺ (X= I ⁻ , Br ⁻ , Cl ⁻ , and F ⁻ , represented by violet, brown, green and light blue colors respectively).	72
Table 5.4.	Calculated hardness values in eV for the studied complex (ground state quartet).	73
Table 5.5.	Comparison of absorption and emission wavelengths in nm (energies in cm ⁻¹) in [Cr ^{III} (NH ₃) ₄ (X) ₂] ⁺ (X= I ⁻ , Br ⁻ , Cl ⁻ , and F ⁻).	74
Table 6.1.	Bonding energy decomposition of ground and excited states of the dimer.	79
Table 6.2.	Comparison of the calculated value λ_{max} of absorption and fluorescence with experimentally reported values.	81
Table 6.3.	HOMO-LUMO gap of monomer unit and hydrogen bonded dimer.	82
