

## LIST OF TABLES

CHAPTERS	TABLES	PAGE NO.
<b>Chapter IV</b>	<p><b>Table 1.</b> Values of surface tension (<math>\gamma</math>) and conductance (<math>k</math>) at the break point with corresponding concentration of 15-C-5 for NTPH at 298.15K<sup>a</sup></p> <p><b>Table 2.</b> Data for the Surface tension study of aqueous 15-C-5+NTPH (concentration of stock solution of NTPH = 10mM, concentration of stock solution of CD = 10mM) at 293.15K<sup>a</sup>.</p> <p><b>Table 3.</b> Data for the Conductivity study of aqueous 15-C-5+NTPH (concentration of stock solution of NTPH = 10mM, concentration of stock solution of CE = 10mM) at 293.15K<sup>a</sup>.</p>	<b>116-118</b>
<b>Chapter V</b>	<p><b>Table 1.</b> Table 1: Source and purity of the chemicals</p> <p><b>Table 2.</b> Table 2: Experimental values of density (<math>\rho</math>) and viscosity (<math>\eta</math>) at different temperatures, refractive index (<math>n_D</math>) at 298.15 K of different mass fraction (<math>w_1</math>) of aq. uric acid mixtures*</p> <p><b>Table 3.</b> Table3: Experimental values of density (<math>\rho</math>) and viscosity (<math>\eta</math>) of Allopurinol in different mass fractions of aqueous uric acid mixture (<math>w_1</math>) at three different temperatures.</p> <p><b>Table 4.</b> Table 4: Apparent molar volume (<math>\varphi_V</math>) and <math>(\eta_r - 1)/\sqrt{m}</math> of Allopurinol in different mass fraction (<math>w_1</math>) of aqueous uric acid mixtures at three different temperatures*</p> <p><b>Table 5.</b> Table 5: Limiting apparent molar volume (<math>\varphi_{V^0}</math>), experimental slope (<math>S_V^*</math>), viscosity A- and B-coefficient of allopurinol in different mass fraction (<math>w_1</math>) of aqueous uric acid mixtures at different temperatures*</p> <p><b>Table 6.</b> Table 6: Values of various coefficients of equation-3 for allopurinol in different aqueous uric acid solutions</p> <p><b>Table 7.</b> Table Limiting apparent molar expansibilities (<math>\varphi_{E^0}</math>) for allopurinol in different mass fraction of aqueous uric acid (<math>w_1</math>) at different temperature</p> <p><b>Table 8.</b> Values of <math>dB/dT</math>, <math>A_1</math> and <math>A_2</math> coefficients for the allopurinol in different mass fraction of aqueous uric acid (<math>w_1</math>) at studied temperatures*</p> <p><b>Table 9.</b> Values of <math>V_1^0</math>, <math>(V_1^0 - V_2^0)</math>, <math>\Delta\mu_1^{0\#}</math>, <math>\Delta\mu_2^{0\#}</math>, <math>T\Delta S_2^{0\#}</math> and <math>\Delta H_2^{0\#}</math> for allopurinol in different mass fraction (<math>w_1</math>) of aqueous uric acid mixture at different temperatures*</p> <p><b>Table 10.</b> Refractive index (<math>n_D</math>), molar refraction (<math>R_M</math>) and limiting molar refraction (<math>R_{M^0}</math>) allopurinol in different mass fraction of aqueous uric acid solutions at 298.15 K.*</p>	<b>135-140</b>

CHAPTERS	TABLES	PAGE NO.
Chapter VI	<p><b>Table1(a).</b> Data for the Surface tension study of aqueous THP VS <math>\alpha</math>-CYD (concentration of stock solution of THP = 10mM, concentration of stock solution of CYD = 10mM) at 298.15K<sup>a</sup>.</p> <p><b>Table1(b).</b> Data for the Surface tension study of aqueous THP VS HP-<math>\beta</math>-CYD (concentration of stock solution of THP = 10mM, concentration of stock solution of CYD = 10mM) at 298.15K<sup>a</sup></p> <p><b>Table 2.</b> Values of Surface tension and at the break point with corresponding concentration of <math>\alpha</math>-CYD and HP- <math>\beta</math>-CYD for THP at 298.15 K</p> <p><b>Table3(a).</b> Variation of Conductivity of THP VS <math>\alpha</math>-CYD at 298.15K</p> <p><b>Table3(b).</b> Variation of Conductivity of THP VS HP- <math>\beta</math>-CYD at 298.15K</p> <p><b>Table 4.</b> Values of Conductivity and at the break point with corresponding concentration of <math>\alpha</math>-CYD and HP- <math>\beta</math>-CYD for THP at 298.15 K</p> <p><b>Table5(a).</b> Data for Job plot obtained from UV-spectroscopy for aqueous THP+ <math>\alpha</math>-CYD system at 298.15K<sup>a</sup></p> <p><b>Table5(b).</b> Data for Job plot obtained from UV-spectroscopy for THP+HP- <math>\beta</math>-CYD system at 298.15K<sup>a</sup></p> <p><b>Table 6.</b> Association constant (<math>K_a</math>) and thermodynamic parameters <math>\Delta H^\circ</math>, <math>\Delta S^\circ</math> and <math>\Delta G^\circ</math> of THP-CYDs inclusion complexes at 293.15K, 298.15K, 303.15K temperatures</p> <p><b>Table 7a.</b> Data for IR spectra of THP</p> <p><b>Table 7b.</b> Data for IR spectra of <math>\alpha</math>-CYD</p> <p><b>Table 7c.</b> Data for IR spectra of HP-<math>\beta</math>-CYD</p> <p><b>Table 7d.</b> Data for IR spectra of <math>\alpha</math>-CYD +[THP] inclusion complex.</p> <p><b>Table 7e.</b> Data for IR spectra of HP-<math>\beta</math>-CYD+THP complex</p> <p><b>Table 8a.</b> Data for NMR spectra of HP-<math>\beta</math>-CYD &amp; THP+ HP- <math>\beta</math>-CYD complex(IC-1)</p> <p><b>Table 8b.</b> Data for NMR spectra of <math>\alpha</math>-CYD and THP+<math>\alpha</math>-CYD complex(IC-2)</p> <p><b>Table8c.</b> Data for NMR spectra for significant peak of THP and THP+ HP-<math>\beta</math>-CYD complex(IC-1)</p> <p><b>Table8d.</b> Data for NMR spectra for significant peak of <math>\alpha</math>-CYD and THP+<math>\alpha</math>-CYD complex(IC-2)</p>	154-162

CHAPTERS	TABLES	PAGE NO.
Chapter VII	<p><b>Table 1.</b> Sample descriptions.</p> <p><b>Table 2.</b> Experimental and literature values of density (<math>\rho</math>), viscosity (<math>\eta_0</math>) and relative permittivity (<math>\epsilon</math>) of the solvents at 293.15 K, 303.15K and 313.15K temperatures and 0.101M Pa pressure.</p> <p><b>Table 3.</b> The concentration (c) and molar conductivity (<math>\Lambda</math>) of IL in 2-methoxy ethanol (ME), 2-ethoxy ethanol (EE) and 2-propoxy ethanol (PE) at 293.15, 303.15 and 313.15K respectively and 0.101MPa Pressure.</p> <p><b>Table 4.</b> Limiting molar conductance (<math>\Lambda_0</math>), association constant (<math>K_A</math>), co-sphere diameter(<math>R</math>) and standard deviations of experimental <math>\Lambda(\delta)</math> obtained from Fuoss conductance equation for IL in ME, EE and PE at 293.15, 303.15, and 313.15 K respectively and 0.101MPa pressure</p> <p><b>Table 5.</b> Walden product (<math>\Lambda_0\eta</math>) and Gibbs energy change (<math>G^\circ</math>) of IL in ME,EE and PE at 293.15 K to 313.15 K temperatures and 0.101MPa pressure.</p> <p><b>Table 6.</b> Ionic limiting molar conductance (<math>\lambda_0^\pm / \text{S m}^2 \text{ mol}^{-1}</math>), ionic Walden product (<math>\lambda_0^\pm\eta / \text{S m}^2 \text{ mol}^{-1} \text{ mPa s}</math>), crystallographic radii (<math>r_c/\text{\AA}</math>) and Stoke's radii (<math>r_s/\text{\AA}</math>) of IL (having <math>[\text{MMIm}]^+</math> cation and <math>[\text{MSO}_4]^-</math> anion) in ME,EE and PE at various Kelvin and 0.101 MPa pressure</p> <p><b>Table 7.</b> Concentration, m, density, <math>\rho</math>, apparent molar volume, <math>\Phi_v</math>, limiting apparent molar volume <math>\Phi_v^0</math>, and exponential slope (<math>S_v^*</math>) for IL in ME, EE and PE at 293.15K, 303.15K and 313.15K respectively and 0.101 M Pa pressure.</p> <p><b>Table 8.</b> Values of empirical coefficients (<math>a_0</math>, <math>a_1</math>, and <math>a_2</math>) of temperature dependent apparent molar volume for IL in different solvents (ME,EE, PE) at 293.15 K to 313.15 K and 0.101MPa pressure.</p> <p><b>Table 9.</b> Limiting apparent molal expansibilities for IL in different solvents (ME, EE, PE) at 293.15 K to 313.15 K and 0.101MPa pressure</p> <p><b>Table 10.</b> Concentration, viscosity, <math>\eta</math>, <math>(\eta / \eta_0 - 1) / \sqrt{c}</math>, viscosity A- and B-coefficients for IL in ME, EE and PE at 293.15 K, 303.15 K and 313.15 K respectively and 0.101MPa pressure.</p> <p><b>Table 11.</b> Stretching frequencies of the functional groups present in the pure solvent and change of frequency of IL in ME, EE and PE.</p>	185-194

CHAPTERS	TABLES	PAGE NO.
<b>Chapter VIII</b>	<p><b>Table 1.</b> Data of Job's plot between ALP and <math>\alpha</math>-CD obtained from UV spectroscopy at 298.15K<sup>a</sup></p> <p><b>Table 2.</b> Data of Job's plot between ALP and <math>\beta</math>-CD obtained from UV spectroscopy at 298.15K<sup>a</sup></p> <p><b>Table 3.</b> Data of Job's plot between ALP and HP-<math>\beta</math>-CD obtained from UV spectroscopy at 298.15K<sup>a</sup></p> <p><b>Table 4.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [ALP]-<math>\alpha</math>-CD systems at (293.15, 298.15, and 303.15) K</p> <p><b>Table 5.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [ALP]-<math>\beta</math>-CD systems at (293.15, 298.15, and 303.15) K</p> <p><b>Table 6.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [ALP]-HP-<math>\beta</math>-CD systems at (293.15, 298.15, and 303.15) K</p>	<b>206-208</b>

CHAPTERS	TABLES	PAGE NO.
Chapter IX	<p><b>Table1.</b> Values of surface tension (<math>\gamma</math>) and at the break point with corresponding concentration of <math>\alpha</math> -CD, <math>\beta</math>-CD for [BMPy]PF<sub>6</sub> at 298.15 K<sup>a</sup></p> <p><b>Table2.</b> Association constants obtained by the Benesi-Hildebrand method (<math>K_a</math>) from UV-Vis study and corresponding thermodynamic parameters and stoichiometry of [BMPy]PF<sub>6</sub> &amp; <math>\alpha</math>-CD and [BMPy]PF<sub>6</sub> &amp; <math>\beta</math>-CD inclusion complexes at 293.15K<sup>a</sup>, 303.15K<sup>a</sup> at 313.15K<sup>a</sup>.</p> <p><b>Table3.</b> Association constants obtained by the nonlinear program (<math>K_a^c</math>) from conductance study and corresponding thermodynamic parameters and stoichiometry of [BMPy]PF<sub>6</sub> &amp; <math>\alpha</math>-CD and [BMPy]PF<sub>6</sub> &amp; <math>\beta</math>-CD inclusion complexes at 293.15K<sup>a</sup>, 303.15K<sup>a</sup> at 313.15K<sup>a</sup>.</p> <p><b>TableS1.</b> Data for Job's Plot performed by UV-Vis spectroscopy for [BMPy]PF<sub>6</sub>-<math>\beta</math>-CD system at 298.15 K<sup>a</sup></p> <p><b>TableS2.</b> Data for Job's Plot performed by UV-Vis spectroscopy for [BMPy]PF<sub>6</sub>-<math>\alpha</math>-CD system at 298.15 K<sup>a</sup></p> <p><b>TableS3.</b> 1H-NMR spectra of [BMPy]PF<sub>6</sub>, <math>\alpha</math>-CD, <math>\beta</math>-CD, and [BMPy]PF<sub>6</sub>+<math>\alpha</math>-CD, [BMPy]PF<sub>6</sub>+<math>\beta</math>-CD complexes.</p> <p><b>Table S4.</b> Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin molecule in two different host-guest complexes in D<sub>2</sub>O at 298.15 K<sup>a</sup>.</p> <p><b>TableS5.</b> Surface Tension (<math>\gamma</math>) values of <math>\alpha</math>-CD&amp;[BMPy]PF<sub>6</sub>, <math>\beta</math>-CD &amp; [BMPy]PF<sub>6</sub>Complexat 298.15 K</p> <p><b>TableS6.</b> Data for the conductivity study of aqueous [BMPy]PF<sub>6</sub>+<math>\alpha</math>-CD and [BMPy]PF<sub>6</sub>+<math>\beta</math>-CD system (concentration of stock solution of IL = 10mM, concentration of stock solution of CD = 10mM) at 293.15K<sup>a</sup>, 303.15K<sup>a</sup>, 313.15K<sup>a</sup></p> <p><b>TableS7.</b> Values of conductivity (<math>\kappa</math>) at the break point with corresponding concentrations of [BMPy]PF<sub>6</sub> and CDs at 298.15 K<sup>a</sup></p> <p><b>TableS8.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF<sub>6</sub>-<math>\alpha</math>-CD and [BMPy]PF<sub>6</sub>-<math>\beta</math>-CD systems at 293.15 K</p> <p><b>TableS9.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF<sub>6</sub>-<math>\alpha</math>-CD and [BMPy]PF<sub>6</sub>-<math>\beta</math>-CD systems at 303.15 K</p> <p><b>TableS10.</b> Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF<sub>6</sub>-<math>\alpha</math>-CD and [BMPy]PF<sub>6</sub>-<math>\beta</math>-CD systems at 303.15 K</p> <p><b>TableS11.</b> Data obtained from FT-IR spectroscopic study of <math>\alpha</math>-CD, <math>\beta</math>-CD, [BMPy]PF<sub>6</sub>,<math>\alpha</math>-CD +[BMPy]PF<sub>6</sub>, <math>\beta</math>-CD+[BMPy]PF<sub>6</sub>.</p> <p><b>TableS12.</b> The observed peaks at different m/z with corresponding ions for the solid inclusion complexes.</p>	245-254

