

## LIST OF FIGURES

	<b>Description</b>	<b>Page</b>
<b>Figure 1</b>	Pattern recognition receptors and their function against antigens	<b>11</b>
<b>Figure 2</b>	Structural model of the full-length TLR3/dsRNA	<b>12</b>
<b>Figure 3</b>	Mammalian TLR signalling pathways	<b>13</b>
<b>Figure 4</b>	TLR signaling and membrane processing in pDCs polymorphic variation of the TLRs in Human populations	<b>16</b>
<b>Figure 5</b>	Geographical distribution of Typhoid fever	<b>43</b>
<b>Figure 6</b>	1.2% agarose gel picture showing genomic DNA of different populations (1.Muslim, 2.Rabha, 3.Gurkha) and (4) HIV disease sample	<b>58</b>
<b>Figure 7</b>	1% agarose gel of TLR bands (after PCR) in case of (5) Rheumatoid arthritis, (6) Human Immunodeficiency virus, and (7) (8) Typhoid fever in different disease patients	<b>59</b>
<b>Figure 8</b>	1% Agarose gel picture of TLR4 genes (after PCR) in different populations of North Bengal region of India	<b>60</b>
<b>Figure 9</b>	1% agarose gel electrophoresis (after PCR) to show the presence of TLR8 gene in 8 Rajbanshi Individuals. The marker used in Lane 1 is Lambda DNA-Hind III double digested $\Phi$ X174 DNA	<b>63</b>
<b>Figure 10</b>	Frequency graph of ten TLR genes of the four populations in the North Bengal region (Kyplot ver-2.0)	<b>72</b>
<b>Figure 11</b>	Principle Component Analyses (PCA) based of the 10 TLR genes in the four ethnic populations of North Bengal (Minitab ver-6)	<b>74</b>
<b>Figure 12</b>	Neighbour joining tree showing relationship among Rajbanshi and three other populations (POPGENE ver- 1.32 and Phylip ver-3.5)	<b>76</b>
<b>Figure 13</b>	Frequency graph of ten TLR genes of rheumatoid arthritis patients (Kyplot ver-2.0)	<b>78</b>
<b>Figure 14</b>	Frequency graph of ten TLR genes of typhoid patients (Kyplot ver-2.0)	<b>83</b>
<b>Figure 15</b>	Frequency graph of ten TLR genes was of HIV positive patients	<b>89</b>