

List of Figures:

Figure 1.1. Schematic diagram of three different types of thermotropic liquid crystals: (a) calamitic, (b) discotic, and (c) bent-core	4
Figure 1.2. General representation of molecular ordering in (a) uniaxial and (b) biaxial nematic phase	6
Figure 1.3. Structural representation of molecular ordering in smectic <i>A</i> phase. The director \hat{n} is parallel to the smectic layer normal (\hat{k})	8
Figure 1.4. Structural representation of molecular ordering in layers for smectic <i>C</i> phase. The director \hat{n} is tilted with respect to the smectic layer normal \hat{k} by a certain angle	9
Figure 1.5. Structural representation of molecular ordering in each layer for hexatic smectic <i>B</i> phase.....	10
Figure 1.6. Structural representation of molecular ordering in each layer for smectic <i>I</i> and smectic <i>F</i> phases	11
Figure 1.7. Molecular geometry of soft crystal <i>E</i> phase	12
Figure 1.8. (a) Molecular representation of cholesteric or chiral nematic phase, (b-c) Optical textures appeared in cholesteric phase at two different temperatures ...	13
Figure 1.9. (a) Helical superstructure of the chiral smectic <i>C</i> (<i>Sm-C*</i>) phase with the helix axis parallel to the smectic layer normal \hat{k} , (b) Orientation of a single molecule in each layer. The polarization vector \vec{P}_s is always perpendicular to the director \hat{n} and the tilt plane	15
Figure 1.10. (a) Molecular representation in a double twist cylinder. Relative orientation of the double twist cylinders in case of the (b) <i>BPII*</i> and (c) <i>BPI*</i> phase.....	17
Figure 1.11. A structural representation of the molecular superstructure with screw dislocations for (a) <i>TGBA</i> and (b) <i>TGBC*</i> phases.....	18
Figure 1.12. Molecular structures of (a) a symmetrical and (b) an asymmetrical bent-core compound.....	21
Figure 1.13. Schematic illustration of molecular geometry in <i>B₁</i> mesophase. a and b give the measure of periodicities in the two perpendicular directions.....	23
Figure 1.14. Representations of molecular geometry in (a) three mutually orthogonal planes and (b) four fundamental variants of <i>Sm-CP</i> phase in the <i>B₂</i> phase	24
Figure 1.15. Schematic representation of the in-plane molecular packing in the <i>B₅</i> phase. Vectors a and b suggests the presence of in-plane order	26

Figure 1.16. Schematic illustration of molecular ordering in the B_6 phase.....	27
Figure 1.17. Sketch of the molecular arrangement in bent-core nematic liquid crystals comprising smectic-like cybotactic clusters	30
Figure 1.18. Structural representation of a hockey stick-shaped compound H-22.5 [details in Chapter 5].....	31
Figure 2.1. Orientations of the molecules in planar (HG) and homeotropic (HT) cells.....	53
Figure 2.2. Schematic diagram of the experimental set-up for refractive index measurement.....	56
Figure 2.3. Schematic diagram of the experimental set-up for optical transmission measurement.....	58
Figure 2.4. Schematic diagram of the experimental set-up for the dielectric study	60
Figure 2.5. Frequency-dependent variation of the real (ϵ') and imaginary (ϵ'') parts of dielectric permittivity for a fixed temperature of a bent-core compound-1/7 (described in chapter 8).....	63
Figure 2.6. Voltage dependent variation of sample filled cell-capacitance due to electrically driven Fredericksz transition in the nematic phase of a binary mixture (described in chapter 4) at 345 K. Inset shows the closer view near the threshold voltage	65
Figure 2.7. Schematic diagram of the experimental set-up for electro-optical study using field reversal polarization technique	69
Figure 2.8. Input and output waveforms of the LC sample for (a) square wave and (b) triangular wave input voltages.....	69
Figure 2.9. Orientational order parameter predicted from Maier-Saupe mean field theory.....	74
Figure 2.10. (a) The free energy density (f) as a function of scalar order parameter S . (b) The nematic order parameter (S) vs. temperature. T^* and T^{**} are the temperatures slightly lower and higher than T_{IN} respectively	83
Figure 2.11. Schematic diagram of the heat capacity variation near the phase transition for the dimensions $d > 4$ and $d < 4$. Red solid line is the mean field heat capacity and green dashed line corresponds to the same including the correction term appears due to the fluctuation of the correlation length.....	90
Figure 3.1. Chemical structure and phase behavior of (a) the hockey stick-shaped compound (SF7) and (b) the rod-like compound (8OCB).....	105

Figure 3.2. Phase diagram for the binary mixture consisting of SF7 and 8OCB. x_{SF7} denotes the mole fraction of SF7. <i>I</i> – isotropic phase, <i>N</i> – nematic phase and Sm-A – smectic-A phase. ■: <i>I</i> – <i>N</i> transition temperature; ●: <i>N</i> –Sm-A transition temperature; □: Melting temperatures measured during heating cycle. Inset depicts variation of nematic range with concentration for the present system. Solid lines are drawn for guidance to eye.....	106
Figure 3.3. Temperature dependence of optical birefringence (Δn) data for all the mixtures along with that for pure 8OCB. Vertical arrows show the <i>N</i> –Sm-A phase transition for all the mixtures	108
Figure 3.4. Refractive indices (n_e, n_o) and birefringence ($\Delta n = n_e - n_o$) as a function of temperature for (a) $x_{\text{SF7}} = 0.012$, (b) $x_{\text{SF7}} = 0.03$, (c) $x_{\text{SF7}} = 0.05$ and (d) $x_{\text{SF7}} = 0.065$. Dashed arrow and solid arrows denote the isotropic–nematic (T_{IN}) and nematic–smectic-A (T_{NA}) transition temperatures respectively.....	109
Figure 3.5. Variation of the temperature-dependent optical birefringence data for all the mixtures including pure 8OCB. Dashed arrow denotes the nematic–smectic-A (T_{NA}) transition temperature. The solid line is a fit to Eq. (3.3)	113
Figure 3.6. Temperature dependent variation of the quotient $Q(T)$ in the vicinity of <i>N</i> –Sm-A phase transition for different mole fractions of SF7. Data are arranged in sequence of increasing mole fraction of SF7 from left to right with 1: 8OCB; 2: $x_{\text{SF7}} = 0.012$; 3: $x_{\text{SF7}} = 0.03$; 4: $x_{\text{SF7}} = 0.05$; 5: $x_{\text{SF7}} = 0.065$. The solid lines are fit to Eq. (3.7).....	117
Figure 3.7. Concentration dependence of the effective critical exponent (α') obtained by fitting $Q(T)$ to Eq. (3.7). The vertical dashed line corresponds to the tricritical point (TCP). The solid line is a 2 nd order polynomial fit to the data....	119
Figure 3.8. Variation of effective critical exponent (α') with McMillan ratio (T_{NA}/T_{IN}). The vertical dashed line corresponds to the tricritical point (TCP). The solid line is a linear fit to the data	119
Figure 3.9. Plot of optical birefringence as a function of reduced temperature $ \tau = 1 - T/T_{NA} $ for different mixtures. The solid lines are fit to Eq. (3.8)	121
Figure 3.10. Variation of order parameter critical exponent (β') value with McMillan ratio (T_{NA}/T_{IN}). The solid line is a second-order polynomial fit to the data	122
Figure 3.11. Optimized molecular structure of (a) H-22.5, (b) SF7	124
Figure 4.1. Temperature variation of the dielectric parameters for (a) $x_{\text{SF7}} = 0.012$, (b) 0.03, (c) 0.05 and (d) 0.065. Solid arrow and dashed vertical arrow indicate the <i>I</i> – <i>N</i> (T_{IN}) and the <i>N</i> –Sm-A (T_{NA}) phase transition temperatures respectively	136

Figure 4.2. Temperature variation of the dielectric anisotropy ($\Delta\varepsilon$) for all the mixtures. Inset depicts the average permittivity (ε_{avg}) of the same. Solid arrow and dashed vertical arrow indicate the $I-N$ (T_{IN}) and the $N\text{-Sm-A}$ (T_{NA}) phase transition temperatures respectively	138
Figure 4.3. Variation of permittivities with temperature for the concentration $x_{\text{SF7}} = 0.012$. Solid lines are fit to Eq. (4.1) and (4.2). The critical temperatures T^* and T^{**} are indicated with star symbols (\star) and the dotted vertical line represents the nematic-isotropic phase transition temperature (T_{IN}).	140
Figure 4.4. Variation of the isotropic permittivity (ε_{iso}) with temperature for all of the mixtures. Solid lines are the fit to Eq. (4.1) and the star symbols (\star) are representing the critical temperatures (T^*)	141
Figure 4.5. Temperature-dependent variation of the quotient $E(T)$ in the vicinity of $N\text{-Sm-A}$ phase transition at different mole fractions of SF7 in the binary mixtures. The solid lines are fit to Eq. (4.4).....	144
Figure 4.6. The variation of the effective critical exponent (α') with (a) molar concentration and (b) McMillan ratio (T_{NA}/T_{IN}), obtained by fitting $E(T)$ to Eq. (4.4). The vertical dashed line corresponds to the tricritical point (TCP). The solid lines are the polynomial fits to the data points.....	146
Figure 4.7. Variation of cell capacitance in planar configuration at different temperatures for $x_{\text{SF7}} = 0.012$. Different temperatures are mentioned in the figure in unit of Kelvin (K).....	148
Figure 4.8. The variations of threshold voltage (V_{th}) with temperature for all the mixtures only in nematic phase. The vertical dashed line and solid line corresponds to the T_{IN} and T_{NA} respectively for each of molar concentration.....	148
Figure 4.9. The variation of the splay elastic constant (K_{11}) with reduced temperature for all the mixtures in the nematic phase. The vertical arrows correspond to T_{NA} for each of molar concentration.....	150
Figure 4.10. (a) Energy optimized molecular structure of SF7 and 8OCB. (b) Mutual alignment of both the hockey-stick shaped (SF7) and calamitic molecule (8OCB) in the nematic phase.	151
Figure 4.11. The variation of the relaxation time (τ_0) with reduced temperature for all the mixtures in the nematic phase. The vertical arrows correspond to T_{NA} for each of molar concentration	152
Figure 4.12. The variation of rotational viscosity (γ_1) with reduced temperature for all the mixtures only in nematic phase. The vertical arrows correspond to T_{NA} for each of molar concentration. Solid lines are fit to Eq. (4.7).	154

- Figure 4.13.** Temperature dependence of optical birefringence data for $x_{SF7} = 0.012$. Dashed arrow denotes the Nematic–Smectic-A transition temperature (T_{NA}). The solid line is a fit to Eq. (4.9)..... 157
- Figure 4.14.** The temperature dependence of orientational order parameter ($\langle P_2 \rangle$) for all the mixtures. Dashed arrow denotes the nematic–smectic-A (T_{NA}) transition temperatures respectively. Inset represents the value of $\langle P_2 \rangle$ at a particular temperature in both nematic and smectic phases for all the mixtures..... 159
- Figure 4.15.** The variations of $\ln(\gamma_1/\langle P_2 \rangle)$ with reciprocal of temperature ($1/T$) for all the concentrations only in nematic phase. Solid lines represent a linear fit to the data points 160
- Figure 5.1.** Partial phase diagram of the binary system comprising of PhP2 and PhP1. x_{PhP1} denotes the mole fraction of PhP1. *I*– isotropic phase, *N*– nematic phase, *Sm-A*–smectic-A phase, *Sm-C*–smectic-C phase, ○: isotropic to nematic transition temperature; ●: nematic to smectic-A transition temperature; ■: smectic-A to smectic-C transition temperature; Inset shows the concentration dependence of the *Sm-A* and nematic ranges for the present system. Solid lines are drawn for guidance to eye... .. 174
- Figure 5.2.** Temperature dependence of birefringence (Δn) data for the (a) Pure PhP2 compound and (b) mixture $x_{PhP1} = 0.1$, (c) 0.2 and (d) 0.3. Vertical arrows show the *I*–*N*, *N*–*Sm-A* and *Sm-A*–*Sm-C* phase transitions for the mixtures. In the insets variation of birefringence in the vicinity of *Sm-A*–*Sm-C* transition [inset (a)] and *N*–*Sm-A* transition [inset (b)] has been presented... .. 176
- Figure 5.3.** The temperature variation of the parameter $Q(T)$ near the *Sm-A*–*Sm-C* phase transition. Solid lines are fit to Eq. (5.2) near the *Sm-A*–*Sm-C* phase transition..... 180
- Figure 5.4.** The temperature dependence of $Q(T)$ in the vicinity of the *Sm-A*–*Sm-C* transition in mixtures of PhP1 and PhP2 for $|\tau|_{\max} = 3 \times 10^{-3}$. 1: $x_{PhP1} = 0.0$; 2: $x_{PhP1} = 0.05$; 3: $x_{PhP1} = 0.1$; 4: $x_{PhP1} = 0.15$; 5: $x_{PhP1} = 0.2$; 6: $x_{PhP1} = 0.3$; 7: $x_{PhP1} = 0.35$; 8: $x_{PhP1} = 0.4$; 9: $x_{PhP1} = 0.45$ 183
- Figure 5.5.** Variation of the effective critical exponent (α') with T_{AC}/T_{NA} for *Sm-A*–*Sm-C* phase transition. The vertical dashed line and solid line corresponds to the tricritical point (TCP) and 3D-XY limit respectively. The solid lines (red) are linear fit to the data points..... 184
- Figure 5.6.** Temperature variation of the parameter $Q(T)$ near the *N*–*Sm-A* phase transition for sample PhP2. Solid lines are fit to Eq. (5.2) near the *N*–*Sm-A* phase transition. 186

Figure 5.7. The temperature dependence of $Q(T)$ in the vicinity of the N – $Sm-A$ transition in mixtures of PhP1 and PhP2. 1: $x_{PhP1} = 0.0$; 2: $x_{PhP1} = 0.05$; 3: $x_{PhP1} = 0.1$; 4: $x_{PhP1} = 0.15$; 5: $x_{PhP1} = 0.2$; 6: $x_{PhP1} = 0.3$; 7: $x_{PhP1} = 0.35$; 8: $x_{PhP1} = 0.4$... 188

Figure 5.8. The variation of effective critical exponent (α') with T_{NA}/T_{IN} for the N – $Sm-A$ phase transition and T_{AC}/T_{NA} for the $Sm-A$ – $Sm-C$ phase transition. The vertical dashed lines and solid line corresponds to the tricritical point (TCP) and 3D-XY limit respectively. The solid lines (red) are linear fit to the data... 189

Figure 5.9. Experimental values of birefringence ($\Delta n = n_e - n_o$) as a function of temperature for (a) PhP2, (b) $x_{PhP1} = 0.05$, (c) $x_{PhP1} = 0.1$ and (d) $x_{PhP1} = 0.3$. Solid lines (Red) represent the fit to Eq. (5.3) and extrapolated to the $Sm-A$ phase..... 191

Figure 6.1. Chemical structure and phase behavior of the hockey-stick-shaped compound (Compound 1: H-22.5) and the rod-like compound (Compound 2: 7OCB) 202

Figure 6.2. Optical texture observed for a mixture with mol fraction $x_{7OCB} = 0.184$ in homogeneous alignment (a-c) and homeotropic alignment (d-f). (a) $Sm-A$ phase at 380 K, (b) $Sm-C_s$ phase at 369 K, (c) $Sm-C_a$ phase at 360 K, (d) $Sm-A$ phase at 378 K, (e) $Sm-C_s$ phase at 371 K, (f) $Sm-C_a$ phase at 365 K. Arrows indicates the direction of the polarizers..... 204

Figure 6.3. Partial phase diagram of the binary system comprising of 7OCB and H-22.5. x_{7OCB} denotes the mole fraction of 7OCB. I – isotropic phase, $Sm-A$ – smectic- A phase, $Sm-C_s$ – smectic- C_s phase, $Sm-C_a$ – smectic- C_a phase \square : N – $Sm-C_s$ phase transition; \blacksquare : I – $Sm-A$ phase transition; \bullet : $Sm-A$ – $Sm-C_s$ phase transition; \circ : $Sm-C_s$ – $Sm-C_a$ phase transition temperatures. The inset shows the concentration dependence of the $Sm-A$ range for the present system. Solid lines are drawn for guidance to the eye. 206

Figure 6.4. Temperature dependence of birefringence (Δn) data for (a) $x_{7OCB} = 0.184$ and (b) $x_{7OCB} = 0.167$. \circ : Birefringence data from optical transmission method, \square : Birefringence data from thin prism technique. Blue solid arrows indicate isotropic to smectic- A (I – $Sm-A$) phase transition temperatures (T_{IA}), black solid arrows indicate the smectic- A to smectic- C ($Sm-A$ – $Sm-C_s$) phase transition temperatures (T_{AC}) and dashed arrows indicate the smectic- C_s to smectic- C_a phase transition temperatures (T_{CC}). Insets show the variation of birefringence in the vicinity of $Sm-A$ – $Sm-C_s$ transition temperature (T_{AC}) 208

Figure 6.5. The temperature dependence of $Q(T)$ in the vicinity of $Sm-A$ – $Sm-C_s$ transition in mixtures of H-22.5 and 7OCB for $|\tau|_{max} = 4 \times 10^{-3}$. 8: $x_{7OCB} = 0.211$; 7: $x_{7OCB} = 0.2$; 6: $x_{7OCB} = 0.184$; 5: $x_{7OCB} = 0.174$; 4: $x_{7OCB} = 0.167$; 3: $x_{7OCB} = 0.158$; 2: $x_{7OCB} = 0.15$; 1: $x_{7OCB} = 0.1$. All of these $Q(T)$ data have been shifted in temperature axis for the guidance of eye as follows: curve 7 by +0.5 K, curve 6 by

+0.5 K, curve 4 by +1 K, curve 3 by +1.5 K, curve 2 by +1.5 K. Solid lines representing the fits to Eq. (6.2).....211

Figure 6.6. Variation of the effective critical exponent against (a) the temperature ratio T_{AC}/T_{IA} and (b) the temperature width of Sm-A phase at the Sm-A–Sm-C phase transition in the present investigated binary mixtures. The vertical dashed lines correspond to the tricritical point (TCP). The solid lines (red) are the 2nd order fit to the data points214

Figure 6.7. The temperature dependent variation of $Q(T)$ in the vicinity of Sm-A–Sm-C transition in mixtures of H-22.5 and 7OCB for (a) fit range-B ($|\tau|_{\max} = 4 \times 10^{-3}$), and (b) fit range-A ($|\tau|_{\max} = 2 \times 10^{-3}$). 5: $x_{7OCB} = 0.174$; 4: $x_{7OCB} = 0.167$; 3: $x_{7OCB} = 0.158$; 2: $x_{7OCB} = 0.15$; 1: $x_{7OCB} = 0.1$. These $Q(T)$ data has been shifted in temperature axis for the guidance of eye as follows: curve 7 by +0.5 K, curve 6 by +0.5 K, curve 4 by +1 K, curve 3 by +1.5 K, curve 2 by +1.5 K.....217

Figure 6.8. The representation of extracted z values with the variation of the amplitude ratio A^-/A^+ over three fit ranges in the vicinity of Sm-A–Sm-C transition. Solid line is a 2nd order polynomial to the data points219

Figure 6.9. Variation of the effective critical exponent (α') with the temperature ratio T_{AC}/T_{IA} for Sm-A–Sm-C_s phase transitions of the present binary system. ○: Values of α' for Sm-A–Sm-C transition from chapter 5; □: Values of α' for N–Sm-A transition from chapter 5. The vertical dashed-dot lines are corresponding to the tricritical point (TCP). The solid lines (red) are the linear fit to the data points220

Figure 7.1. Chemical structure and phase behavior of the chiral ferroelectric compound (2H6R).....231

Figure 7.2. Optical textures of N^* phase by lowering the temperature as: (a) nematic droplets at 98.5 °C, (b) cholesteric Grandjean texture of 2H6R at 98.3 °C, (c) cholesteric fan-like texture at 98 °C, (d-e) oily streaks texture at 97.6 °C and 97.2 °C, (f-g) color domain texture of different twist and twist-inversion states at 96.8 °C and 96.6 °C, (h) texture with cholesteric finger at 96.2 °C, (i) oily streaks textures appears again at 95.5 °C, (j-k) texture with cholesteric finger in homeotropic cell at 96.5 °C, 96 °C respectively, (l) coexisting of fingerprint texture with cholesteric finger at 94.3 °C for $x_{H-22.5} = 0.306$ 233

Figure 7.3. Optical textures of TGBA phase by lowering the temperature as follows: (a-b) transition from N^* to TGBA phase in planar configuration at 93.3 °C and 93.25 °C, (c) Grandjean texture of TGBA phase at 92.1 °C, (d-e) filament texture at the transition from N^* to TGBA phase in homeotropic configuration at 93.25 °C and 93.2 °C, (f) appearance of black domain for TGBA phase at 93 °C235

Figure 7.4. Optical textures of TGBC* phase by lowering the temperature: (a) appearance of cross-grid pattern in planar aligned Grandjean texture of TGBA phase at 84.6 °C, (b) well-formed texture of TGBC* phase at 84.5 °C, (c) appearance of modulated texture of TGBC* phase at 80 °C for $x_{H-22.5} = 0.306$, (d) transition to TGBA–TGBC* phase at 84.8 °C in homeotropic cell, (e) well-formed TGBC* phase texture at 82 °C in homeotropic cell	236
Figure 7.5. Optical textures of lower temperature Sm- B^*_{hex} phase at (a) 50 °C for pure 2H6R compound in planar cell, (b) 48.6 °C for $x_{H-22.5} = 0.202$ in homeotropic alignment	237
Figure 7.6. DSC thermograms on both heating and cooling cycle for the mixture concentration $x_{H-22.5} = 0.202$	238
Figure 7.7. (a) Phase diagram for the composite mixture consisting of H-22.5 and 2H6R. $x_{H-22.5}$ denotes the mole fraction of H-22.5. <i>I</i> - isotropic phase, N^* - chiral nematic phase, Sm- C^* - chiral Sm- <i>C</i> phase, TGBA- frustrated smectic- <i>A</i> phase and TGBC*- frustrated smectic- <i>C</i> phase. ○: <i>I</i> – N^* transition temperature; ■: N^* –TGBA transition temperature; □: TGBA–TGBC* transition temperature; ★: N^* –Sm- C^* transition temperature; ●: TGBC*–Sm- B^*_{hex} phase transition temperature. (b) Variation of N^* , TGBA and TGBC* phase range with concentration for the present system. Solid lines are drawn for guidance to the eye	240
Figure 7.8. Temperature-dependent measured optical transmission intensity for compound (a) 2H6R, (b) $x_{H-22.5} = 0.101$, (c) $x_{H-22.5} = 0.202$ and (d) $x_{H-22.5} = 0.306$ respectively in planar (open circle) cell and also homeotropic (closed square) cell configuration. Vertical arrows represent the corresponding phase transition.....	242
Figure 7.9. Temperature-dependent dielectric parameters ($\epsilon_{ }$, ϵ_{\perp} , ϵ_{avg}) for compound compound (a) 2H6R, (b) $x_{H-22.5} = 0.101$, (c) $x_{H-22.5} = 0.202$ and (d) $x_{H-22.5} = 0.306$ in planar or HG (open circle) cell and also homeotropic or HT (open square) cell configuration. The dashed line represents the average permittivity (ϵ_{avg}). Inset depicts the magnified version of permittivity inversion region. Vertical arrows represent the corresponding phase transition	245
Figure 7.10. (a) Variation of dielectric anisotropy ($\Delta\epsilon$) against the reduced temperature ($T-T_{I-N^*}$). Vertical arrows represent the corresponding phase transition temperatures(b) The concentration variation of positive anisotropy ($\Delta\epsilon$) range for all the mixture including the pure chiral 2H6R mesogen	248
Figure 7.11. Value of spontaneous polarization (P_s) as a function of reduced temperature for pure compound (2H6R) and all the mixture components. The solid line is a fitting curve of Eq. (7.2).....	250

Figure 7.12. Reduced temperature dependent value of response time (τ) for pure compound (2H6R) including all the mixture components.....	252
Figure 7.13. (a) Reduced temperature dependent value of effective torsional bulk viscosity (η) for all of the mixtures including the pure compound (2H6R). (b) Arrhenius plot of effective torsional bulk viscosity vs. ($1000/T$). Inset represents the linear fit to Eq. (7.3) to experimental data of $x_{H-22.5} = 0.306$	254
Figure 7.14. (a) Reduced temperature dependent value of effective torsional bulk viscosity (η) for all of the mixtures including the pure compound (2H6R). (b) Arrhenius plot of effective torsional bulk viscosity vs. ($1000/T$). Inset represents the linear fit to Eq. (7.3) to experimental data of $x_{H-22.5} = 0.306$	256
Figure 7.15. (a) Arrhenius plot of the relaxation frequency (f_R) and (b) the variation of dielectric strength ($\delta\epsilon$) at different mesophases for the pure compound 2H6R	257
Figure 7.16. (a) Arrhenius plot of the relaxation frequency (f_R) and (b) the variation of dielectric strength ($\delta\epsilon$) at different mesophases for the mixture concentration $x_{H-22.5} = 0.202$	258
Figure 8.1. Frequency dependence of the real (ϵ') and imaginary (ϵ'') parts of the permittivity for compound 1/7 in (a) HG or planar and (b) HT or homeotropic cell configurations. Red solid lines represent the fitting curve to data points with Eq. (8.2)..	276
Figure 8.2. Temperature dependence of relaxation frequency (f_R) and dielectric strength ($\delta\epsilon$) for compound 1/7 in (a,c) planar and (b,d) homeotropic cells respectively. Vertical arrows define the corresponding phase transition temperatures	276
Figure 8.3. Frequency dependence of the imaginary (ϵ'') parts of the permittivity for compound 1/9 in (a) HG or planar and (b) HT or homeotropic cell configurations.....	279
Figure 8.4. Temperature dependence of relaxation frequency (f_R) and dielectric strength ($\delta\epsilon$) for compound 1/9 in (a,c) planar and (b,d) homeotropic cells respectively. Vertical arrows define the corresponding phase transition temperatures	280
Figure 8.5. The variation of frequency-dependent imaginary (ϵ'') part of permittivity with the bias voltage for compound 1/9 in (a) planar and (b) the homeotropic alignment cells	282
Figure 8.6. Frequency dependence of the imaginary (ϵ'') parts of the complex permittivity for compound 1/10 in (a) HG or planar and (b) HT or homeotropic configuration cells.....	284

Figure 8.7. Temperature dependence of relaxation frequency (f_R) and dielectric strength ($\delta\varepsilon$) for the compound **1/10** in (a,c) planar and (b,d) homeotropic cells respectively. Vertical arrows define the corresponding phase transition temperatures. 285

Figure 8.8. The variation of frequency-dependent imaginary (ε'') part of the complex permittivity with the bias voltage for compound **1/10** in (a) planar and (b) the homeotropic alignment cells. 285

Figure 8.9. Switching current response curves of compound **1/9** and **1/10** for a triangular wave voltage (Red dashed line) at different temperatures; (a-c) Compound **1/9** in HG cell, (d) Compound **1/10** in HG cell, (e) Compound **1/10** in HT cell, (f-g) Temperature-dependent polarization value for the compound **1/9** and **1/10** respectively.. 289

Figure 8.10. Optical textures of the compound **1/9** and **1/10** for an applied square wave AC voltage at different mesophases; (a-c): compound **1/9** in HG cell, (d-f): compound **1/9** in HT cell, (g-j): compound **1/10** in HG cell, (k-n): compound **1/10** in HT cell. Corresponding insets are the textures before the application of the field, and arrows indicate the direction of the analyzer and polarizer... 290