

LIST OF TABLES

CHAPTERS	TABLE. CAPTIONS	PAGE NO
CHAPTER-I	Table.1. Physical properties of Cyclodextrins.	3
CHAPTER-IV	Table.1. Sample Description.	57
	Table.2. Values of density (ρ) and viscosity (η) of aqueous Dopamine Hydrochloride with Standard Deviation (SD) and Average Absolute Deviation (AAD) in different mass fraction (w1) at T= (293.15-313.15) Ka and (p = 0.1MPa) respectively.	59
	Table.3. Experimental values of density (ρ) of amino acids in different mass fraction of aqueous Dopamine Hydrochloride with Standard Deviation (SD) and Average Absolute Deviation (AAD) at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	59-60
	Table.4. Experimental values of viscosity (η) of amino acids in different mass fraction of aqueous Dopamine Hydrochloride with Standard Deviation (SD) and Average Absolute Deviation (AAD) at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	64-65
	Table.5. Limiting apparent molar volume (Φ_V^0), experimental slopes (Sv*) and viscosity A and B-coefficient of amino acids in different mass fraction of aqueous Dopamine Hydrochloride at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	65-66
	Table.6. Limiting apparent molar expansibilities(Φ_E^0) of amino acids in different mass fraction of aqueous Dopamine Hydrochloride at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	67
	Table.7. Values of dB/dT for Aspartic Acid and Glutamic Acid in different mass fraction of aqueous Dopamine Hydrochloride (w1) at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	67

CHAPTERS	TABLE. CAPTIONS	PAGE NO
	Table.8. Limiting molar refractions (R_M^0) values of amino acids in different mass fraction of aqueous Dopamine Hydrochloride at T= (293.15 to 313.15) Ka and (p = 0.1MPa) respectively.	69
	Table.9. Conductivity values for the Aspartic Acid and Glutamic Acid in aqueous Dopamine Hydrochloride at three different temperature (K_a) and (p = 0.1MPa).	71
	Table.10. Data for the Benesi-Hildebrand double reciprocal plot performed by UV-Vis spectroscopy for DH-Aspartic Acid system.	74
	Table.11. Data for the Benesi-Hildebrand double reciprocal plot performed by UV-Vis spectroscopy for DH-Glutamic Acid system.	75
	Table.12. Values of Association Constant (K_a) and free energy change (ΔG^0) of the two system (Asp+DH) and (Glu+DH).	75
	Table.13. Stretching frequency of N-H band in Raman Spectroscopy	78
	Table.S1. Apparent molar volume(Φ_V) and $\frac{(n_r-1)}{\sqrt{m}}$ for amino acids in different mass fraction of aqueous Dopamine Hydrochloride at different temperatures and (p = 0.1MPa).	64-86
	Table.S2. Values of empirical coefficients (a_0 , a_1 , and a_2) of Equation 3 of amino acid in different mass fraction of aqueous Dopamine Hydrochloride at T= (293.15 to 313.15) K and (p = 0.1MPa).	86
	Table.S3. Values of Refractive Index (n_D) and Molar Refraction (R_M) of Aspartic Acid and Glutamic Acid in different mass fraction of aqueous Dopamine Hydrochloride (w_1) at different temperatures (298.15 Ka, 303.15Ka & 313.15Ka) and (p = 0.1MPa) respectively	87-90

CHAPTERS	TABLE. CAPTIONS	PAGE NO
CHAPTER-V	Table.1. Details of the chemicals used.	95
	Table.2. Association constants (K_a) obtained by the Benesi–Hildebrand method at 293.15K, 303.15K, 313.15K, and corresponding Gibb's free energy change (ΔG^0), enthalpy change (ΔH^0) and entropy change (ΔS^0) of AC. β -CD system.	99
	Table.3. The calculated values and obtained m/z values of the solid inclusion complex in methanol.	103
	Table.S1. Stretching frequencies obtained from FTIR spectroscopic study of pure β -CD, AC, AC. β -CD complex.	105
	Table.S2. Data for Benesi-Hildebrand double reciprocal plot performed by UV-Visible Spectroscopic Study for of AC. β -CD system at 293.15K.	105
	Table.S3. Data for Benesi-Hildebrand double reciprocal plot performed by UV-Visible Spectroscopic Study for of AC. β -CD system at 303.15K.	106
	Table.S4. Data for Benesi-Hildebrand double reciprocal plot performed by UV-Visible Spectroscopic Study for of AC. β -CD system at 303.15K.	106
	Table.S5. Data for Job plot performed by UV-Visible Spectroscopic Study for of AC. β -CD system at 293.15K.	107
	Table.S6. Data for Job plot performed by UV-Visible Spectroscopic Study for of AC. β -CD system at 293.15K.	108
	Table.S7. Data for Benesi-Hildebrand double reciprocal plot (assuming the stoichiometry to be 1:2) performed by fluorescence Spectroscopic Study for of AC. β -CD system at 293.15K.	108
CHAPTER-VI	Table.1. Density (ρ), viscosity (η) and relative permittivity (ϵ) of the different concentration (m) of aqueous D(-)fructose and D(+)galactose at 298.15, 303.15 and 308.15 K respectively.	115

CHAPTERS	TABLE. CAPTIONS	PAGE NO
	Table.2. Molar conductivities (Λ) of [bmp]Cl in aqueous D(-)fructose solutions as a function of ionic liquid molality (m) at different temperatures.	117
	Table.3. Molar conductivities (Λ) of [bmp]Cl in aqueous D(+)galactose solutions as a function of ionic liquid molality (m) at different temperatures.	118
	Table.4. Ion association constants (KA), limiting molar conductivities (Λ_0), distance parameters (R), Walden product($\Lambda_0\eta$) and standard deviations (δ) obtained from Fuoss conductance equation of IL in aqueous D(-)fructose and D(+)galactose solutions as a function of ionic liquid molality (m) at different temperatures.	121-122
	Table.5. The values of coefficients in Eq.(11) A_0 , A_1 and A_2 at different solvent compositions.	122
	Table.6. Thermodynamic functions (ΔG_A^0 , ΔS_A^0 , ΔH_A^0) of IL in aqueous D(-)fructose and D(+)galactose solutions as a function of ionic liquid molality (m) at different temperatures.	122-123
	Table.7. Limiting apparent molar volume (Φ_V^0), experimental slope (S_V^*), viscosity-B and A co-efficients of IL in aqueous D(-)fructose and D(+)galactose solutions at different temperatures.	126
	Table.8. Values of $\Phi_V^0(\text{aq})$, $\Phi_V^0(\text{aq})_{\text{tr}}$, B(aqueous), ΔB for IL in different solvent systems at different temperatures.	126-127
	Table.9. Values of empirical coefficients (a_0 , a_1 , and a_2) of eqn (17) for IL in different solvent systems.	128
	Table.10. Limiting apparent molal expansibilities (Φ_E^0) and $(\delta\Phi_E^0/\delta T)_P$ for IL in different solvent systems at different temperatures.	128
	Table.11. Values of dB/dT, A_1 and A_2 coefficient of equation (22) for the IL in different solvent systems.	130

CHAPTERS	TABLE. CAPTIONS	PAGE NO
	Table.12. ¹ H NMR data of [bmp]Cl, D(-)fructose, D(+)-galactose and IL- carbohydrates mixture	132
	Table.1. Association constants (K_a), Gibb's free energy (ΔG^0), enthalpy (ΔH^0) and entropy (ΔS^0) of (AA+CD) systems from UV-Visible spectroscopy	142
	Table.2. The calculated values and obtained m/z values of the solid inclusion complexes.	146
	Table.S1. Details of chemicals used	149
	Table.S2. Data for the conductivity study of aqueous [AA]. α -CD and [AA]. β -CD system (concentration of the stock solution of [α -CD] and [β -CD] = 10Mm at 298.15Ka.	149
	Table.S3. Data obtained from FT-IR spectroscopic study of α -CD, β -CD, AA, AA. α -CD and AA. β -CD complex.	150
	Table.S4. Data obtained from FT-IR spectroscopic study of α -CD, β -CD, AA, AA. α -CD and AA. β -CD complex.	150
	Table.S5. Data obtained from FT-IR spectroscopic study of α -CD, β -CD, AA, AA. α -CD and AA. β -CD complex.	151
	Table.S6. Data obtained from FT-IR spectroscopic study of α -CD, β -CD, AA, AA. α -CD and AA. β -CD complex.	151
	Table.S7. Data obtained from FT-IR spectroscopic study of α -CD, β -CD, AA, AA. α -CD and AA. β -CD complex.	151
	Table.S8. Data for the Benesi-Hildebrand double reciprocal plot obtained from UV-VIS spectroscopic study for (AA+ β -CD) and (AA+ α -CD) systems at 313.15 K.	152
	Table.S9. ¹ H-NMR spectra of AA, α -CD, β -CD, and (AA+ α -CD), (AA+ β -CD) inclusion complexes.	152
CHAPTER-VIII	Table.1. Association constants (K_a) obtained by the Benesi-Hildebrand method at 303.15Ka at 313.15Ka, 323.15Ka, and corresponding Gibb's free energy (ΔG^0), enthalpy (ΔH^0) and	167

CHAPTERS	TABLE. CAPTIONS	PAGE NO
	entropy(ΔS^0) of (NBTU+ β -CD) system.	
	Table.2. The calculated values and obtained m/z values of the solid inclusion complex.	171
	Table.S1. Details of the chemicals used.	174
	Table.S2. Stretching frequency of functional groups present in pure NBTU, pure β -CD and NBTU. β -CD inclusion complex.	174-175
	Table.S3. Stretching frequency of functional groups present in pure NBTU, pure β -CD and NBTU. β -CD inclusion complex.	175
	Table.S4. Stretching frequency of functional groups present in pure NBTU, pure β -CD and NBTU. β -CD inclusion complex.	176
	Table.S5. Association constant data for NBTU & β -CD obtained using Fluorescence method.	176
	Table.S6. ¹ H-NMR spectra of pure NBTU, pure β -CD, (NBTU+ β -CD) inclusion complexes.	179
CHAPTER-IX	Table.1. Comparison between the stretching frequencies (cm^{-1}) of the different functional groups of free compound and their complexes.	186-187
	Table.2. Limiting apparent molar volume (Φ_V^0) and viscosity B-coefficient of the sulfa drug in a different mass fraction of different crown ethers in ACN at T= (293.15 to 308.15) K.	196-198
	Table.3. Limiting apparent molal expansibilities (Φ_E^0) of the sulfa drug in a different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= (293.15 to 308.15) K.	198-199
	Table.4. Values of dB/dT of sulfa drug in different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= (293.15 to 308.15) K respectively.	199
	Table.5. Limiting molar refractions (R_M^0) values of sulfa drug in different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= 298.15 K respectively.	201
	Table.6. Values of Association constant (K_a) and free	202

CHAPTERS	TABLE. CAPTIONS	PAGE NO
	energy change (ΔG^0) of the three SA-CEs complexes.	
	Table.S1. Data for the Job plot performed by UV-Vis spectroscopy for SA-DC18C6 system.	203
	Table.S2. Data for the Job plot performed by UV-Vis spectroscopy for SA-18C6 system.	203
	Table.S3. Data for the Job plot performed by UV-Vis spectroscopy for SA-DB18C6 system.	204-206
	Table.S4. Experimental values of density (ρ) and viscosity (η) of sulfa drug in different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= (293.15 to 308.15) K.	206
	Table.S5. Values of empirical coefficients (a_0 , a_1 , and a_2) of Equation 14 of sulfa drug in different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= (293.15 to 308.15) K.	207-208
	Table.S6. Values of Refractive Index (n_D) and Molar Refraction (R_M) of sulfa drug in different mass fraction of DC18C6 (w_1), 18C6 (w_2) and DB18C6 (w_3) in ACN at T= 298.15 K respectively.	208
	Table.S7. Data for the Bensi-Hildebrand double reciprocal plot performed by UV-Vis spectroscopy for SA-DC18C6 system.	208
	Table.S8. Data for the Bensi-Hildebrand double reciprocal plot performed by UV-Vis spectroscopy for SA-18C6 system.	209
	Table.S9. Data for the Bensi-Hildebrand double reciprocal plot performed by UV-Vis spectroscopy for SA-DB18C6 system.	209