

LIST OF FIGURES

CHAPTERS	FIGURES	PAGE NO.
Chapter IV	<p>Figure 1. Molecular structures of (a) SNP, (b) PEH, (c) Cyclodextrins, showing the exterior and the interior protons, here, $n = 6$ to 8 for the α, β and γ-cyclodextrins respectively.</p> <p>Figure 2. (a,b,c,d) UV-Vis spectra for the generation of Job plots of (a) SNP+α-CD and (b) SNP+β-CD systems at $\lambda_{\max} = 209$ nm, and (c) PEH+α-CD and (d) PEH+β-CD systems at $\lambda_{\max} = 219$ nm.</p> <p>Figure 3. Probable host:guest stoichiometric ratio of the inclusion complexes.</p> <p>Figure 4. (a,b,c,d) Job plots of the (a) SNP+α-CD and (b) SNP+β-CD systems at $\lambda_{\max} = 209$ nm and (c) PEH+α-CD and (d) PEH+β-CD systems at $\lambda_{\max} = 219$ nm, at 298.15K. $\Delta A =$ absorbance difference of SNP/PEH without and with CD, $R = [DGs]/([DGs] + [CD])$.</p> <p>Figure 5. (a,b,c,d) Variations in the surface tension of aqueous SNP with increasing concentration of (a) α-CD, (b) β-CD and the variations in the same of aqueous PEH with increasing concentration of (c) α-CD, (d) β-CD at 298.15 K.</p> <p>Figure 6. Schematic representation of the host:guest inclusion complexation through the more favorable wider rim of the cyclodextrin molecules.</p> <p>Figure 7. (a,b,c,d) Variations in the conductivity of aqueous SNP with increasing concentration of (a) α-CD, (b) β-CD and</p>	122-133

the variations in the same of aqueous PEH with increasing concentration of (c) α -CD, (d) β -CD at 298.15 to 308.15 K.

Figure 8. 2D ROESY NMR spectra of the solid (SNP+ α -CD) system.

Figure 9. 2D ROESY NMR spectra of the solid (SNP+ β -CD) system.

Figure 10. 2D ROESY NMR spectra of the solid (PEH+ α -CD) system.

Figure 11. 2D ROESY NMR spectra of the solid (PEH+ β -CD) system.

Figure 12. (a,b) HRMS spectra of the (a) SNP+ α -CD, PEH+ α -CD and (b) SNP+ β -CD, PEH+ β -CD ICs.

Figure 13. (a,b,c) FTIR spectra of (a) SNP+ α -CD, (b) SNP, (c) α -CD.

Figure 14. (a,b,c) FTIR spectra of (a) SNP+ β -CD, (b) SNP, (c) β -CD.

Figure 15. (a,b,c) FTIR spectra of (a) PEH+ α -CD, (b) PEH, (c) α -CD.

Figure 16. (a,b,c) FTIR spectra of (a) PEH+ β -CD, (b) PEH, (c) β -CD.

Figure 17. (a,b,c,d,e,f) SEM images of (a) α -CD, (b) (SNP+ α -CD) physical mixture, (c) (SNP+ α -CD) inclusion complex (d) β -CD, (e) (SNP+ β -CD) physical mixture, (f) (SNP+ β -CD) inclusion complex.

Figure 18. (a,b,c,d,e,f) SEM images of (a) α -CD, (b) (PEH+ α -CD) physical mixture, (c) (PEH+ α -CD) inclusion complex (d)

	<p>β-CD, (e) (PEH+β-CD) physical mixture, (f) (PEH+β-CD) inclusion complex.</p> <p>Figure 19. Antimicrobial activity analysis α-CD, β-CD on Gram-negative E. coli by Agar Cup method. No zone of inhibition was observed. Double distilled water was taken as the control. [Marker points (red) for the verified samples taken in the plates (1. SNP, 2. PEHB, 3. PEHA, 4. SNPA, 5. PEH, 6. SNPB) and Marker points (black) for the model organism taken in the plates]</p> <p>Figure 20. Antimicrobial activity analysis α-CD, β-CD on Gram-positive B. subtilis by Agar Cup method. No zone of inhibition was observed. Double distilled water was taken as the control. [Marker points (red) for the verified samples taken in the plates (1. SNP, 2. PEHB, 3. PEHA, 4. SNPA, 5. PEH, 6. SNPB) and Marker points (black) for the model organism taken in the plates]</p>	
<p>Chapter V</p>	<p>Figure 1: Molecular structures of (a) thiamine hydrochloride, (b) cyclodextrin</p> <p>Figure 2(a,b): Spectra for the generation of Job plot of (a) THC+α-CD and (b) THC+β-CD systems at 236 and 263 nm.</p> <p>Figure 3(a,b,c,d): Job plots of the THC+α-CD system at (a) λ_{\max}=236 nm, (b) λ_{\max}=263 nm and THC+β-CD system at (c) λ_{\max}=236 nm, (d) λ_{\max}=263 nm</p> <p>Figure 4. 2D ROESY NMR spectra of IC-1 (THC+α-CD)</p> <p>Figure 5. 2D ROESY NMR spectra of IC-2 (THC+β-CD)</p> <p>Figure 6: Plausible mechanism of inclusion complexation through wider rim of cyclodextrin.</p>	<p>164-172</p>

	<p>Figure 7(a,b,c): DSC thermos-gram of (a) THC, (b) THC+α-CD, (c) THC+β-CD</p> <p>Figure 8(a,b): ESI- Mass spectra of (a) IC-1 (THC+α-CD), (b) IC-2 (THC+β-CD)</p> <p>Figure 9(a,b,c): FTIR spectra of IC-1 (THC+α-CD), IC-2(THC+β-CD), THC, β-CD and α-CD</p> <p>Figure 10(a,b,c,d): SEM images of (a) α-CD, (b) THC, (c) (THC+α-CD) IC and (d) (THC+α-CD) physical mixture.</p> <p>Figure 11(e,f,g,h): SEM images of (e) α-CD, (f) THC, (g) (THC+α-CD) IC and (h) (THC+α-CD) physical mixture.</p> <p>Figure 12(a, b, c): Fluorescence spectra of HSA molecule in presence of (a) THC, (b) THC along with β-CD (c) THC along with α-CD at the various concentrations.</p>	
<p>Chapter VI</p>	<p>Figure 1. Molecular structures of (a) 1,2,5,6,9,10-hexabromocyclododecane (b) β-cyclodextrin and (c) hydroxypropyl-β-cyclodextrin.</p> <p>Figure 2. (a) 2D ROESY spectra of inclusion complex of HBCDD and β-CD in DMSO-d₆(correlation signals are marked by red circles) (b) 2D ROESY spectra of inclusion complex of HBCDD and HP-β-CD in DMSO-d₆(correlation signals are marked by red circles).</p> <p>Figure 3(a, b, c): FTIR spectra of (a) (HBCDD+β-CD) inclusion complex, (b) HBCDD, (c) β-CD.</p> <p>Figure 4(a, b, c): FTIR spectra of (a) (HBCDD+HP β-CD) inclusion complex, (b) HBCDD, (c) HP β-CD</p>	<p>186-198</p>

	<p>Figure 5. HRMS spectra of (a) HBCDD-β-CD inclusion complex and (b) HBCDD-HP-β-CD inclusion complex.</p> <p>Figure 6. Variation of surface tension with mole ratio of β-CD and HP-β-CD respectively.</p> <p>Figure 7. SEM images of (a) β-CD, (b) IC-1, (c) HBCDD, (d) HP-β-CD and (e) IC-2.</p> <p>Figure 8. EDX images of (a) β-CD, (b) IC-1, (c)HBCDD, (d) HP-β-CD and (e) IC-2</p> <p>Figure 9: DSC thermograms of (a) β-CD, (b) HP-β-CD, (c) HBCDD, (e) IC-1, (d) IC-2.</p> <p>Figure 10. Solubility of (a) HBCDD, (b) HBCDD-β-CD inclusion complex and (c) HBCDD-HP-β-CD inclusion complex in water.</p> <p>Figure 11. Representative TLC plats for degradation studies of IC-1 and IC-2.</p> <p>TLC-I: Rf (I-a): 0.77; Rf (I-b): 0.77, 0.86, 0.96; Rf (I-c): 0.77; Rf (I-d): 0.98.</p> <p>TLC-II: Rf (II-b): 0.21, 0.60, 0.96; Rf (II-c): 0.60; Rf (II-d): 0.98.</p> <p>Figure S1. ^1H NMR spectra of HBCD in DMSO-d_6 at 298.15 K</p> <p>Figure S2. ^1H NMR spectra of β-cyclodextrin in DMSO-d_6 at 298.15 K</p> <p>Figure S3. ^1H NMR spectra of HP-β-cyclodextrin in DMSO-d_6 at 298.15 K</p> <p>Figure S4. ^1H NMR spectra of IC-1 in DMSO-d_6 at 298.15 K</p>	
--	---	--

	Figure S5. ¹ H NMR spectra of IC-2 in DMSO-d ₆ at 298.15 K	
Chapter VII	<p>Figure 1: UV-vis spectra for the generation of Job's plot (a,c) ALVC-α-CD inclusion complexes and (b,d) ALVC-β-CD inclusion complexes.</p> <p>Figure 2: Mass spectra of (a) ALVC/α-CD inclusion complex, (b) ALVC/β-CD inclusion complex.</p> <p>Figure 3: 2D ROESY NMR spectra of ALVC/α-CD inclusion complex.</p> <p>Figure 4: 2D ROESY NMR spectra of ALVC/β-CD inclusion complex.</p> <p>Figure 5(a, b, c, d, e): IR frequencies of (a) ALVC, (b) α-CD, (c) β-CD, (d) ALVC/α-CD inclusion complex, (e) ALVC/β-CD inclusion complex.</p> <p>Figure 6(a, b, c, d, e): PXRD spectra of (a) ALVC, (b) α-CD, (c) β-CD, (d) ALVC/α-CD inclusion complex, (e) ALVC/β-CD inclusion complex.</p> <p>Figure 7(a, b, c): DSC thermogram of the (a) ALVC, (b) ALVC/α-CD inclusion complex, (c) ALVC/β-CD inclusion complex.</p> <p>Figure 8(a, b, c, d, e): SEM images of (a) α-CD, (b) β-CD, (c) ALVC, (d) ALVC/ α-CD inclusion complex, (e) ALVC/ β-CD inclusion complex.</p> <p>Figure 9: Fluorescence spectra and plot for obtaining binding constant.</p>	218-223
Chapter VIII	Figure 1: Graphical presentation of Well diffusion assay (Diameter of inhibition in mm) of Scyt and SBz, singly and in	264-271

	<p>combination with IL1 (BTEACl) and IL2 (BTMACl) applied to <i>B. subtilis</i> and <i>E. coli</i>.</p> <p>Figure 2: Well diffusion assay of Scyt and SBz, singly and in combination with IL1 (BTEACl) and IL2 (BTMACl).</p> <p>Figure 3: Well diffusion assay of Scyt and SBz, singly and in combination with IL1 (BTEACl) and IL2 (BTMACl).</p> <p>Figure 4. (a,b,c,d,e,f) Plot of (R_M) Vs \sqrt{c} for (BTEACl+SBz+H₂O), (BTEACl+SCyt+H₂O) systems in aqueous BTEACl solutions of mass fractions (a, b) $W_1=0.001$, (c, d) $W_2=0.003$, (e, f) $W_3=0.005$ respectively at 298.15 K, 303.15 K, 308.15 K, 313.15 K and 318.15 K.</p> <p>Figure 5. Variation of $\frac{dB}{dT}$ with the mass fraction of BTEACl and BTMACl separately at their four different combinations, BTEACl+SBz, BTEACl+Scyt, BTMACl+SBz, BTMACl+Scyt.</p> <p>Figure 6 (a, b, c, d) Spectra for the generation of Benesi-Hildebrand double reciprocal plot of (a) BTEACl+SBz, (b) BTEACl+Scyt systems and corresponding Benesi-Hildebrand double reciprocal plot of (c) BTEACl+SBz, (d) BTEACl+Scyt systems.</p> <p>Figure 7 (a, b, c, d) Spectra for the generation of Benesi-Hildebrand double reciprocal plot of (a) BTMACl+SBz, (b) BTMACl+Scyt systems and corresponding Benesi-Hildebrand double reciprocal plot of (c) BTMACl+SBz, (d) BTMACl+Scyt systems.</p> <p>Figure 8 (a,b): Optimised geometry of (a) BTMACl+SBz and (b) BTMACl+Scyt systems.</p>	
--	---	--