

## LIST OF TABLES

Table No.	Caption	Page No.
<b>Introduction</b>		
<b>Table 1.</b>	Commonly used saturated and unsaturated fatty acids.	2
<b>Chapter I</b>		
<b>Table 1.</b>	Iatrosan data of serum: the components present in the serum and their percent composition.	62
<b>Table 2.</b>	Summary of the equilibrium surface tension data for the adsorbed BLES films in the presence of different additives at 25 °C.	64
<b>Chapter II</b>		
<b>Table 1.</b>	Temperature for maximum heat flow ( $T_m$ ), the width at half peak height ( $\Delta T_{1/2}$ ), change in enthalpy ( $\Delta H$ ), heat capacity ( $\Delta C_p$ ) and percentage of crystallinity (C.I.) of blank as well as LIDO and PRO.HCl loaded NLC (5mM; Span 65+SLC+SA, 2:2:1 M/M/M).	103
<b>Table 2.</b>	Drug release kinetics profiles of LIDO and PRO.HCl loaded NLCs.	109
<b>Chapter III</b>		
<b>Table 1.</b>	Mean size ( $d_h$ / nm), PDI, zeta potential (mV), Entrapment Efficiency (EE %) and Loading Capacity (LC %) values of empty and ursolic acid - loaded NLC.	124
<b>Table 2.</b>	Temperature for maximum heat flow ( $T_m$ ), the width at half peak height ( $\Delta T_{1/2}$ ), change in enthalpy ( $\Delta H$ ), and heat capacity ( $\Delta C_p$ ) of blank as well as UA loaded NLC.	129
<b>Chapter IV</b>		

<b>Table 1.</b>	Photon correlation spectroscopy dimensional analysis, entrapment efficiency and drug loading capacity of blank as well as orcinol glucoside loaded non - PEGylated and PEGylated NLCs.	142
<b>Table 2.</b>	Temperature for maximum heat flow ( $T_m$ ), change in enthalpy ( $\Delta H$ ), the width at half peak height ( $\Delta T_{1/2}$ ), heat capacity ( $\Delta C_p$ ) and crystallinity index of orcinol glucoside loaded PEGylated and non - PEGylated NLCs.	150
<b>Table 3.</b>	Drug release kinetics profiles of orcinol glucoside loaded PEGylated NLCs and non -PEGylated NLCs.	154
<b>Table 4.</b>	IC <sub>50</sub> value of pure orcinol glucoside (OG), OG loaded non-PEGylated NLC (OG-NLC), PEG-25-SA coated OG loaded NLC (OG-P <sub>1</sub> -NLC) and PEG-55-SA coated OG loaded NLC (OG-P <sub>2</sub> -NLC) for Hepatocellular carcinoma cell line (HepG2), hepatocyte-derived carcinoma cell line (Huh-7), colorectal carcinoma cell line (HCT-116) and gastric adenocarcinoma (AGS) cells.	155

---