

List of figures

Chapter 1

Figure 1.1	<i>Molecular configuration of nematic, smectic A and smectic C phases.</i>	4
Figure 1.2	<i>Symmetries in SmC and SmC* phases</i>	7
Figure 1.3	<i>Bistable rotation of molecule in SmC* phase</i>	8
Figure 1.4	<i>Helical structure in SmC* phase</i>	9
Figure 1.5	<i>Helical structure in SmC*_A phase</i>	10
Figure 1.6	<i>Nonhelical local model of chiral smectic phases</i>	11

Chapter 2

Figure 2.1	<i>Schematic diagram of a) homogeneously and b) homeotropically aligned cell.</i>	32
Figure 2.2	<i>Schematic diagram of ideal SSFLC cell</i>	34
Figure 2.3	<i>The experimental setup for polarizing microscopy</i>	35
Figure 2.4	<i>Typical dielectric spectra of Debye type dielectric material</i>	38
Figure 2.5	<i>Cole-Cole plot for Debye type spectra</i>	39
Figure 2.6	<i>Cole -Cole type spectra</i>	41
Figure 2.7	<i>A schematic diagram of both GM and sm relaxation observed together</i>	43
Figure 2.8	<i>Theoretically predicted temperature variation for different β of a) dielectric increment of GM b) dielectric increment of SM c) critical frequency of GM d) critical frequency of SM.</i>	44
Figure 2.9	<i>Experimental setup for dielectric measurement</i>	46
Figure 2.10	<i>Schematic diagram of X-ray diffraction photographs of a) SmC phase and b) SmA phase</i>	47
Figure 2.11	<i>Observed X-ray diffraction photograph of a) SmC* and b) SmA* phase of a chiral liquid crystal mixture.</i>	48
Figure 2.12	<i>Fitted intensity profile of SAXS data a) at low and b) high angle.</i>	49
Figure 2.13	<i>Experimental set up for X-ray scattering experiment.</i>	51
Figure 2.14	<i>Schematic diagram of the experimental setup to</i>	

	<i>measure spontaneous polarization and response time.</i>	53
Figure 2.15	<i>Hysteresis loop of P_s at different phases.</i>	55
Figure 2.16	<i>Polarization bump in SmC^* phase undersquare pulse.</i>	56
Chapter 3		
Figure 3.1	<i>Molecular structure of 1F3R</i>	67
Figure 3.2	<i>DSC thermogram of 1F3R in heating and cooling cycle</i>	70
Figure 3.3	<i>Observed textures of 1F3R at a) SmA^* (143°C) b) SmA^* during heating (130°C) c) SmA^* to SmX^* transition (127°C), d) SmX^* (125°C), e) SmB^* (112°C) and e) SmE^* (91°C).</i>	71
Figure 3.4	<i>Optimised geometry of 1F3R</i>	72
Figure 3.5	<i>Diffraction photographs of (a) SmE^* (89.5°C) (b) $SmB^*_{hex}{}^{mo}$ (107°C) (c) $SmB^*_{hex}{}^{lo}$ (SmX^*) (125°C)(d) SmA^*(140°C) phases</i>	73
Figure 3.6	<i>Intensity profile of inner ring(a) SmE^*(89.5°C), (b) $SmB^*_{hex}{}^{mo}$ (107°C) (c) $SmB^*_{hex}{}^{lo}$ (125°C) and (d)SmA^*(140°C).</i>	74
Figure 3.7	<i>Intensity profile of outer ring (a) SmE^*(89.5°C), (b) $SmB^*_{hex}{}^{mo}$ (107°C) (c) $SmB^*_{hex}{}^{lo}$ (125°C) and (d)SmA^*(140°C).</i>	75
Figure 3.8	<i>Intensity of the first order (a) inner and (b) outer diffraction peaks with temperature.</i>	76
Figure 3.9	<i>Layer spacing of all phases as a function of temperature.</i>	77
Figure 3.10	<i>Variation of average intermolecular spacing (D) with temperature</i>	77
Figure 3.11	<i>Temperature variation of correlation length (a) across the layer and (b) within the layer</i>	79
Figure 3.12	<i>Hexagonal lattice with herringbone structure</i>	80
Figure 3.13	<i>Temperature dependence of real part of dielectric permittivity ϵ'_{\perp} at 100 Hz.</i>	82
Figure 3.14	<i>Fitted spectra of (a) MW mode in SmB^* phase at 105°C and of (b) SA mode in SmE^* phase at 81°C.</i>	83
Figure 3.15	<i>Fitted Cole-Cole plot of (a) MW mode in SmB^* phase at 105°C and of (b) SA mode in SmE^* phase at 81°C</i>	84
Figure 3.16	<i>Temperature variation of (a) critical frequency</i>	86

	<i>and (b) dielectric increment.</i>	
Figure 3.17	<i>Real parts of the dielectric spectra in different phases</i>	87
Figure 3.18	<i>Imaginary parts of the dielectric spectra in different phases</i>	88
Figure 3.19	<i>Arrhenius plot for 1F3R</i>	88
 Chapter 4		
Figure 4.1	<i>Molecular structures of (a) 2F3R and (b) 3F3R</i>	96
Figure 4.2	<i>Optimised geometry of a) 2F3R and b) 3F3R</i>	99
Figure 4.3	<i>Textures of 2F3R at a) SmJ* (80°C) b) SmF* (84°C) c) SmC* (125°C) and d) SmA* (132°C).</i>	101
Figure 4.4	<i>Textures of 3F3R at a) SmG* (70°C) b) SmF* (74°C) and c) SmC* (130°C).</i>	101
Figure 4.5	<i>Schematic diagram of the structures and X-ray patterns of SmF* and SmI* phases.</i>	103
Figure 4.6	<i>Diffraction pattern of 2F3R at a) SmJ* (80°C), b) SmF* (84°C) and c) mixed phase (100°C).</i>	104
Figure 4.7	<i>Diffraction pattern of 3F3R at a) SmG* (70°C), b) SmF* (75°C) and c) mixed phase (83°C).</i>	104
Figure 4.8	<i>Intensity profiles of outer maxima in different phases of a) 2F3R and b) 3F3R.</i>	105
Figure 4.9	<i>Intensity profile of inner maxima of a) 2F3R and b) 3F3R</i>	107
Figure 4.10	<i>Temperature variation of correlation lengths of 2F3R a) across and b) within the layer.</i>	108
Figure 4.11	<i>Temperature variation of correlation lengths of 3F3R a) across and b) within the layer.</i>	108
Figure 4.12	<i>Temperature variation of layer spacing of a) 2F3R and b) 3F3R</i>	110
Figure 4.13	<i>Temperature variation of X-ray tilt angle of a) 2F3R and b) 3F3R</i>	111
Figure 4.14	<i>Temperature variation of average intermolecular spacings (D) of a) 2F3R and b) 3F3R</i>	112
Figure 4.15	<i>Fitted dielectric spectra showing a) SM relaxation in SmA* phase of 2F3R and b) MW, BOO and tilt phason relaxations in SmF* phase of 3F3R.</i>	115
Figure 4.16	<i>Cole-Cole plots of dielectric spectra of a) 2F3R at 134°C and b) 3F3R at 74°</i>	115

Figure 4.17	<i>Temperature variation of dielectric increment of a) 2F3R and b) 3F3R.</i>	116
Figure 4.18	<i>Temperature variation of Critical frequency of a) 2F3R and b) 3F3R.</i>	116
Figure 4.19	<i>Temperature variations of spontaneous polarizations of a) 2F3R and b) 3F3R.</i>	118
Figure 4.20	<i>Temperature variation of rotational viscosities of 2F3R and 3F3R.</i>	119
Figure 4.21	<i>Temperature variation of response time of 2F3R and 3F3R</i>	120
Figure 4.22	<i>Temperature variation of helical pitch of a) 2F3R and b) 3F3R</i>	121
Figure 4.23	<i>Temperature variations of elastic constants of a) 2F3R and b) 3F3R</i>	122
 Chapter 5		
Figure 5.1	<i>Molecular structure of a) 5F3R and b) 6F3R</i>	132
Figure 5.2	<i>Optimised geometry of a) 5F3R and b) 6F3R</i>	134
Figure 5.3	<i>Optical texture of 5F3R at a) Crystalline state b) SmC* phase</i>	135
Figure 5.4	<i>Fitted spectra with fitted parameters of a) 5F3R (95°C) and b) 6F3R (86°C).</i>	137
Figure 5.5	<i>Fitted Cole-Cole plot of a) 5F3R (95°C) and b) 6F3R (86°C)</i>	137
Figure 5.6	<i>Temperature dependence of critical frequency of a) 5F3R and b) 6F3R</i>	138
Figure 5.7	<i>Temperature variation of dielectric increment of a) 5F3R and b) 6F3R</i>	138
Figure 5.8	<i>Temperature variation of spontaneous polarization of a) 5F3R and b) 6F3R</i>	142
Figure 5.9	<i>Temperature variation of response time of a) 5F3R and b) 6F3R</i>	142
Figure 5.10	<i>Temperature variation of optical tilt angle of a) 5F3R and b) 6F3R</i>	144
Figure 5.11	<i>Temperature variation of rotational viscosity of a) 5F3R and b) 6F3R</i>	146

Figure 5.12	<i>Temperature variation of rotational viscosity measured both by electro-optic method and dielectric method of a) 5F3R and b) 6F3R</i>	146
--------------------	---	------------

Chapter 6

Figure 6.1	<i>The molecular structures of a) 4F4R and b) 4F5R</i>	154
Figure 6.2	<i>Optimised geometry of a) 4F4R and b) 4F5R</i>	157
Figure 6.3	<i>Optical textures of 4F4R in a) crystal b) SmX* and c) SmC* phases</i>	158
Figure 6.4	<i>Optical textures of 4F5R in a) crystal and b) SmC* phase</i>	158
Figure 6.5	<i>Variation of real part of dielectric permittivity of 4F4R with temperature (a) at 1kHz, (b) at 10 kHz</i>	160
Figure 6.6	<i>Fitted dielectric spectra for 4F4R showing GM: (a) SmX* phase (96°C), (b) SmC* phase (110°C)</i>	160
Figure 6.7	<i>Fitted dielectric spectra in SmC* phase at 115°C for 4F5R showing GM.</i>	161
Figure 6.8	<i>Fitted cole cole plots of a) 4F4R in SmX* phase (96°C), b) SmC* phase (110°C) and c) 4F5R in SmC* phase (115°C)</i>	162
Figure 6.9	<i>Temperature variations of (a) critical frequency and (b) dielectric increment of 4F4R.</i>	162
Figure 6.10	<i>Temperature variations of (a) critical frequency and (b) dielectric increment of 4F5R.</i>	162
Figure 6.11	<i>Dependence of Ps on temperature (a) 4F4R, (b) 4F5R</i>	164
Figure 6.12	<i>Rotational viscosity (γ_ϕ) of a) 4F4R and b) 4F5R.</i>	166
Figure 6.13	<i>Optical tilt angles of a) 4F4R and b) 4F5R</i>	167
Figure 6.14	<i>Response time of a) 4F4R and b) 4F5R</i>	168

Chapter 7

Figure 7.1	<i>Molecular structure of 7F3R</i>	174
Figure 7.2	<i>Optical texture of 7F3R at a) crystal (74°C) b) SmC* (114°C) and c) SmA* (149°C) phases</i>	177
Figure 7.3	<i>Optimised geometry of 7F3R</i>	178
Figure 7.4	<i>Diffraction photographs of 7F3R at a) 122°C (SmC*) and b) 149°C (SmA*)</i>	179

Figure 7.5	<i>Temperature variation of layer spacing and average intermolecular spacing of 7F3R</i>	180
Figure 7.6	<i>Temperature variation of d_C/d_A near transition</i>	181
Figure 7.7	<i>Temperature variation of X-ray tilt angle of 7F3R</i>	182
Figure 7.8	<i>Correlation lengths of 7F3R a) across the layer and b) within layer</i>	183
Figure 7.9	<i>Fitted dielectric spectra in SmC* phase at 147.7°C showing both GM and SM.</i>	184
Figure 7.10	<i>Fitted cole-cole plot at 147.7°C</i>	185
Figure 7.11	<i>Temperature variation of a) dielectric increment and b) critical frequency of 7F3R</i>	185
Figure 7.12	<i>Temperature variation of spontaneous polarisation of 7F3R.</i>	186
Figure 7.13	<i>Temperature variation of rotational viscosity and response time</i>	187
Figure 7.14	<i>Temperature variation of optical tilt angle</i>	188
Figure 7.15	<i>DSC thermogram of the mixture during cooling</i>	192
Figure 7.16	<i>Textures of the mixture at temperature a) 33°C (SmC*) b) 63°C (SmC* with changed birefringence) c) 85°C (SmA*) and d) 93°C (SmA* with isotropic domain).</i>	193
Figure 7.17	<i>Diffraction photograph of HM at SmC (60°C)</i>	194
Figure 7.18	<i>Diffraction photograph of the mixture at a) SmC* (66°C) and b) SmA* (84°C) phases.</i>	194
Figure 7.19	<i>The intensity profile of inner maxima of X-ray photograph at a) SmC* (66°C) and b) SmA* (84°C) phases.</i>	195
Figure 7.20	<i>The intensity profile of outer maxima of X-ray photograph at a) SmC* (66°C) and b) SmA* (84°C) phases.</i>	195
Figure 7.21	<i>Temperature variation of a) layer spacing and b) tilt angle of HM, H1, H2, H3 and H4.</i>	195
Figure 7.22	<i>Temperature variation of a) layer spacing and b) tilt angle of the mixture, HM and 7F3R.</i>	196
Figure 7.23	<i>Temperature variation of layer spacing of the mixture</i>	197
Figure 7.24	<i>Temperature variation of X-ray tilt of mixture with and without bias.</i>	199
Figure 7.25	<i>Correlation lengths of the mixture and the dopant a) across the layers and b) within the layers.</i>	200
Figure 7.26	<i>Temperature variation of P_S of the mixture.</i>	202

Figure 7.27	<i>Field variation of P_s in a) SmC^* and b) SmA^* phases.</i>	203
Figure 7.28	<i>Dependence of effective dipole moment on applied field in SmA^* phase (77°C)</i>	203
Figure 7.29	<i>Temperature variation of optical tilt of the mixture</i>	204
Figure 7.30	<i>Bias dependence of optical tilt of the mixture at temperature 89°C (SmA^*).</i>	205
Figure 7.31	<i>The X-ray and optical tilt of mixture and 7F3R.</i>	205
Figure 7.32	<i>Temperature variation of rise time of the mixture.</i>	206

Chapter 8

Figure 8.1	<i>a) Optimised molecular length and b) Dipole moments of compounds $nFmR$. For 4F4R and 4F5R, number of oligomethylene spacers is plotted along x-axis.</i>	215
Figure 8.2	<i>Span of SmC^* phase of different compounds. For 4F4R and 4F5R, number of oligomethylene spacers is plotted along x-axis.</i>	217
Figure 8.3	<i>Variation of Goldstone mode a) dielectric increment ($\Delta\epsilon$) and b) critical frequency (f_c) of $nFmR$. For 4F4R and 4F5R, number of oligomethylene spacers is plotted along x-axis.</i>	220
Figure 8.4	<i>Highest value of spontaneous polarization of all compounds having SmC^* phase.</i>	221
Figure 8.5	<i>Highest value of response time of compounds $nFmR$. For 4F4R and 4F5R, number of oligomethylene spacers is plotted along xaxis</i>	222