

List of Figures

1.1	Luminosity distance vs. cosmological redshift variation curve for type Ia Supernovae	10
1.2	Flat rotation curve of a spiral galaxy	11
3.1	Geometrical configuration of time delay/advancement of photon/-particle in gravitational field. O is the Centre of the spherically symmetric mass distribution, A and B are two arbitrary points. r_A and r_B are radial distances of X and Y from O respectively.	56
4.1	Lensing diagram - the source S emits light rays which reach the observer O after being gravitationally deflected by the lens L. . . .	74
4.2	Lensing diagram - the source S emits light rays which reach the observer O after being gravitationally deflected by the lens L which is an isolated global monopole. The image position is denoted as I. . . .	78
4.3	Magnification in weak gravitational lensing - the solid line and dotted line respectively denote μ_+ and μ_- due to lensing by a Schwarzschild black hole that swallowed a global monopole, the dashed line and long dashed line respectively denote μ_+ and μ_- due to lensing by a Schwarzschild black hole.	82
4.4	Variation of deflection angle with mass due to a pure Schwarzschild model (dotted line) and a global monopole swallowed Schwarzschild model (solid line). The closest distance parameter is taken 100 kpc.	85
5.1	The variation of β_ϕ^4 with M_B for several galaxies. The dotted line represents the least squared fit of the data.	99
6.1	Variation of observed total baryonic mass with rotation velocity. The filled (blue) circle represent the observed data, solid (black) line gives the fitting of the data for standard Rindler acceleration (power index fixed at 4) and the dotted (red) line shows the fitting of the data with generalized Rindler acceleration under Grumiller's modified gravity model.	105
6.2	Frequency distribution of estimated Rindler acceleration parameter.	107