

LIST OF FIGURES

Figure No.	Title	Page No.
Figure I.A.1	Model of a C ₆₀ buckyball	3
Figure I.A.2	Structure of monolayer graphene, graphene oxide and reduced graphene oxide	4
Figure I.A.3	Molecular representations of (a) SWCNT and (b) MWCNT	5
Figure I.A.4	Schematic illustration for the synthesis of graphene by ‘top-down’ and ‘bottom-up’ approaches	6
Figure I.A.5	Different routes for the functionalization of graphene	7
Figure I.A.6	Different structural models of GO	14
Figure I.B.1	Bioactive sulfur functionalized indoles	23
Figure I.B.2	Recyclability of GO for the sulfenylation of indole	35
Figure I.B.3	FT-IR spectra of GO before and after first, second and fourth run	35
Figure I.B.4	(a) XRD patterns and (b) Raman spectra of GO fresh and after the first run	36
Figure I.B.5	The pH potentiometric titration curve of GO fresh and after the first run	36
Figure I.B.6	Scanned copy of ¹ H NMR spectrum of 3-(pentylthio)-1 <i>H</i> -indole (3f)	48
Figure I.B.7	Scanned copy of ¹³ C NMR spectrum of 3-(pentylthio)-1 <i>H</i> -indole (3f)	48
Figure I.B.8	Scanned copy of HRMS spectrum 3-(pentylthio)-1 <i>H</i> -indole (3f)	49
Figure I.C.1	Examples of clinical drugs bearing functionalized 1,4-dihydropyridine unit	54
Figure I.C.2	FT-IR spectra of GO and AFGONs	61
Figure I.C.3	Raman spectra of GO and AFGONs	62
Figure I.C.4	X-ray diffraction patterns of GO and AFGONs	62
Figure I.C.5	SEM images of (a) GO and (b) AFGONs	63
Figure I.C.6	EDS images of (a) GO and (b) AFGONs	63
Figure I.C.7	Recyclability of AFGONs in the synthesis of 1,4-dihydropyridines	71
Figure I.C.8	(a) FT-IR and (b) Raman spectra of AFGONs fresh and after the third run	71
Figure I.C.9	X-ray diffraction patterns of AFGONs fresh, after first and third runs	71
Figure I.C.10	SEM images AFGONs (a) fresh and (b) after the third run	72

Figure I.C.11	EDS images of AFGONs (a) fresh and (b) third run	72
Figure I.C.12	Scanned copy of ¹ H NMR spectra of methyl 4-(4-isopropylphenyl)-2,7,7-trimethyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carboxylate (6c)	89
Figure I.C.13	Scanned copy of ¹³ C NMR spectra of methyl 4-(4-isopropylphenyl)-2,7,7-trimethyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carboxylate (6c)	90
Figure I.C.14	Scanned copy of HRMS spectra of methyl 4-(4-isopropylphenyl)-2,7,7-trimethyl-5-oxo-1,4,5,6,7,8-hexahydroquinoline-3-carboxylate (6c)	90
Figure I.D.1	Schematic diagram of a continuous flow set-up	95
Figure I.D.2	Examples of active pharmaceutical ingredients synthesized via flow chemistry	96
Figure I.D.3	Functionalized thioethers possessing bioactive properties	97
Figure I.D.4	Experimental setup for continuous flow two-step synthesis of thioether: (a) schematic, (b) digital image	102
Figure I.D.5	Recyclability of GO flow reactor bed in Stadler-Ziegler reaction	107
Figure I.D.6	FT-IR spectra of GO catalyst: fresh and after the fifth run	108
Figure I.D.7	Scanned copy of ¹ H NMR spectrum of (4-chlorophenyl)(4-methoxyphenyl)sulfane (3c)	114
Figure I.D.8	Scanned copy of ¹³ C NMR spectrum of (4-chlorophenyl)(4-methoxyphenyl)sulfane (3c)	114
Figure II.B.1	Isomeric and tautomeric forms of various triazoles	125
Figure II.B.2	Biologically active triazoles and their potential activities	126
Figure II.B.3	Clinical drugs based on triazole unit	126
Figure II.B.4	Typical example of a triazole based dendrimer	127
Figure II.B.5	FT-IR spectra of GO, NaY zeolite, GO-zeolite and Ni-rGO-zeolite	136
Figure II.B.6	Raman spectra of Ni-rGO-zeolite catalyst and GO-zeolite (as control)	137
Figure II.B.7	XRD patterns of GO-zeolite and Ni-rGO-zeolite composites	137
Figure II.B.8.1	XPS survey scan of Ni-rGO-zeolite catalyst	139
Figure II.B.8.2	XPS spectra of Ni-rGO-zeolite composite: (a) high resolution C1s spectrum (b) high resolution Ni2p spectrum	139
Figure II.B.9	(a) SEM image of GO-zeolite, (b) SEM image of Ni-rGO-zeolite, (c) EDS of Ni-rGO-zeolite and (d) TEM image of Ni-rGO-zeolite	140
Figure II.B.10	Recyclability of Ni-rGO-zeolite in the click synthesis of triazoles	145

Figure II.B.11	XPS analysis of Ni-rGO-zeolite after first run: (a) survey scan (b) high resolution Ni2p spectrum	145
Figure II.B.12	Scanned copy of ¹ H NMR spectrum of 1-benzyl-4-phenyl-1 <i>H</i> -1,2,3-triazole (3a)	154
Figure II.B.13	Scanned copy of ¹³ C NMR spectrum of 1-benzyl-4-phenyl-1 <i>H</i> -1,2,3-triazole (3a)	154
Figure II.C.1	FT-IR spectra of GO, GO-SiO ₂ and Cu@GO-SiO ₂	161
Figure II.C.2	PXRD patterns of GO-SiO ₂ and Cu@GO-SiO ₂	162
Figure II.C.3	Raman spectra of GO-SiO ₂ and Cu@GO-SiO ₂	162
Figure II.C.4	(a-d) HR-TEM images and (e) EDS image of Cu@GO-SiO ₂	163
Figure II.C.5	XPS analysis of Cu@GO-SiO ₂ nanocomposite	164
Figure II.C.6	Recyclability of Cu@GO-SiO ₂ nanocomposite in C-S cross-coupling reaction	173
Figure II.C.7	Scanned copy of ¹ H NMR spectrum of 1-nitro-3-(<i>p</i> -toloxy)benzene (7c)	182
Figure II.C.8	Scanned copy of ¹³ C NMR spectrum of 1-nitro-3-(<i>p</i> -toloxy)benzene (7c)	182

