

List of Figures

<u>Figures</u>	<u>Page No.</u>
 <u>Materials and Methods</u>	
Figure 3.1. Detailed map of the rivers of the study area with the sampling sites marked on it.	57
Figure 3.2. Map of the study area with the hospitals marked on it whose effluents was screened for the study.	58
Figure 3.3. Map of the study area with the towns marked on it whose municipality drinking water and domestic effluents was screened for the study.	59
 <u>Results and Discussion</u>	
Figure 4.1. Some sample collection sites in Rivers Chenga, Tepu, Balason, Panchanoi and Mahananda in different administrative blocks, districts and town of North Bengal. GPS locations are depicted within parentheses.	102
Figure 4.2. Some sample collection sites in Rivers Mahananda and Teesta in different administrative blocks, districts and towns of North Bengal and Siliguri government hospital. GPS locations are depicted within parentheses.	103
Figure 4.3. Some sample collection sites in Rivers Karala and Ghish in different administrative blocks of Jalpaiguri District of North Bengal and Rajbari dighi at Jalpaiguri town. GPS locations are depicted within parentheses.	104
Figure 4.4. Growth of bacteria obtained after growing water samples after collecting from different locations and sites (river Chenga, Naxalbari-Matigara block and North Bengal Medical College sewage	110

line, Siliguri), spread in different dilutions on *Aeromonas* Selective media. Dilutions used: Raw, 1/10 dilution, 1/100 dilution and 1/1000 dilution.

Figure 4.5. Growth of bacteria obtained after growing water samples after collecting from different locations and sites (North Bengal Medical College sewage line and river Balason, Palpara), spread in different dilutions on *Aeromonas* Selective media. Dilutions used: Raw, 1/10 dilution, 1/100 dilution and 1/1000 dilution. **111**

Figure 4.6. Growth of bacteria obtained after growing water samples after collecting from different locations and sites (river Balason, Palpara and river Chenga Naxalbari-Matigara block), spread in different dilutions on *Aeromonas* Selective media. Dilutions used: Raw, 1/10 dilution, 1/100 dilution and 1/1000 dilution. **112**

Figure 4.7. Representative *Aeromonas* starch DNA agar plates showing isolated single colonies of *Aeromonas* sp. obtained from hospital effluents. **113**

Figure 4.8. Representative Gram-stained slides of *Aeromonas* sp isolated from rivers and hospital effluents observed under oil immersion at 100 X magnification. **114**

Figure 4.9. Catalase test. Bubble formation indicates positive result; absence of bubbles indicates negative result. Black arrows indicate the formed bubbles. **116**

Figure 4.10. Starch hydrolysis test. Clear zones around the bacterial growth indicate the positive results; absence of clear zone indicates negative result. Arrow heads indicate clear zones formed. **117**

Figure 4.11. Gelatinase Test. Clear zones around the bacterial growth indicate positive results; absence of clear zones indicate negative results. Black arrow-heads in the lower panel indicate clear zones formed. **118**

Figure 4.12. Glucose fermentation Test; Yellow colourations indicate positive results; absence of yellow colours indicates negative results. Black arrow heads indicate occurrence of fermentation. **119**

Figure 4.13. Indole test. Formation of rosy-red rings indicate positive results; absence of rosy-red rings indicates negative result. Black arrow-heads indicate formation of indole.	120
Figure 4.14. Methyl Red test. Formation of red colouration indicates positive result; absence of red colour indicates negative result. Black arrow-heads indicate formation of acid.	121
Figure 4.15. Voges Proskauer test. Formation of red colouration indicates positive result; absence of red colour indicates negative result. Black arrow-heads indicate formation of the red coloured end-product of the reaction.	122
Figure 4.16. Oxidase test. Formation of blue colouration indicates positive result; absence of blue colour indicates negative result. Black arrow-heads indicate production of Cytochrome c oxidase enzyme by the bacteria.	123
Figure 4.17. Citrate test. Formation of blue colouration indicates positive result; absence of red colour indicates negative result. Black arrow-heads indicate production of citrase enzyme by the bacteria.	124
Figure 4.18. Lipase test. Clear zone around bacterial growth indicates positive result; absence of clear zone indicates negative result. Black brrow heads indicate clear zones formed around the bacterial isolates due to hydrolysis of lipid by the enzyme lipase produced by the bacterial isolates.	125
Figure 4.19. Representative agarose gels showing the isolated genomic DNA of <i>Aeromonas</i> isolates.	129
Figure 4.20. 16S rDNA Polymerase Chain Reaction. A. Lanes M: 100 bp ladder; 1 to 6: PCR product (953 bp). B. Lanes M: 100 bp ladder; 1 to 9: PCR product (953 bp). C. Lanes M: 100 bp ladder; 1 and 3 to 20: PCR product (953 bp). White arrow-heads indicate 953 bp PCR products.	131
Figure 4.21. Restriction fragment length polymorphism of 16S rRNA gene amplification product; A. Lanes M: 100 bp ladder; 1: uncut PCR product (953 bp); 2: AluI digest (361, 211,177 and 162 bp). B. Lanes	134

M: 100 bp ladder; 1: uncut PCR product; 2: AluI digest (361, 211, 177 and 162 bp); 3: AluI+MboI digest (346, 195, 165, 54 and 33 bp). C. Lane M: 100 bp ladder; lanes 1, 2, 3, 5 and 8: AluI+MboI (195, 165 and 157 bp); lane 4: AluI+MboI digest (346, 165, 157 and 138 bp); lane 6: AluI+MboI digest (165 and 157 bp); lanes 7 and 9: AluI+MboI digest (346, 165 and 157 bp); lane 10: Uncut PCR product (953 bp). White arrow heads indicate DNA bands.

Figure 4.22. Restriction fragment length polymorphism of 16S rRNA gene; D. Lanes M: 100 bp Ladder; Lanes 1 and 5: uncut PCR product (953 bp); lanes 2 and 6: AluI digests (211, 188, 177 and 173 bp); lanes 3 and 7: MboI digests (400, 350 and 200 bp); lanes 4 and 8: AluI+MboI digests (188, 177 and 173 bp); white arrow heads indicate DNA bands. **135**

Figure 4.23. Molecular Phylogenetic analysis based on *Aeromonas* rDNA sequences by Maximum Likelihood method. The tree is based on a comparison of 953 nucleotides using the neighbour-joining method. Sequences enclosed in red boxes are of the *Aeromonas* isolates found during the study. **142**

Figure 4.24. Mueller Hinton agar plates showing resistance-vs-susceptibility (clear zones) patterns in different *Aeromonas* natural isolates (AR7, AR10, AR12 and AD1 in plates A, B, C and D respectively) against ampicillin (25 µg) (1), nitrofurantoin (50 µg) (2), ticarcillin (75 µg) (3), tetracycline (100 µg) (4), nalidixic acid (30 µg) (5), trimethoprim (2.5 µg) (6), sulphamethoxazole (50 µg) (7) and gentamycin (10 µg) (8) (Himedia UTI-VII). Clear zones indicate zones of inhibition (black arrow heads) against a particular antibiotic; absence of clear zone was observed in cases of isolates (white arrow heads) resistant to a particular antibiotic. Refer to **Table 4.4** for a complete picture of antibiotic resistance/susceptibility shown by different natural *Aeromonas* isolates. **145**

Figure 4.25. Mueller Hinton agar plates showing resistance-vs-susceptibility (clear zones) pattern in different *Aeromonas* natural **146**

isolates [AR8 (plates A and C), AR9 (plate B) and AH1 (plate D)] against [cefotaxime (30 µg) (1), nalidixic acid (30 µg) (2), nitrofurantoin (300 µg) (3), norfloxacin (10 µg) (4), netillin (30 µg) (5), ofloxacin (5 µg) (6), ceftazidime (30 µg) (7) and ciprofloxacin (5 µg) (8) (Himedia G-XVIII-minus)], [ceftazidime (30 µg) (1), azithromycin (15 µg) (2), ampicillin/sulbactam (10/10 µg) (3), cefoperazone/sulbactam (75/30 µg) (4), cefepime (30 µg) (5), cefpirome (30 µg) (6), amikacin (30 µg) (7) and amoxyclav (30 µg) (8) (Himedia Combi 94)] and [amoxyclav (30 µg) (1), gentamycin (10 µg) (2), co-trimoxazole (25 µg) (3), norfloxacin (10 µg) (4), oxytetracycline (30 µg) (5) and cefuroxime (30 µg) (6) (HimediaHexa UTI-I)]. Clear zones indicate zones of inhibition (black arrow heads) against a particular antibiotic; absence of clear zone was observed in cases of isolates (black arrow heads) resistant to a particular antibiotic. The **Table 4.4** presents a complete picture of antibiotic resistance/susceptibility as shown by different natural *Aeromonas* isolates from the study region.

Figure 4.26. Mueller Hinton agar plates showing resistance-vs-susceptibility (clear zones) patterns in different *Aeromonas* natural isolates [AR6 (plates B and D) and AR9 (plates A and C)] against [co-trimoxazole (25 µg) (1), levofloxacin (5 µg) (2), nitrofurantoin (50 µg) (3), ampicillin (10 µg) (4), cefazolin (30 µg) (5), nalidixic acid (30 µg) (6), norfloxacin (10 µg) (7) and ciprofloxacin (5 µg) (8) (Himedia Combi 71)], [cephalothin (30 µg) (1), chloramphenicol (30 µg) (2), clindamycin (2 µg) (3), erythromycin (15 µg) (4), gentamycin (10 µg) (5), oxacillin (1 µg) (6), vancomycin (30 µg) (7) and ampicillin (10 µg) (8) (Himedia Combi IV)] and [streptomycin (10 µg) (1), chloramphenicol (30 µg) (2), ampicillin (10 µg) (3), tetracycline (30 µg) (4), gentamycin (10 µg) (5), kanamycin (30 µg) (6), co-trimoxazole (25 µg) (7), amikacin (30 µg) (8) (Himedia G-XIX-minus)]. Clear zones indicate zones of inhibition (black arrow heads) against a particular antibiotic; absence of clear zone was

observed in cases of isolates (white arrow heads) resistant to a particular antibiotic. Refer to **Table 4.4** for a complete picture of antibiotic resistance/susceptibility shown by different natural *Aeromonas* isolates.

Figure 4.27. Natural plasmids obtained from *Aeromonas* isolates. **A.** 166
Lanes M1: supercoiled DNA ladder; lane 1: Natural plasmids (8066, 2600 and 2000 bp approximately indicated by white arrow heads). **B.**
Lane M1: Supercoiled DNA ladder; lanes 1 and 2: natural plasmids (10102, 6030, 2972, 2067 and 1900 bp approximately; indicated by white arrow heads); M2: Lambda HindIII digest.

Figure 4.28. Restriction endonuclease digestion of natural plasmids 169
obtained from *Aeromonas* isolates. **A.** Lane M1: Supercoiled plasmid DNA ladder; lane 1: uncut plasmid; black arrow indicates a plasmid of 12138 bp; lane 2: EcoRI digest; lane 3: BamHI digest; lane 4: HindIII digest; lane M2: Lambda HindIII digest. **B.** Lane M2: Lambda HindIII digest; lanes 1 and 3: Uncut natural plasmids; lanes 2 and 4: Eco RI digest (9416, 5500, 3500, 2027 bp and another less than 2 kb).

Figure 4.29. Ampicillin plates showing the growth of transformed *E. coli* JM109 colonies; **A:** experimental plates and negative control. **B:** 172
negative and positive controls.

Figure 4.30. Ampicillin plates showing the growth of transformed 173
bacterial colonies; C, D and E: experimental plates; F: negative control; G and H: positive control.

Figure 4.31. Plasmids obtained from *E. coli* previously transformed 175
with natural plasmids, isolated from *Aeromonas* and their restriction digestion patterns. **A.** Lanes 1 to 5: EcoRI digests (23130, 6557 and 4361 approximately indicated by white arrow heads); lanes 6, 7 to 10: PstI digests (23130, 6557, 4361 and 2686 bp approximately indicated by white arrow heads); lane 11: pBR322 (uncut); lane 12: pUC19 (uncut); lane 13: pBR322-EcoRI digest; lane 14: pBR322-PstI digest; lane 15: pUC19-EcoRI digest; lane 16: pUC19-PstI digest; lane 17:

Lambda HindIII digest. **B.** Lane 1: supercoiled DNA ladder; lanes 2 to 15: PstI digests (14174, 7500 and 4361 bp approximately indicated by white arrow heads); lane 16: pBR322-PstI digest; lane 17: pUC 19-PstI digest.

Figure 4.32. Mueller Hinton agar plates showing antibiotic resistance transfer from *Aeromonas* to *E. coli* through transformation; **A:** *Aeromonas*; **B:** *E. coli* (JM109) transformed with plasmid from A; 1: Oxytetracycline (30 µg); 2: Cefuroxime (30 µg); 3: Amoxyclav (30 µg); 4: Gentamycin (10 µg); 5: Co-trimoxazole (25 µg); 6: Norfloxacin (10µg); resistance to amoxyclav got transferred from *Aeromonas* to *E. coli* (JM109). **178**

Figure 4.33. Mueller Hinton agar plates showing antibiotic resistance transfer from *Aeromonas* to *E. coli* through transformation; **A:** *Aeromonas*; **B:** *E. coli* (JM109) transformed with plasmid from A; 1: Chloramphenicol (30 µg); 2: Clindamycin (2 µg); 3: erythromycin (15 µg); 4: Gentamycin (10 µg); 5: Oxacillin (1 µg); 6: Vancomycin (30 µg); 7: Ampicillin (10 µg); 8: Cephalothin (30 µg). Resistance to 3 antibiotics (Cephalothin, ampicillin and oxacillin) got transferred from *Aeromonas* to *E. coli* (JM109). **179**

Figure 4.34. Mueller Hinton agar plates showing antibiotic resistance transfer from *Aeromonas* to *E. coli* through transformation; **A:** *Aeromonas*; **B:** *E. coli* (JM109) transformed with plasmid from A; 1: Streptomycin (10 µg); 2: Chloramphenicol (30 µg); 3: Ampicillin (10 µg); 4: Tetracycline (30 µg); 5: Getamycin (10 µg); 6: Kanamycin (30 µg); 7: Co-Trimoxazole (25 µg); 8: Amikacin (30 µg). Resistance to ampicillin got transferred from *Aeromonas* to *E. coli* (JM109). **180**

Figure 4.35. Mueller Hinton agar plates showing antibiotic resistance transfer from *Aeromonas* to *E. coli* through transformation; **A:** *Aeromonas*; **B:** *E. coli* (JM109) transformed with plasmid from A; 1: Cefepime (30 µg); 2: Cefpirome (30 µg); 3: Amikacin (30 µg); 4: Amoxyclav (30 µg); 5: Ceftazidime (30 µg); 6: Azithromycin (15 µg); 7: Ampicillin/Sulbactum (10/10 µg); 8: Cefoperazone/sulbactum **181**

(75/30 µg). Resistance to 2 antibiotics (Amoxyclav and ampicillin/sulbactam) got transferred from *Aeromonas* to *E. coli* (JM109).