

LIST OF SCHEMES

Scheme No.	Title	Page No.
Scheme I.1.	Synthesis of functionalized pyrroles in gluconic acid aqueous solution	5
Scheme I.2.	One-pot three-component synthesis of 2,4,6-triarylpyridines using triflimide	5
Scheme I.3.	Ruthenium catalyzed annulation of <i>N</i> -Cbz hydrazones <i>via</i> C–H/N–N bond activation for synthesis of isoquinolines	6
Scheme I.4.	Synthesis of selenopyridines in PEG-400 from aldehydes	6
Scheme I.5.	Copper-catalyzed aerobic synthesis of 2,4,5-trisubstituted imidazoles	7
Scheme I.6.	Copper-Catalyzed synthesis of substituted pyrazoles at room temperature	7
Scheme I.7.	Synthesis of 2 <i>H</i> -indazole derivatives catalysed by Cu ₂ O-Np	7
Scheme I.8.	Catalyst-free synthesis of benzodiazepines and benzimidazoles using glycerol	8
Scheme I.9.	Synthesis of unsymmetrical pyrazines from α -diazo oxime ethers	8
Scheme I.10.	Synthesis of quinoxaline from 2-iodo benzoic acid using organo Cu(II) catalyst	9
Scheme I.11.	Synthesis of 2,3-dihydroquinazolin-4(1 <i>H</i>)-one derivatives in ethanol	9
Scheme I.12.	Acidic ionic liquid [Bu ₃ NH][HSO ₄] catalysed synthesis of 1 <i>H</i> -pyrazolo[1,2- <i>b</i>]phthalazine-5,10-diones	10
Scheme I.13.	Synthesis of 1, 4-disubstituted-1,2,3-triazoles	10
Scheme I.14.	One-pot synthesis of 1,2,4-triazoles from hydrazides and secondary amides	10
Scheme I.15.	Copper (I)-catalyzed cascade addition-oxidative cyclization for synthesis of 1,2,4-triazoles	11
Scheme I.16.	PEG-600 as a reusable solvent system for synthesis of 1,3,5-triazines	11

Scheme I.17.	Synthesis of tetrazoles from nitrile using cuttlebone	12
Scheme I.18.	Synthesis of 1-substituted 1 <i>H</i> -1,2,3,4-tetrazole from amine and orthoformate	12
Scheme I.19.	Synthesis of 2,5-disubstituted oxazoles	12
Scheme I.20.	Tandem synthesis of isoxazoles from 1-copper (I) alkynes	13
Scheme I.21.	Synthesis of benzoxazoles in ethanol at room temperature	13
Scheme I.22.	One-pot synthesis of 3,5-disubstituted 1,2,4-oxadiazole using potassium fluoride	13
Scheme I.23.	I ₂ mediated synthesis of 1,3,4-oxadiazoles from aldehydes and hydrazides	14
Scheme I.24.	Synthesis of 2-amino-4 <i>H</i> -pyrans catalysed by WEMFSA	14
Scheme I.25.	PEG-600 mediated synthesis of 2-aminothiophene derivatives	14
Scheme I.26.	Synthesis of benzothiazoles using graphene oxide	15
Scheme I.27.	PhI catalyzed intramolecular oxidative coupling for synthesis of 2-amino-1,3,4-thiadiazoles	15
Scheme I.28.	One-pot synthesis of 3,5-disubstituted 1,3,4-thiadiazole from dithioesters	15
Scheme I.29.	Glycerol assisted one-pot pseudo five-component synthesis of 4 <i>H</i> -thiopyran	15
Scheme I.30.	Graphene oxide (GO): A carbocatalyst for the synthesis of 1,4-benzothiazines	16
Scheme I.31.	Synthesis of novel 2,5-diarylselenophenes using Woollin's reagent	16
Scheme I.32.	Copper catalyzed selenylation of imidazo[1,2- <i>a</i>]pyridines with selenium powder	17
Scheme I.33.	Copper promoted synthesis of 2-alkyl-1,2-benzisoselenazol-3(2 <i>H</i>)-ones	17
Scheme I.34.	Synthesis of 2-aryl-1,3-benzoselenazoles by reaction of bis(2-aminophenyl) diselenides with aldehydes	17
Scheme II.1.	Synthesis of tetrazole as proposed by Dimroth and Fester	22
Scheme II.2.	Synthesis of tetrazoles from nitrile using cuttlebone	24

Scheme II.3.	Cu(II) immobilized on Fe ₃ O ₄ @SiO ₂ @L-Arginine catalyst in the preparation of tetrazole	24
Scheme II.4.	A simple synthesis of tetrazole using amino acid	24
Scheme II.5.	Cu(OAc) ₂ in deep eutectic solvent (DES) for synthesis of tetrazole	25
Scheme II.6.	Bi(OTf) ₃ : A Lewis acid catalysed synthesis of tetrazole	25
Scheme II.7.	Synthesis of tetrazole derivatives using zeolite-based heterogeneous catalyst	25
Scheme II.8.	Cu-MCM-41 nanoparticles induced three-component synthesis of tetrazole	26
Scheme II.9.	Utilisation of dehydrating agent P ₂ O ₅ for development of tetrazole	26
Scheme II.10.	One-pot tandem reaction for synthesis of tetrazole moiety	26
Scheme II.11.	Synthesis of tetrazole by high molecular weight polymer	26
Scheme II.12.	Synthesis of tetrazole from oxime using Lewis acid catalyst	27
Scheme II.13.	A heterogeneous Pd-nanoparticle for synthesis of tetrazole from iodobenzene in presence of K ₄ [Fe(CN) ₆]	27
Scheme II.14.	Synthesis of 1-substituted 1 <i>H</i> -1,2,3,4-tetrazole from amine and orthoformate	27
Scheme II.15.	ONO pincer type Pd(II) complex: A heterogenous catalyst for synthesis of tetrazole from phenyl boronic acid	28
Scheme II.16.	Synthesis of tetrazole derivative using ionic liquid azide	28
Scheme II.17.	Cyanogens azide: A source of azide for synthesis of 5-amino tetrazoles	28
Scheme II.18.	CAS as a catalyst in organic synthesis	29
Scheme II.19.	Synthesis of substituted benzothiazoles by using nano-ceria	30
Scheme II.20.	CAN: A Lewis acid catalyst for synthesis of unsymmetrical bis(indolyl)alkane	30
Scheme II.21.	One pot three-component synthesis of 5-substituted 1 <i>H</i> -tetrazole from aldehyde	31
Scheme II.22.	Plausible mechanism for the synthesis of 5-substituted 1 <i>H</i> -tetrazoles	37
Scheme III.1.	Synthesis of nitrile using Sandmeyer reaction	50

Scheme III.2.	Rosenmund-von Braun reaction	51
Scheme III.3.	S. L. Buchwald <i>et al.</i> method for synthesis of aryl nitrile	51
Scheme III.4.	Ammoxidation reaction for the synthesis of acrylonitrile	51
Scheme III.5.	Friedel-Crafts synthesis of benzonitrile	51
Scheme III.6.	Houben-Fischer synthesis of aromatic nitrile	52
Scheme III.7.	Synthesis of nitrile using $K_4[Fe(CN)_6]$ as cyanide source in presence of biosynthesized Ag NPs	53
Scheme III.8.	Synthesis of nitrile from aldehyde using hydroxylamine- <i>O</i> -sulfonic acid	53
Scheme III.9.	Synthesis of nitrile from aldehyde in DES	54
Scheme III.10.	Catalyst-free synthesis of nitrile in glycerol	54
Scheme III.11.	Synthesis of nitrile utilizing acetohydroxamic acid	54
Scheme III.12.	Transformation of aldehyde into nitriles using $NH_3.H_2O/FeCl_2/NaI/Na_2S_2O_8$	54
Scheme III.13.	Transoximation reaction for synthesis of nitrile utilizing <i>O</i> -protected oxime	55
Scheme III.14.	Synthesis of nitrile assisted by $HCl.DMPU$	55
Scheme III.15.	Ionic liquid catalysed synthesis of nitrile	55
Scheme III.16.	Synthesis of nitrile in presence of Fe_3O_4 -CTAB NPs	56
Scheme III.17.	Synthesis of nitriles <i>via</i> Schmidt reaction using triflic acid	56
Scheme III.18.	Synthesis of nitrile in ionic liquid <i>via</i> Schmidt type protocol	57
Scheme III.19.	$POCl_3$ mediated synthesis of nitrile using sodium azide	57
Scheme III.20.	Synthesis involving transformation of aldoxime into nitrile	57
Scheme III.21.	Catalytic Appel-type reaction for synthesis of nitrile from amide	58
Scheme III.22.	Oxidative conversion of alcohols into nitriles	58
Scheme III.23.	Transnitration for conversion of acid into nitrile	58
Scheme III.24.	Synthesis of nitriles <i>via</i> cyanation of aryl iodides	59
Scheme III.25.	Synthesis of nitriles from phenyl boronic acid using various thiocyanate derivatives	59
Scheme III.26.	<i>p</i> -TsOH mediated 1,3-dipolar cyclo-addition for the synthesis	60

	of 1,2,3-triazoles	
Scheme III.27.	<i>p</i> -TsOH catalysed synthesis of substituted 2,4-diaryl-3-sulfonyl quinolines under microwave condition	61
Scheme III.28.	Silica mediated synthesis of indolinoxazolidine	61
Scheme III.29.	Synthesis of bis(indolyl)methanes using silica as surface	61
Scheme III.30.	<i>p</i> -TsOH mediated synthesis of nitrile from aldehyde	62
Scheme III.31.	Plausible mechanism for transformation of aldehydes into nitriles	66
Scheme IV.1.	PEG-400: Metal-free synthesis of 2-amino-3-cyanopyridine in water	76
Scheme IV.2.	HBF ₄ : An oxidizing catalyst for synthesis of pyridine derivatives	77
Scheme IV.3.	Synthesis of pyridine derivative in presence of heterogeneous catalyst FePO ₄	77
Scheme IV.4.	NMS: A protic ionic liquid for synthesis of 2-aminonicotino nitriles	77
Scheme IV.5.	[Yb(PFO) ₃] promoted synthesis of 2-amino-3-cyanopyridine	78
Scheme IV.6.	Trifluoroethanol mediated synthesis of 2-aminonicotinonitriles	78
Scheme IV.7.	Ultrasound-promoted method for designing nicotinonitriles in water	78
Scheme IV.8.	Cellulose sulfuric acid: An efficient catalyst for preparation of 2-amino-3-cyanopyridine in water	79
Scheme IV.9.	Nano Fe ₃ O ₄ catalysed synthesis of 2-amino-3-cyanopyridine	79
Scheme IV.10.	Graphene oxide: A carbocatalyst mediated synthesis of 2-amino nicotinonitriles in water	79
Scheme IV.11.	β-Cyclodextrin catalysed synthesis of thioamide derivatives	80
Scheme IV.12.	One-pot synthesis of 2-amino-4 <i>H</i> -chromen-4-yl phosphonate derivatives using β-cyclodextrin in water	81
Scheme IV.13.	Water mediated synthesis of 2,3-dihydroquinazolin-4(1 <i>H</i>)-one	81
Scheme IV.14.	Sulfonated graphene oxide catalyzed synthesis of pyrano[2,3- <i>c</i>] pyrazole in water	82
Scheme IV.15.	One-pot four-component synthesis of 2-amino-4,6-diphenyl nicotinonitriles	82

Scheme IV.16.	Plausible mechanism for formation of pyridine motif	86
Scheme V.A.1.	Fe ₃ O ₄ /SiO ₂ /PTS/AMTA nanocomposite: A heterogeneous catalyst for synthesis of bis(indolyl)methane derivatives	96
Scheme V.A.2.	Oleic acid catalysed synthesis of bis(indolyl)methane derivatives	96
Scheme V.A.3.	Synthesis of trisubstituted methanes catalyzed by sulfated polyborate	97
Scheme V.A.4.	Enzymatic approach to cascade synthesis of bis(indolyl)methanes	97
Scheme V.A.5.	Graphene oxide: A recyclable solid acid catalysed synthesis of bis(indolyl)methanes	97
Scheme V.A.6.	Thiamine hydrochloride as a recyclable organocatalyst for the synthesis of bis(indolyl)methanes	98
Scheme V.A.7.	Synthesis of bis(indolyl)methanes in aqueous ethyl lactate	98
Scheme V.A.8.	Meglumin assisted synthesis of bis(indolyl)methanes	98
Scheme V.A.9.	Synthesis of bis(indolyl)methanes using humic acid	99
Scheme V.A.10.	Possible mechanism for bis(indolyl)methanes assisted by humic acid	103
Scheme V.B.1.	PEG-SO ₃ H: A homogeneous polymeric catalyst for the synthesis of 4,4'-(arylmethylene)-bis(3-methyl-1-phenyl-1 <i>H</i> pyrazol-5-ol)	116
Scheme V.B.2.	Phosphomolybdic acid catalysed synthesis of bis(pyrazolyl) methanes	116
Scheme V.B.3.	Caffeine-H ₃ PO ₄ promoted synthesis of 4,4'-(arylmethylene) bis(1 <i>H</i> pyrazol-5-ols)	116
Scheme V.B.4.	Synthesis of bis(pyrazolyl)methanes utilizing Zirconium@ guanine@MCM-41 nanoparticles	117
Scheme V.B.5.	SDBS assisted reaction of aromatic aldehydes with 1-phenyl -3-methyl-5-pyrazolone in aqueous media	117
Scheme V.B.6.	Sulfonated carbon/nano-metal oxide composites: A recyclable solid acid catalyst for synthesis of bis(pyrazolyl)methanes	118
Scheme V.B.7.	Application of {Fe ₃ O ₄ @SiO ₂ @(CH ₂) ₃ -thiourea dioxide-SO ₃ H/HCl} catalyst for synthesis of bis(pyrazolyl)methanes	118
Scheme V.B.8.	Aqueous extract of <i>Balanites roxburghii</i> fruit promoted synthesis	119

	of bis(pyrazolyl)methanes	
Scheme V.B.9.	Synthesis of tetrazoles by humic acid	119
Scheme V.B.10.	[Bu ₃ NH][HSO ₄] catalysed synthesis of 1 <i>H</i> -pyrazolo[1,2- <i>b</i>]phthalazine-5,10-diones	120
Scheme V.B.11.	Synthesis of 1,4-benzothiazines using graphene oxide	121
Scheme V.B.12.	Solvent-free synthesis of bis(pyrazolyl)methanes using humic acid	121
Scheme V.B.13.	Plausible mechanism for synthesis of bis(pyrazolyl)methanes	124
Scheme VI.A.1.	Synthesis of thioamide derivatives from acid using ammonium phosphorodithioates	135
Scheme VI.A.2.	Thionation of amides using a solid-supported P ₂ S ₅ reagent	135
Scheme VI.A.3.	Photocatalyst promoted synthesis of thioamides from aliphatic amine	135
Scheme VI.A.4.	[DBUH][OAc]: An ionic liquid catalysed solvent-free synthesis of thioamides from nitriles	136
Scheme VI.A.5.	Synthesis of thioamide derivatives from thiols	136
Scheme VI.A.6.	Chemoselective one-pot transformation of aldoximes	136
Scheme VI.A.7.	Oxidative coupling of disulphide to design thioamide	136
Scheme VI.A.8.	Microwave-assisted Kindler thioamide synthesis	137
Scheme VI.A.9.	Willgerodt-Kindler reaction in presence of catalytic amount of Na ₂ S.9H ₂ O	137
Scheme VI.A.10.	Sulfonic acid functionalized nano γ -Al ₂ O ₃ mediated synthesis of thioamides	138
Scheme VI.A.11.	Sulfated polyborate promoted synthesis of thioamides	138
Scheme VI.A.12.	β -Cyclodextrin catalysed Kindler type reaction	138
Scheme VI.A.13.	Water mediated synthesis of thioamides	138
Scheme VI.A.14.	Synthesis of thioamides from aldehydes and amines	139
Scheme VI.A.15.	Plausible mechanism for synthesis of thioamide derivatives	143
Scheme VI.B.1.	Synthesis of thiopyran derivatives using piperidine	155
Scheme VI.B.2.	Microwave irradiation for synthesis of thiopyran derivatives	156
Scheme VI.B.3.	Ionic liquid promoted synthesis of thiopyrans	156

Scheme VI.B.4.	L-proline: A green organocatalyst for synthesis of thiopyrans	156
Scheme VI.B.5.	Chitosan-grafted-poly(4-vinylpyridine) catalysed synthesis of thiopyran derivatives	157
Scheme VI.B.6.	One-pot pseudo five-component synthesis of thiopyran derivatives	157
Scheme VI.B.7.	MNPs-BPAT promoted synthesis of thiopyran derivatives	158
Scheme VI.B.8.	Synthesis of thiopyran derivatives through electrolysis	158
Scheme VI.B.9.	Glycerol mediated synthesis of 4,4'-(arylmethylene)bis(3-methyl-1 <i>H</i> -pyrazol-5-ols) and dihydroquinazolin-4(1 <i>H</i>)-ones	159
Scheme VI.B.10.	Catalyst-free synthesis of 4-(aryl)-3-(phenylsulfonyl)-4 <i>H</i> -benzo [<i>h</i>]chromen-2-amine derivatives using glycerol	160
Scheme VI.B.11.	One-pot multi-component synthesis of 4 <i>H</i> -thiopyran	160
Scheme VI.B.12.	Plausible mechanism for synthesis of thiopyran derivatives	163