

TABLE OF CONTENTS

	Page No.
Abstract (4 pages)	i
Preface (1 page)	v
<hr/>	
List of Tables (1 page)	xiii
List of Schemes (8 pages)	xv
List of Figures (2 pages)	xxiii
List of Appendices (1 page)	xxv
Appendix A: List of Research Publications (2 page)	xxvii
Appendix B: Poster & Oral Presentations (1 page)	xxix
Abbreviation (4 pages)	xxxi

Chapter I

A brief review on approaches towards the synthesis of bioactive heterocyclic compounds

I.1. Introduction	3
I.2. Approach towards the synthesis of heterocyclic compounds	5
I.2.1. Synthesis of <i>N</i> -containing heterocyclic compounds	5
I.2.2. Synthesis of <i>O</i> -containing heterocyclic compounds	12
I.2.3. Synthesis of <i>S</i> -containing heterocyclic compounds	14
I.2.4. Synthesis of <i>Se</i> -containing heterocyclic compounds	16
I.3. Conclusion	18
I.4. References	18

Chapter II

One-pot three-component synthesis of 5-substituted 1*H*-tetrazoles from aldehydes

II.1. Introduction	21
II.2. Background and Objectives	22
II.2.1. Classical method for synthesis of tetrazole	22

II.2.2. Modern methods for synthesis of 5-substituted 1 <i>H</i> -tetrazole	22
II.2.2.1. Synthesis of tetrazole derivatives from nitriles	23
II.2.2.2. Synthesis of tetrazole derivatives from aldehydes <i>via</i> one-pot three-component protocol	24
II.2.2.3. Synthesis of tetrazole moiety from various synthons	27
II.2.3. Role of various cerium components as catalyst in organic synthesis	28
II.3. Present work: Result and Discussion	30
II.3.1. Optimization of the reaction conditions	31
II.3.2. Recyclability of catalyst	35
II.3.3. Scale-up experiment	36
II.3.4. Mechanism	36
II.4. Conclusion	37
II.5. Experimental Section	38
II.5.1. General Information	38
II.5.2. General procedure for the synthesis of tetrazole from aldehyde	38
II.5.3. Characterization data of various tetrazole derivatives	38
II.5.4. Scanned copies of ¹ H and ¹³ C NMR, HRMS spectra	43
II.6. References	45

Chapter III

***p*-TsOH assisted solvent and metal catalyst-free synthesis of nitriles from aldehydes *via* Schmidt reaction**

III.1. Introduction	49
III.2. Background and Objectives	50
III.2.1. Synthesis of nitrile using classical methods	50
III.2.2. Modern methods for synthesis of nitrile	52
III.2.2.1. Synthesis of nitriles from aldehydes	53
III.2.2.2. Synthesis of nitriles utilizing Schmidt reaction from aldehydes	56
III.2.2.3. Synthesis of nitrile from various substrates	57
III.2.3. Organic synthesis catalyzed by <i>p</i> -toluene sulphonic acid	59
III.2.4. Silica gel supported organic synthesis	61

III.3. Present Work: Result and Discussion	62
III.3.1. Optimization of the reaction conditions	62
III.3.2. Mechanism	65
III.4. Conclusion	66
III.5. Experimental Section	66
III.5.1. General Information	66
III.5.2. General Procedure for the synthesis of nitriles from aldehydes	66
III.5.3. Characterization data of various nitrile derivatives	67
III.5.4. Scanned copies of IR, ¹ H and ¹³ C NMR spectra of benzonitrile	70
III.6. References	72

Chapter IV

β-Cyclodextrin: A supramolecular catalyst for metal-free approach towards the synthesis of 2-amino-4,6-diphenylnicotinonitriles

IV.1. Introduction	75
IV.2. Background and Objectives	76
IV.2.1. One-pot four-component synthesis of 2-amino-3-cyanopyridine	76
IV.2.2. β-Cyclodextrin mediated organic synthesis	80
IV.2.3. Water as green reaction medium	81
IV.3. Present work: Result and Discussion	82
IV.3.1. Optimization of the reaction conditions	82
IV.3.2. Mechanism	85
IV.4. Conclusion	86
IV.5. Experimental Section	87
IV.5.1. General Information	87
IV.5.2. General procedure for the synthesis of 2-amino-4,6-disubstituted nicotinonitriles	87
IV.5.3. Characterization data of various nicotinonitrile derivatives	87
IV.5.4. Scanned copies of ¹ H and ¹³ C NMR spectra of 2'-amino-6'-phenyl-[3,4'-bipyridine]-3'-carbonitrile	91

Chapter V

Section A

Humic acid: An organocatalyst for solvent-free benign synthesis of bis(indolyl)methane

V.A.1. Introduction	95
V.A.2. Background and Objectives	95
V.A.2.1. Synthesis of bis(indolyl)methanes from aldehydes	95
V.A.3. Present Work: Result and Discussion	98
V.A.3.1. Optimisation of reaction conditions	99
V.A.3.2. Mechanism	102
V.A.3.3. Recyclability experiment	103
V.A.4. Conclusion	104
V.A.5. Experimental Section	104
V.A.5.1. General Information	104
V.A.5.2. General procedure for the synthesis of bis(indolyl)methanes	104
V.A.5.3. Characterization data of various bis(indolyl)methane derivatives	105
V.A.5.4. Scanned copies of ^1H and ^{13}C NMR spectra of 3-((1 <i>H</i> -indol-3-yl)(phenyl)methyl)-1 <i>H</i> -indole	110
V.A.6. References	111

Section B

Humic acid mediated solvent-free synthesis of bis(pyrazolyl)methane

V.B.1. Introduction	115
V.B.2. Background and Objectives	115
V.B.2.1. Synthesis of bis(pyrazolyl)methanes from aldehydes	115
V.B.2.2. Application of humic acid in organic synthesis	119
V.B.2.3. Solvent-free protocols in organic synthesis	120
V.B.3. Present Work: Result and Discussion	121
V.B.3.1. Optimisation of reaction conditions	121

V.B.3.2. Mechanism	124
V.B.4. Conclusion	125
V.B.5. Experimental Section	125
V.B.5.1. General Information	125
V.B.5.2. General procedure for the synthesis of bis(pyrazolyl)methanes	125
V.B.5.3. Characterization data of various bis(pyrazolyl)methane derivatives	126
V.B.5.4. Scanned copies of ¹ H and ¹³ C NMR spectra of 4,4'-[(4-methoxyphenyl) methylene]bis(3-methyl-1-phenyl-1 <i>H</i> -pyrazol-5-ol)	129
V.B.6. References	130

Chapter VI

Section A

Glycerol: A benign solvent for catalyst-free synthesis of thioamides from aldehydes

VI.A.1. Introduction	133
VI.A.2. Background and Objectives	134
VI.A.2.1. Synthesis of thioamide derivatives from various synthons	134
VI.A.2.2. Willgerodt-Kindler Reaction for synthesis of thioamide derivatives	136
VI.A.3. Present Work: Result and Discussion	139
VI.A.3.1. Optimisation of reaction conditions	139
VI.A.3.2. Mechanism	143
VI.A.4. Conclusion	144
VI.A.5. Experimental Section	144
VI.A.5.1. General Information	144
VI.A.5.2. General procedure for the synthesis of thioamide derivatives	144
VI.A.5.3. Characterization data of various thioamide derivatives	144
VI.A.5.4. Scanned copies of ¹ H and ¹³ C NMR spectra of (4-fluorophenyl) (pyrrolidin-1-yl)methanethione	151
VI.A.6. References	152

Section B

Glycerol assisted catalyst-free one-pot multi-component synthesis of thiopyran from aldehyde

VI.B.1. Introduction	155
VI.B.2. Background and Objectives	155
VI.B.2.1. Synthesis of thiopyran derivative from active methylene compound	155
VI.B.2.2. Synthesis of thiopyran <i>via</i> one-pot pseudo five-component method	157
VI.B.2.3. Glycerol mediated organic synthesis	158
VI.B.3. Present Work: Result and Discussion	160
VI.B.3.1. Optimisation of reaction conditions	160
VI.B.3.2. Mechanism	163
VI.B.4. Conclusion	164
VI.B.5. Experimental Section	164
VI.B.5.1. General Information	164
VI.B.5.2. General procedure for the synthesis of 4 <i>H</i> -thiopyran	164
VI.B.5.3. Characterization data of various thiopyran derivatives	164
VI.B.5.4. Scanned copies of ¹ H and ¹³ C NMR spectra of 2,6-diamino-4-phenyl-4 <i>H</i> -thiopyran-3,5-dicarbonitrile	167
VI.B.6. References	168
Bibliographic References	
References for Chapter I	169
References for Chapter II	170
References for Chapter III	173
References for Chapter IV	175
References for Chapter V A	177
References for Chapter V B	179
References for Chapter VI A	181
References for Chapter VI B	182
Index	185